


ENDOCRINE BLOCK SIXTH WEEK



Done by: Noor Alzahrani , Norah alajmi , Atheer Alduaij

Design by: khalid alosaimi



Table of Content

Biochemistry	3
Medicine	4
psychiatry	
Pharmacology	





[9]Obesity

q1) which of the following hormones will be Suppressed in starvation?

- A. LEPTIN
- B. RESISTEN
- C. GHRELIN
- D. ADEPONECTIN

Ans:A

Q2) which of the following hormones will stimulate the appetite?

- A. ADEPONECTIN
- B. LEPTIN
- C. RESISTEN
- D. GHRELIN

ANS:D

Q3) which of the following hormones will decrease in dieting?

- A. ADEPONECTIN
- B. LEPTIN
- C. RESISTEN
- D. GHRELIN

Ans:B

[10] Metabolic syndrome

Q1)Thiazolidinediones (TZDs) is used to treat which one of the following?

A-INSULIN RESISTANCE AND TYPE2 DM

B-REDUCE LIPID LEVEL

C- HYPERTENSION

D-HYPERLIPIDAEMIA

ANS:A

Q2) which of the following drug can be risk factor of metabolic syndrome ?-

AMETFORMIN.

B-RIFAMPICIN.

C-STATINS

D-AMIODARONE

ANS:B

Q3)insulin resistance in obese leads to?

A- INHIBIT HORMONE SENSITIVE LIPASE

B- DECREASE VLDL

C-INCREASE HDL-D-INCREASE LEVEL OF FATTY ACID

ANS:D





Medicine

obesity

Q1: Which one of the following is true regarding BMI & fat?

- A. At same body mass index men tend to have more fat than women
- B. At same body mass index women tend to have more fat than men
- C. At same body mass index young adult tend to have more fat than elderly
- D. Athlete has high BMI because of increase fat

Answer B

Q2:Which one of the following is true regarding BMI?

- A. It is the the square of height divided by the Wight
- B. Obesity in childhood increase risk of diabetes in adulthood
- C. If the BMI=> 25 it is consider obesity

Answer B

Q3:Why obesity accelerating in developing country?

- A. Increase consumption of energy dense nutrient poor food
- B. Reduce physical activity
- C. A & B

Answer C

Q4:causes of obesity?

- A. necessarily caused by over eating
- B. biochemical difference
- C. basal body temp
- D. B & C

Answer D

Not necessarily caused by over eating

Q5:genetic role in obesity?

- A. less quantity and sensitivity to satiety hormone
- B. increase leptin production, and sensitivity to satiety

answer A

Q6:which of the following is true?

- A. men tend to have visceral obesity
- B. waist hip ratio more than 1 in men means peripheral obesity
- C. waist hip ratio less than 0.8 in women means visceral obesity

Revision Test Team

answer A

in men (>1 visceral obesity ,,<1peripheral obesity) in women (>0.8 visceral obesity ,, <0.8 peripheral obesity)

Q7:loss of 10 kg in 100kg patient with obesity related co morbidity?

- A. decrease mortality (diabetes and obesity) related deaths
- B. decrease blood pressure
- C. decrease LDL, cholesterol, triglycerides
- D. increase HDL
- E. All above

answer E



psychiatry



www.shutterstock.com - 114848119

Q1:treatment goal in diabetes?

- A. Prevent Short and long term complications
- B. Symptom free
- C. Quality of life
- D. All above

Answer D

Q2:What is the most important factor &preventive approach of the psychological difficulty in case of diabetes?

- A. Peers & media
- B. Health care team
- C. Educative supportive Family

Answer C

Q3:Which of the following pattern in Yong adult that make difficult follow up?

- A. Confidentiality
- B. trust
- C. Possess skills to independently ask physician
- D. Inconsistent behavior

Answer D

Q4:The best clinic for emerging adult is?

- A. Adult clinic
- B. Children clinic
- C. Transitional clinic

answer C

Q5: Which is the most important program for transitional population which is expected to influence HbA1c level?

- A. Training to Fluently counting carbohydrate every time they eat
- B. Chang their inappropriate behavior such as uncertainty
- C. Improve life style

Answer A





Pharmacology

- 1) which one of the following drugs is better to be given for a patient with hypercholestremia?
- A- Tolbutamide
- **B-** Metformin
- C- Acarbosr

Ans:"B"

- 2) which one of the following drugs is better to be given for Impaired glucose tolerance ?
- A- Tolbutamide
- **B- Metformin**
- C- Acarbosr

Ans:"C"

- 3) which one of the following drugs should be avoided for a patient with C.H.F "Congestive Heart Failure"?
- A- Pioglitazone
- B- Repaglitnide
- C- Acarbosr

Ans: "A"

- 4) Patient with diabetes comes to complain about diarrhea and flutalence. which one of the following drugs has been prescribed for him?
- A- Pioglitazone
- B- Repaglitnide
- C- Acarbosr

Ans:"C"

* * * *

- 5) which one of the following drugs can induce Hypoglycemia?
- A- Glipizide
- B- Sitagliptan
- C- Metformin

Ans:"A"

* * * *

6) which one of the following drugs can induce Hepatotoxicity?



A- Glipizide

B- Sitagliptan

C- Pioglitazone

Ans: "C"

* * * *

7) which one of the following sulfanourease can be given safely to renal patient and would not induce Hypoglycemia?

A- Glipizide

B- Tolbutamide

C- Glyburide

Ans:"B"

* * * *

8) which one of the following drugs acts via PPAR-gamma receptors?

A- Glitazones

B- DPP-4-Inhibitors

C- alpha-Glucosidase-Inhibitors

Ans: "A"

If you have any questions you want to add, please send it to

Revisiontest432@gmail.com

Good luck

Team Leaders : Khalid Al-Osaimi& Lulu Al-Obaid

