

Hormones affecting the breast

Dr. Aida Korish


Objectives

- Hormonal requirement for breast development (Mamogenesis)
- Hormones involved in the process of lactation (Lactogenesis) and their physiological action
- Physiological basis of suckling reflex and its role in lactation
- Galactopoeisis
- Involution (the termination of milk production).






Where does milk come from?










thelial cells secrete milk into the lumen. Contraction of the surrounding myoepithelial cells ejects the secreted milk out through the duct.

- During puberty
 - Estrogen stimulate proliferation of ducts and deposition of fat
 - Progesterone stimulate development of lobules






- During pregnancy
 - Complete development of glandular tissue



Copyright © 2010 by Saunders, an imprint of Elsevier Inc. All rights reserved

- Endocrine system plays a major role in synchronizing development (mamogenesis) and function (lactogenesis) of mammary gland with reproduction
- Three categories of hormones:
 - Reproductive hormones (endocrine)
 - Estrogen, progesterone, prolactin, oxytocin and hPL
 - Metabolic hormones (endocrine)
 - GH, corticosteroids, thyroxin, PTH and insulin
 - Mammary hormones (autocrine)
 - GH, prolactin, parathyroid hormone-related protein (PTHrP) and leptin

- Reproductive hormones (direct effect)
 - Estrogen (placenta)
 - Growth & branching of ductal system (with GH)
 - Fat deposition in the stroma
 - Progesterone (placenta)
 - Growth of lobule-alveolar system(budding of alveoli and secretory changes in epithelial cells)
- Although estrogen and progesterone are essential for physical development of the breasts, they inhibit actual secretion of milk



PREGNANT

• **Prolactin** (anterior pituitary)

- Its level increases during pregnancy (10-20 times)
- Its main function is milk production
- Sudden drop in E & P after delivery allows milk production
- It is controlled mainly by hypothalamic hormone
 - PIH (Dopamine)
- Human placental lactogen (placenta)
 - Facilitate mammogenesis
 - Delay milk production

Lactogenesis


- Lactogenesis: cellular changes by which mammary epithelial cells are converted from a non secretory state to a secretory state: 2 stages
- Lactogenesis 1: (Cytologic and enzymatic differentiation of alveolar epithelial cells).
 - Starts in mid pregnancy and characterized by expression of many genes involved in synthesis of milk components (increase in the uptake transport systems for amino acids, glucose, and calcium required for milk synthesis).
 - Hormones involved:
 - 1. Progesterone (suppresses milk secretion)
 - 2. **Prolactin** and/or placental lactogen
 - 3. Growth hormone
 - 4. Glucocorticoids (**Cortisol**)

Lactogenesis

- Lactogenesis 2: (Copious secretion of all milk components).
 - Around parturition withdrawal of progesterone + high level of prolactin leads to:
 - Further increase in expression of milk protein genes
 - Glands absorb increased quantities of metabolic substrates from the blood.
 - Movement of cytoplasmic lipid droplets and casein into alveolar Lumina
 - Transfer of immunoglobulins
 - Secretion of colostrum followed by milk


Lactogenesis

- Lactogenesis 2:
 - Suckling stimulates further increase in expression of genes involved in milk secretion with expansion of alveolar epithelium
 - Lactation is maintained by removal of milk
 - 2 hormones involved
 - Prolactin (milk production)
 - Oxytocin (milk let-down)



Alveolus of Mammary Gland

Stages of lactogenesis



Mamogenesis

Lactogenesis

and

Galctopoesis

Hormonal regulation of lactogenesis

- Metabolic hormones (direct effect)
 - GH
 - Can be produced locally
 - Its secretion is stimulated by progesterone
 - Increases production of IGF-1 by the liver
 - Mediate cell survival and ductal growth
 - Corticosteroids
 - Increases during pregnancy (five fold)
 - Involved in breast development (permissive action on milk protein synthesis)


Hormonal regulation of lactogenesis

Thyroxin


- Essential for milk production
- Thyroxin & TSH level decreases during lactation
- TRH increases leading to stimulation of PRL (nasal administration to treat inadequate lactation)
- Insulin
 - Low during lactation
 - Shunt of nutrients from storage depots to milk synthesis

Hormonal regulation of lactogenesis

- Mammary hormones
 - GH
 - Progesterone stimulates its secretion
 - Leptin
 - Increases during pregnancy (increase adipose tissue)
 - Decreases with lactation
 - PTHrP
 - Increases during lactation
 - Mobilizes bone calcium
 - Increase in alkaline phosphatase




Copyright © 2009 by Saunders, an imprint of Elsevier Inc. All rights reserved.




The pathways for milk secretion and synthesis by the mammary epithelial cell. I: Exocytosis of milk protein, lactose, and other components of the aqueous phase in Golgi-derived secretory vesicles. II: Milk fat secretion via the milk fat globule. III: Direct movement of mono valent ions, water, and glucose across the apical membrane of the cell. IV: Transcytosis of components of the interstitial space. V: The paracellular pathway for plasma components and leukocytes. Pathway V is open only during pregnancy, involution, and in inflammatory states such as mastitis. SV = Secretory vesicle; RER = Rough endoplasmic reticulum; BM = Basement membrane; MFG = Milk fat globule; CLD = Cytoplasmic lipid droplet; N = Nucleus; PC = Plasma cell; FDA = Fat-depleted adipocyte; TJ = Tight junction; GJ = Gap junction; D = Desmosome; ME = Myoepithelial cell.


Galactopoeisis

- **Definition:** Galactopoeisis is defined as the maintenance of lactation once lactation has been established.
- Role of Hormones
 - **Prolactin**: milking-induced surge is a direct link between the act of nursing (or milk removal) and the galactopoeitic hormones involved in maintaining lactation.
 - **Growth Hormone:** support increase in synthesis of lactose, protein, and fat in the mammary gland
 - **Glucocorticoids:** galactopoeitic in physiological doses
 - Thyroid Hormones: galactopoeitic
 - Ovarian Hormones :
 - Estrogen in very low doses is galactopoietic
 - Progesterone alone has no effect on galactopoeisis because there are no progesterone receptors in the mammary gland during lactation





Suckling reflex




• Milk production is a

"use it or lose it" process. The more often and effectively the baby nurses, the more milk will be produced

• Milk production <100 ml/day in day 1 postpartum

- Milk production by day 3 reaches 500 ml/day
- Milk composition changes dramatically(↓Na⁺²& Cl⁻) due to closure of tight junctions that block paracellular pathway



- يقول تعالى (والوالدات يرضعن أو لادهن حولين كاملين لمن أراد أن يُتمَ
 الرضاعة) [البقرة :٢٣٣] .
- وفي إشارة علمية دقيقة أخرى للقرآن الكريم نراه يحدد مدة الرضاعة بما يقرب من الحولين ، كما جاء في الآية رقم (١٤) في سورة [لقمان]:
 (ووصينا الإنسان بوالديه حملته أمه وهنا على وهن وفصاله في عامين)
 ، والآية (١٥) في سورة [الأحقاق] : (حملته أمه كرها ووضعته كرها وحمله وفصاله ثلاثون شهرا) ، ويُفهم من هذا أن إرضاع الحولين ليس حتماً ، بل هو التمام ، ويجوز الاقتصار على ما دونه ، كما أشارت الأحكام وحماء أرادا إرضاعة بما وحمله وفصاله في عامين)

AAP Recommendations

- Exclusive breast feeding for the first six months of life
- Continued breast feeding for at least one year, 'As long as is desired by mother and child'.



American Academy of Pediatrics (2005). "Breastfeeding and the Use of Human Milk." Pediatrics 115(2): 496-506.

Failure of breast feeding

