




Physiology team

# Lecture 2

## Physiology of androgens and control of male sexual functions



### Sources:

- ✓ Female slides
- ✓ Male slides

# Objectives

1. Understand the functions of the male reproductive organs and glands
2. Describe the synthesis, secretion, metabolism and effects of testosterone
3. Explain how the hypothalamus and anterior pituitary gland regulate male reproductive function
4. Describe the major testicular abnormalities
5. Discuss the normal mechanism of the male sexual act

# Sperms

## ◆ Spermatogenesis

- formation of sperm & occur in the **seminiferous tubules** during active sexual life due to stimulation by **AP-Gonadotropic Hormones**, begin at **age of 13 years**, continue throughout life & decrease in old age.

## ◆ Maturation of sperm in the epididymis



- After formation in the seminiferous tubules (takes 74 days), the sperm require several days to pass through the **epididymis (still non-motile)**.
- - After the sperm have been in the epididymis for some 18 to 24 hours, they **develop the capability of motility** (some inhibitory proteins in the epididymal fluid prevent final motility until after ejaculation).

## ◆ Physiology of mature sperm

- . Mature sperm are **motile** & **capable of fertilizing the ovum** & their activity is enhanced in a neutral & slightly **alkaline** medium & **depressed in mildly acidic medium**. The life expectancy of ejaculated sperm in the female genital tract is **only 1 to 2 days**.

## ◆ Storage of sperm

- The 2 testis of adult human formed up to **120 million** sperm each day.
  1. **Small amount** stored in the epididymis
  2. **the majority stored in the vas deferens**, maintaining their fertility for at least a month. **The sperm are kept inactive state by multiple inhibitory substances in the secretion of the ducts.**



# Cont.

## ◆ After ejaculation

- The sperm become motile & capable of fertilizing the ovum **called maturation.**
- The Sertoli cells and epithelium of the epididymis secrete nutrient fluid which contains (testosterone & estrogens), enzymes & nutrients essential for sperm maturation.

| Sertoli Cells   | Leydig Cell   |
|---|---|
| large with overflowing cytoplasmic envelopes that <u>surround the developing spermatogonia</u> around the central lumen of the seminiferous tubules | <ul style="list-style-type: none"><li>• <b><u>lie with interstitium between the seminiferous tubules.</u></b></li><li>• They are <u>non-existent</u> in the testis during childhood when the testis secrete almost no testosterone,</li><li>• Numerous in the newborn male infants for the first few months of life</li><li>• Active at puberty &amp; throughout adult life &amp; secrete testosterone.</li></ul> |



|           | seminal vesicles   | prostate gland  |
|-----------|--|---|
| Secretion | Muroid material containing fructose, citric acid & nutrient substances & <u>large quantities of prostaglandins</u> & fibrinogen  | Thin milky fluid contains Ca <sup>2+</sup> ion, citrate ion, phosphate ion, a clotting enzyme & profibrinolysin.  |
| Functions | <p>The <u>prostaglandins are important help in fertilization:</u></p> <p>1- by reacting with the female cervical mucus making <u>it more receptive</u> to sperm movement.</p> <p>2- by causing <u>backward reverse peristaltic contractions</u> of the uterus &amp; fallopian tubes to move the ejaculated sperm toward the ovaries. There is receptors for prostaglandins that will cause contraction of the smooth muscles and help the movement of the sperms when prostaglandins bind to it.</p> | <p>The alkaline prostatic fluid is important for</p> <ol style="list-style-type: none"> <li><u>successful fertilization of the ovum.</u></li> <li>Help to <u>neutralize</u> the slightly acidic fluid of the vas deferens (due to the presence of citric acid and metabolic &amp; product of the sperm inhibit sperm fertility)</li> <li>helps to neutralize <u>the acidic vaginal secretions</u> (pH 3.5-4.0) to optimize it for better sperm motility (pH 6.0-6.5)</li> </ol> |

- The glands are important .

## Semen

- Ejaculated semen during sexual act is composed of the fluid & sperm from the vas deferens (~10%)+fluid from the prostate gland (~30%),+fluid from the seminal vesicles (~60%) + small amounts from the mucous glands the bulbourethral glands.
- The average pH is about 7.5, the alkaline prostatic fluid help to neutralize the mild acidity of other portions of the semen & gives the semen a milky appearance
- fluid from the seminal vesicles & mucous glands give the semen muroid consistency.

# Capacitation of the spermatozoa:

Making it possible for them to penetrate the ovum

Sperm in the epididymis is kept inactive by multiple inhibitory factors secreted by the genital duct epithelia & they activated in female genital tract, for the processes of fertilization. These activation changes are called capacitation of the spermatozoa (require 1 to 10 hrs.).

1


Uterine & fallopian fluids wash away the inhibitory factors which suppress the sperm activity in the male genital ducts.

2

While the spermatozoa remain in the fluid of the genital ducts, they exposed to many floating vesicles from the seminiferous tubules containing large amount of cholesterol. This cholesterol is added to the cellular membrane covering the acrosome making it more rough & prevent the release of its enzyme. After ejaculation the sperm removed from the cholesterol vesicles & this makes the membrane of the sperm & head becomes weaker

3

The sperm membrane becomes more permeable to  $Ca^{2+}$  ion which increase their movements & help to release the proteolytic enzymes from acrosome which aid in penetrating the ovum




Acrosome enzymes, the “Acrosome Reaction” and penetration of the ovum:

The acrosome of the sperm stored large quantities of **hyaluronidase and proteolytic enzymes**. Hyaluronidase depolarizes hyaluronic acid polymers in the intracellular cement that hold the ovarian granulose cells together. Also the proteolytic enzymes digest the proteins.

# Hormonal factors that stimulate spermatogenesis:

| Hormone                       | Secreted by   | Functions   |
|-------------------------------|---|---|
| Testosterone                  | The <u>Leydig Cells</u> Which Located In The <u>Interstitial</u> Of The Testis,           | Essential For The <u>Growth</u> And <u>Division</u> Of The Testicular Germinal Cells  |
| Luteinizing hormone (LH)      | Anterior Pituitary Gland  | <u>Stimulates The Leydig Cells</u> To Secrete Testosterone  |
| Follicle stimulating hormone: | Anterior Pituitary Gland  | <u>Stimulates The Sertoli Cells</u> , Stimulate The Conversion Of Spermatids To Sperm (Also Important For Spermatogenesis)  |
| Estrogen                      | Formed From Testosterone By Aromatase enzyme in <u>Sertoli Cell</u> Under FSH Stimulation | <u>Essential For Spermatogenesis</u>  |
| Growth hormone                | Anterior Pituitary Gland  | Is Necessary For <u>Controlling Metabolic Functions</u> Of The Testis. GH Promotes Early Division Of Spermatogonias In Its Absence (Pituitary Dwarfs), The Spermatogenesis Is Severely Deficient Or Absent → <u>Infertility</u> . |



# Male sexual act

## Stages of male sexual act:

### 1-Penile erection

- Erection is caused by **parasympathetic** impulses that pass from the **sacral portion** of the spinal cord through the pelvic nerves to the penis.

### 2-Lubrication

- **Parasympathetic** impulses cause the urethral glands & bulbourethral glands to secrete mucous

### 3-Emission and ejaculation

- Function of the **sympathetic** nerves. Emission begins by **contraction of the vas deferens & ampulla** to cause **expulsion of the sperm** in the internal urethra. **Contraction of the prostate & seminal vesicles** to **expel their fluid** in the urethra. All these fluid mix in the internal urethra with the mucous secreted by the bulbourethral glands to form the semen. This process at this point is called **emission**.



- **Filling of the internal urethra** with semen causes sensory impulses through pudendal nerves to the **sacral region** of the cord. Fullness of the internal urethra causes rhythmical contractions of the internal genital organs which increases their pressure to ejaculate the semen to the outside called **ejaculation**.



# Testosterone

## Testosterone and other male sex chromosomes: secretion, metabolism and chemistry of the male sex hormone:

- Secretion of testosterone by the interstitial cell of Leydig in the testis. The testis secretes several male sex hormones called **androgens** including testosterone, dihydrotestosterone and androstenedione.
- **Testosterone** is the **most abundant** form while **dihydrotestosterone** is **most active** and **testosterone** is converted into dihydrotestosterone in the target cells.

## Secretion & chemistry of androgens in the body:

- From the adrenal glands & synthesized either **from cholesterol** or **directly** from acetylcoenzyme A.


## Metabolism of testosterone :

- Testosterone bound with **beta globulin** and circulates in the blood for 30 minutes to several hours and is converted to estrogen in the liver and excreted either into the gut through liver bile or into the urine through the kidneys.

# Testosterone

## Functions of testosterone:

- It is responsible for the characteristic **masculine body**.
- During **fetal life** the testis are **stimulated by placenta chorionic gonadotropin** to produce testosterone throughout fetal life & the 10 weeks after birth then no more testosterone production during childhood & at puberty under the **anterior pituitary gonadotropic hormones** stimulation throughout life & then decline beyond 80 years to 50%.



## Function of testosterone during fetal development:

- Testosterone secreted by the genital widges & later by the fetal testis is responsible for **development of the male body characteristics** including the formation of penis & scrotum. & prostate gland, seminal vesicles & male genital ducts & **suppressing the formation of female genital organs**.

## Effect of testosterone to cause descent of the testis:

- The testis descend into the scrotum during the **last 2 to 3 months of gestation** when the testis begin secreting reasonable quantities of testosterone.

# Testosterone

## Effect of testosterone on development of adult primary and secondary sexual characteristics:

|   |   |
|---|---|
| <p><b>1-After puberty,</b> the increasing amounts of testosterone cause enlargement of the penis, scrotum &amp; testis &amp; secondary sexual characteristics.</p>  | <p><b>6-Testosterone increased protein formation and muscle development:</b><br/>Increase muscular development after puberty by 50% in muscle mass over that in female. Also increase in protein in non-muscle parts of the body. These effect due to the <b>anabolic effect</b> of testosterone.</p>   |
| <p><b>2- Effect on the distribution of body hair:</b><br/>Testosterone causes growth of hair:</p> <ol style="list-style-type: none"><li>1. over the pubis,</li><li>2. upward along the linea alba of the abdomen to the umbilicus;</li><li>3. on the face;</li><li>4. on the chest; 5) less often on other regions such as the back.</li></ol>  | <p><b>7- Testosterone increases bone matrix and causes Ca<sup>2+</sup> retention:</b><br/>Bones grown thicker &amp; deposit additional Ca<sup>2+</sup>. Thus it increases the total quantity of bone matrix &amp; causes Ca<sup>2+</sup> retention (anabolic effect). Testosterone has specific effect on the pelvis</p> <ol style="list-style-type: none"><li>1. narrow the pelvic outlet.</li><li>2. lengthen it.</li><li>3. cause the funnel-like shape instead of the broad ovoid shape of the female pelvis.</li></ol> <p>It <b>causes the epiphyses of the long bones to unite with the shafts</b> of the bones &amp; early closure of the epiphyses.</p> |
| <p><b>3-Baldness:</b><br/>Testosterone <u>decreases</u> the growth of hair on the <b>top of the head</b> (two factors 1) genetic background; 2) large quantities of androgenic hormones.</p>  | <p><b>8-Testosterone increases basal metabolism:</b><br/>It increases the basal metabolic rate by about 15% (indirectly as a result of the anabolic effect).</p>  |
| <p><b>4-Effect on voice:</b><br/>It causes <u>hypertrophy of the laryngeal mucosa</u>, enlargement of the larynx (typical adult masculine voice)</p>  | <p><b>9-Effect on red blood cells:</b><br/>It increases red blood cells/ml (due to increase metabolic rate).</p>  |
| <p><b>5-Testosterone increases thickness of the skin and can contribute to development of acne:</b><br/>Testosterone increases the thickness of skin over the body &amp; subcutaneous tissues. Also it increases the secretion of the sebaceous glands &amp; sebaceous glands of the face causing acne.</p>   | <p><b>10-Effect on electrolyte and water balance:</b><br/>It <u>increase</u> the <b>reabsorption of Na<sup>+</sup></b> in the distal tubules of the kidneys.</p>  |
| <p><b>The basic intracellular mechanism of action of testosterone:</b></p> <ul style="list-style-type: none"><li>• It increases the rate of protein synthesis in target cells. <u>Testosterone converted by the intracellular enzyme 5 <math>\alpha</math> reductase to dihydrotestosterone</u>, then it binds with cytoplasmic “receptor protein”. This combination moves to the nucleus where it binds a nuclear protein and induces protein formation.</li></ul> |   |

# Abnormalities of male sexual function:

## ✓ Effect of sperm count on fertility:

The quantity of ejaculated semen during coitus about 3-5 ml, each milliliter contains about 120 million sperm (normal male count vary between 35 million to 200 million sperm). Sperm count below 20 million leads to infertility.

## ✓ Effect of sperm morphology and motility on fertility:

Sometimes sperm count is normal but still infertile when about one half of the sperm having abnormal shape. If there is one sperm in an appropriate shape it will affects the normal others .

-Sometimes the shape of the sperm is normal but they either relatively non-motile or entirely non-motile which causes infertility.

| 1. Prostate gland and its abnormalities   | 2. Hypogonadism in male  |
|---|--|
| <ul style="list-style-type: none"><li>• Benign prostatic fibroadenoma in older age due to overgrowth of prostate tissue (<b>not caused by testosterone</b>).</li><li>• Cancer of the prostate gland caused by stimulation of cancerous cells by testosterone.</li></ul> | <ul style="list-style-type: none"><li>• During fetal life when the <b>testis are non-functional</b>, none of the male sexual characteristics develop in the fetus. <b>Instead female organs are formed.</b></li><li>• If the boy loses his testis <u>before puberty</u>, a state <u>eunuchism</u> (he have infantile sex organs &amp; infantile sexual characteristics) &amp; the height of an adult eunuch is slightly greater than normal because of slow union of the epiphyses.</li><li>• If a man is castrated <u>after puberty</u>, sexual organ regress in size and voice regress.</li><li>• <u>Adiposogenital syndrome, Frohlich's syndrome or hypothalamic eunuchism</u>: hypogonadism due to genetic <b>inability of the hypothalamus</b> to secrete normal amount of GnRH &amp; abnormality of the feeding center of the hypothalamus <u>result in obesity</u> with eunuchism</li></ul> |




## Cont. :


| <b>3. Cryptorchidism</b>  | <b>4. Testicular tumors and hypergonadism in male</b>   |
|---|---|
| <ul style="list-style-type: none"><li>• Failure of the testes to descend in the scrotum which normally occur during fetal life.</li><li>• 10% of newborn males and it falls to 2% at age 1 year</li><li>• 0.3% after puberty</li><li>• They should be treated before puberty because of higher incidence of malignant tumors.</li></ul> | <ul style="list-style-type: none"><li>• Interstitial leydig cell tumors (rare), over production of testosterone. In children, causes rapid growth of the musculature and bones and early uniting of the epiphyses and causes excessive development of male sexual organs.</li><li>• Tumor of the germinal epithelium (more common).</li></ul> |

# Summery

## Hormone Control for Male Reproduction



## TESTOSTERONE



# MCQs

**1Q: Formation of sperms occur in :**

- A- seminiferous tubules .
- B- sertoli cell .
- C- leyding cell .
- D- spermatic cord .

**2Q: majority of sperms are stored in :**

- A- vas deference .
- B- epididymis .
- C- prostate .
- D- seminal vesicle .

**3Q: sertoli cells and epithelium of the epididymis secrete fluid which contains :**

- A- Estrogene.
- B- Nutrients & enzymes .
- C- testosterone .
- D- all .

**4Q: The acrosome of the sperm stores large quantities of :**

- A- hyaluronidase.
- B- proteolytic enzymes.
- C- A & B .
- D- IL . ( INTER LUKINS ) .

**5Q : Which one of the following hormones stimulate spermatogenesis :**

- A- Testosterone & androgens .
- B- seminal fluid .
- C- none .
- D- a & b .

**6Q : necessary for controlling metabolic functions of the testis:**

- A- GH .
- B- PROLACTIN .
- C- ACTH.
- D- none .

**7Q : Which of the following stimulate sertoli cell :**

- A- FSH .
- B- LH .
- C- testosterone .
- D- semen .

**8Q: Alkaline prostate fluid function :**

- A- fertilizes the ovum .
- B- sperm movement .
- C- peristaltic contractions .
- D- secrete mucoid material .

**9Q: It is responsible for the characteristic masculine body :**

- A- Testosterone .
- B- norepinephrine .
- C- acetyl-coA .
- D- none .

**10Q: the testis descend into the scrotum during the last 2 to 3 months of gestation when the testis begin secreting reasonable quantities of :**

- A- testosterone .
- B- cholesterol .
- C- CHD.
- D- estrogen .

1-A 2-A 3- D 4-C 5- A 6-A 7- A 8-A 9- A 10-A



**@PhysiologyTeam**



**Pht433@gmail.com**



Physiology team

**Done by :**

**Aisha Alraddadi**

**Awatif Alenazi**

**Shahad Almuhaideb**

**Hajar Alotaibi**

**Revised by:**

**Mojahed otayf**

**Rahma Alshehri**

**Reproductive Block**