

# Physiology of Motor Tracts

Dr Syed Shahid Habib  
*Professor*  
*Dept. of Physiology*  
*King Saud University*

## OBJECTIVES

At the end of this lecture the student should be able to:


- (A) Appreciate what is upper motor neuron and lower motor neuron .
- (B) List the main differences between the pyramidal and extrapyramidal systems .
- (C) explain the origin , course and functions of the following motor tracts :
  - (1) corticospinal.
  - (2) tectospinal .
  - (3) rubrospinal .
  - (4) vestibulospinal .
  - (5) reticulospinal .
  - (6) olivospinal .



## MOTOR SYSTEM LAYOUT


**Motor = movement**  
**For movement we need**  
**nervous system and muscles.**

### Components of motor system are:


- **Motor cortex**
- **Upper and lower motor neurons**
- **Cerebellum**
- **Basal ganglia**
- **Spinal cord**
- **Muscles**









**FIGURE 12.7** Control of voluntary movement. Commands for voluntary movement originate in cortical association areas. The cortex, basal ganglia, and cerebellum work cooperatively to plan movements. Movement executed by the cortex is relayed via the corticospinal tracts and corticobulbar tracts to motor neurons. The cerebellum provides feedback to adjust and smooth movement.





## DESCENDING TRACTS


### TWO MAJOR GROUPS

1. CORTICOSPINAL OR PYRAMIDAL TRACTS
2. EXTRAPYRAMIDAL TRACTS

## PYRAMIDAL TRACTS

### Corticospinal tract originates from:

- 30% from primary motor cortex (Motor Area 4)
- 30% from premotor & supplementary motor areas
- 40% from somatosensory areas (somatic sensory area 3,1,2)




## PYRAMIDAL TRACTS

❖ **Originating Cells: Giant pyramidal cells**

❖ **80% to 90% of the pyramidal fibers then cross in medulla to the opposite side and descend into the lateral corticospinal tracts**

❖ **Anterior or ventral corticospinal tract is formed by uncrossed fibers and is primarily concerned with the control of fine movements that require dexterity.**



## COMPONENTS OF MOTOR NEURONS

**Upper motor neuron (corticospinal & corticobulbar).**

Starts from motor cortex and ends in

1. Cranial nerve nucleus (corticobulbar).
2. Anterior horn of spinal cord in opposite side (corticospinal tracts).

**Lower Motor Neuron**

Starts from anterior horn of spinal cord and & cranial motor neurons in the brain stem that innervate muscles directly ends in appropriate muscle of the same side.

### LATERAL AND VENTRAL CORTICOSPINAL TRACTS

- ❖ The majority of the pyramidal fibers (80-90%) then cross in the lower medulla to the opposite side and descend into the lateral corticospinal tracts and ends.....
  - directly on motor neurons (AHC) of the opposite side
  - interneurons of the grey matter
  - some ends at sensory neurons of dorsal horn
- ❖ **Functions:** Controls distal limb muscles for skilled voluntary movements (Fingers & Toes)

- ❖ Anterior or ventral corticospinal tract is formed by uncrossed fibers and many of these cross at level of termination to synapse with interneurons, specially in in the neck or in the upper thoracic region.
- ❖ **Functions:** These fibers may be concerned with control of bilateral postural movements by the supplementary motor cortex. Pass medially in ventral horn so control axial & proximal limb muscles in coordination with tectospinal, reticulospinal and vestibulospinal tracts.

### FUNCTIONS OF CORTICOSPINAL TRACTS

- 1-Initiation of fine ,discrete, skilled voluntary movements  
(on which side?)
- 2- lateral corticospinal tracts (main bulk of the tract) control distal muscles of limb as fingers & thumb& toes
- 3- Ventral corticospinal tracts control posture
- 4- Effect on stretch reflex:-
  - Facilitate muscle tone through gamma motor neurons
- 5- Fibers originate from parietal lobe are for sensory-motor coordination
- 6- corticobulbar tracts /control face & neck muscles & facilitate their tone, and are involved in facial expression, mastication, swallowing.



## Effect of Lesions in the Motor Cortex or in the Corticospinal Pathway—The “Stroke”

The motor control system can be damaged by the “stroke.”-

-Removal of the Primary Motor Cortex (Area Pyramidalis)

- If the caudate nucleus and adjacent premotor and supplementary motor areas are not damaged, gross postural and limb “fixation” movements can still occur, but there is loss of voluntary control of discrete movements of the distal segments of the limbs, especially of the hands and fingers.
- Area pyramidalis is essential for what??

essential for voluntary initiation of finely controlled movements, especially of the hands and fingers

## Muscle Spasticity Caused by Lesions That Damage Large Areas Adjacent to the Motor Cortex.

The primary motor cortex normally exerts a continual tonic stimulatory effect on the motor neurons of the spinal cord; when this stimulatory effect is removed, *hypotonia* results.

-Most lesions of the motor cortex, especially those caused by a *stroke*, involve not only the primary motor cortex but also adjacent parts of the brain such as the basal ganglia. In these instances, *muscle spasticity* almost invariably occurs in the afflicted muscle areas on the *opposite side* of the body.

This spasm results mainly from damage to accessory nonpyramidal pathways. These pathways normally inhibit the vestibular and reticular brain stem motor nuclei. When these nuclei cease their state of inhibition (i.e., are “disinhibited”), they become spontaneously active and cause excessive spastic tone in the involved muscles.

## **EXTRAPYRAMIDAL MOTOR TRACTS**

### **Originate in the midbrain and brain stem regions**

**If the pyramidal tracts of an experimental animal are cut, electrical stimulation of the cerebral cortex, cerebellum, and basal nuclei can still produce movements.**

The term extrapyramidal motor system is widely used in clinical circles to denote all those portions of the brain and brain stem that contribute to motor control but are not part of the direct corticospinal-pyramidal system.

### **Components of extrapyramidal system**

- 1. Basal Ganglia**
- 2. Midbrain giving rise to following bulbo-spinal tracts.**
  - A. Rubrospinal tract.**
  - B. Vestibulospinal Tract.**
  - C. Reticulospinal Tract**
  - D. Tectospinal Tract.**
  - E. Olivospinal Tract.**

**(1) sets the postural background needed for performance of skilled movements**

**(2) controls subconscious gross movements**

20

**Excitation of the Spinal Cord Motor Control Areas by the Primary Motor Cortex and Red Nucleus (Dynamic and Static signals)**

- Cells of the motor cortex are organized in vertical columns with thousands of neurons in each column
- Each column has 6 distinct layers and Pyramidal cells originate from Layer 5
- Each Column of neurons operate as an integrative processing and amplifying system (50-100 Pyramidal cells), using information from multiple input sources to determine the output response from the column for purposeful activities.
- Each column of cells excites two populations of pyramidal cell neurons, one called *dynamic neurons* and the other *static neurons*.

**Excitation of the Spinal Cord Motor Control Areas by the Primary Motor Cortex and Red Nucleus (Dynamic and Static signals) Cont.**

- The dynamic neurons are excessively excited for a short period at the beginning of a contraction, causing the initial rapid *development of force*. Then the static neurons fire at a much slower rate, but they continue firing at this slow rate to *maintain the force* of contraction as long as the contraction is required.
- Greater percentage of dynamic neurons is in the red nucleus and a greater percentage of static neurons is in the primary motor cortex


**Red Nucleus Serves as an Alternative Pathway for Transmitting Cortical Signals to the Spinal Cord**

▪ **RUBROSPINAL TRACTS**


- Red Nucleus receives large number of direct fibers from the primary motor cortex through the *corticorubral tract* & some branching fibers from the corticospinal tract as it passes through the mesencephalon.
- These fibers synapse in the *magnocellular portion*, containing large neurons like betz cells and give origin to Rubrospinal tract.
- The red nucleus also has close connections with the cerebellum

- **LATERAL MOTOR SYSTEM** : the corticospinal and rubrospinal tracts together
- **MEDIAL MOTOR SYSTEM**: vestibuloreticulospinal system, lies mainly medially in the cord

The corticorubrospinal pathway serves as an accessory route for transmission of relatively discrete signals from the motor cortex to the spinal cord.






Corticorubrospinal pathway for motor control, showing also the relation of this pathway to the cerebellum.



**Figure 48-8**

Cross section of the spinal cord, showing principal ascending tracts on the right and principal descending tracts on the left.






## Vestibulospinal tracts

-Originates from vestibular nucleus in pons (which receive inputs from inner ear, Vestibular Apparatus and cerebellum)

-Axons descend in the ipsilateral ventral white column of spinal cord

### Functions:-

- 1- Controls Postural & righting reflexes.
- 2- Excitatory to ipsilateral spinal motor neurons-that supply axial & postural muscles
- 3- Control head & eye movements



## Functions of Vestibulospinal Tracts

### The lateral vestibulospinal tract

- Cells of origin : Lateral Vestibular Nucleus
- Axons descend in the ventral white column of spinal cord .

**Functions:** It activates motor neurons to antigravity muscles (eg, proximal limb extensors) to control posture and balance.

### The medial vestibulospinal tract

- Cells of origin : Medial and inferior Vestibular Nuclei
- As its axons descend in the ventral white column of spinal cord to end at the cervical segments of the spinal cord, some fibers form part of the Medial Longitudinal Fasciculus fibres in brain stem that link vestibular nuclei to nuclei supplying the extra-ocular muscles.

**Functions:** Coordination of head and eye movements

## Reticulospinal Tract

- The reticular formation makes up a central core of the brainstem. It contains many different neuronal groups.
- Pontine and medullary nuclei projects to the AHCs of the spinal cord via Reticulospinal Tract

### Functions:

- 1-Influence motor functions as voluntary & reflex movement
- 2-Excitatory or inhibitory to muscle tone





Figure 55-8

Vestibulospinal and reticulospinal tracts descending in the spinal cord to excite (*solid lines*) or inhibit (*dashed lines*) the anterior motor neurons that control the body's axial musculature.



### Two types of reticulospinal tracts

#### (1) Pontine (Medial) Reticulospinal Tract (anterior column of spinal cord)

- Transmit excitatory signals downward into the cord → medial anterior motor neurons that excite the axial muscles of the body, which support the body against gravity. In addition, they receive strong excitatory signals from the vestibular nuclei & from deep nuclei of the cerebellum.

- Axons descend in anterior (ventral) white column of spinal cord

- **Functions:** Excite the antigravity muscles of the body and increases Gamma efferent activity

#### (2) Medullary (Lateral) Reticulospinal Tract:

Axons descend in lateral white column of spinal cord on both sides

It receive strong input from (1) the corticospinal tract, (2) the rubrospinal tract, and (3) other motor pathways → activate the medullary reticular inhibitory system to counterbalance the excitatory signals from the pontine reticular system

- **Functions:** counterbalance the excitatory signals from the pontine reticular system and inhibits Gamma efferent activity

The excitatory and inhibitory reticular nuclei constitute a controllable system that is manipulated by motor signals from the cerebral cortex to provide necessary background muscle contractions for standing against gravity



### Tectospinal tracts

It arises from superior (VISUAL) & inferior colliculi (AUDITORY) of midbrain and Ends on Contralateral cervical motor neurons

**Functions:** Mediate/facilitate turning of the head in response to visual or Auditory stimuli

### Olivospinal Tract

It arises from inferior olivary N of the medulla & is found only in the cervical region of the spinal cord (supply neck muscles). secondary olivocerebellar fibers transmit signals to multiple areas of the cerebellum.

**Functions:** Exact function is not known but it is believed to be involved in reflex movements arising from proprioceptors.




Figure 55-6

Convergence of different motor control pathways on the anterior motor neurons.