




# Thalamus & Limbic System



Prof. Saeed Abuel Makarem &  
Dr.Sanaa Alshaarawy


# Objectives

***By the end of the lecture, you should be able to:***

- Describe the anatomy and main functions of the thalamus.
- Name and identify different nuclei of the thalamus.
- Describe the main connections and functions of thalamic nuclei.
- Name and identify different parts of the **limbic system**.
- Describe main functions of the **limbic system**.
- Describe the effects of lesions of the **limbic system**.

# Thalamus

- It is the **largest nuclear mass** of the whole body.
- It is the **largest part** of the **diencephalon**
- It is formed of **two oval masses** of **grey matter**.
- It is the **gateway to the cortex**.
- Together **with** the **hypothalamus** they form the **lateral wall of the 3<sup>rd</sup> ventricle**.



- It sends received information to the cerebral cortex from diverse brain regions.

- Axons from every sensory system (**except olfaction**) synapse in the thalamus as the **last relay site** '**last pit stop**' before the information reaches the cerebral cortex.

- There are some thalamic nuclei that receive input from:

1. Cerebellar nuclei,
2. Basal ganglia- and
3. Limbic-related **brain regions**.

# Thalamus



It has 4 surfaces & 2 ends.

## Surfaces

### Lateral:(L)

Posterior limb of the **internal capsule**

### Medial:

The **3<sup>rd</sup> ventricle**

It is connected to the thalamus of the opposite side by the **interthalamic connexus**, **(adhesion) or Massa intermedia**.


### Superior: (s)

Lateral ventricle and **fornix**.

### Inferior: (I)

**Hypothalamus**, anteriorly & **Subthalamus** posteriorly.

## Relations



## Anterior end:



Forms a projection, called the **anterior tubercle**.

It lies just behind the *interventricular foramen*.

## Posterior end: Broad

Forms a projection called **Pulvinar** which lies above the *superior colliculus* and the lateral & medial *Geniculate bodies*.

Thalamus



## ● White matter:


### ● External medullary lamina:

- Covers the lateral surface.
- It consists of thalamocortical & corticothalamic fibers.

### ● Internal medullary lamina:


- Bundle of Y-shaped myelinated (afferent & efferent) fibers.
- It divides the thalamus into: **anterior , medial, lateral nuclear groups.**
- Each of these groups is subdivided into a number of named nuclei.

## Internal Structure



- It is divided into:  
Dorsal & Ventral tiers
- Dorsal tier:**
- which contains:**
  - Lateral Dorsal (LD)&
  - Lateral Posterior (LP)
  - Pulvinar.
- Ventral tier,**
- which contains :**
  - Ventral Anterior (VA)
  - Ventral Lateral (VL)
  - Ventral Intermediate (VI)
  - Ventral Posterior (VP)  
(PLVNT, PMVNT)
  - Lateral & Medial  
Geniculate nuclei.


## Lateral Nuclear Group



# Projection of thalamic nuclei


## Anterior Thalamic Nucleus

- **Afferent:** Mammillary body.
- **Efferent:** Cingulate gyrus, (limbic system)



## Medial Nucleus

- **Afferent:** Hypothalamus.
- **Efferent:** Frontal & Prefrontal cortex.



# Projection of thalamic nuclei


## Ventral Anterior Nucleus

- **Afferent:** Globus pallidus body.
- **Efferent:** Premotor cortex.
- 


## Ventral Lateral Nucleus

- **Afferent:** Dentate Nucleus
- **Efferent:** primary motor cortex.

## Input of Ventral Thalamic Nuclei



## Output of thalamic nuclei



# Projection of thalamic nuclei


## Ventral Posterior Lateral Nucleus

- **Afferent:** Medial and spinal lemnisci.
- **Efferent:** Sensory cortex.
- 


## Ventral Posterior Medial Nucleus

- **Afferent:** Trigeminal Lemniscus
- **Efferent:** Sensory cortex.

### Input of Ventral Thalamic Nuclei



### Output of thalamic nuclei




# Projection of thalamic nuclei


- **Lateral geniculate body :**
- **Afferent :** optic tract.
- **Efferent :** visual cortex

- **Medial geniculate body :**
- **Afferent :** lateral lemniscus.
- **Efferent :** auditory cortex.


## Input of Ventral Thalamic Nuclei



## Output of thalamic nuclei




# INPUT AND OUTPUT OF THALAMIC NUCLEI



# LIMBIC SYSTEM

- The term "limbic" is from the Latin word ***Limbus***, for "border" or "edge".
- It separates the medial surface of the **cerebral cortex** from the **diencephalon**
- It consists of a number of **cortical & subcortical** structures with **looped connections** that all project to the **hypothalamus** (particularly mammillary bodies).



# WHAT IS THE FUNCTION OF THE LIMBIC SYSTEM?

It control a variety of functions including:

- ❖ **Emotions:**
- ❖ Emotional responses
- ❖ Behaviour & Mood  
(happy, cry, laugh, sad, afraid, aggression, depression)
- ❖ Motivation.
- ❖ **Memory.**
- ❖ **Visceral & Motor responses** involved in (sex, **pleasure**, hunger, and reproduction).
- ❖ **Olfaction.**



MEMORY

Pleasure sensation

OLFACTION



# The limbic system is a set of brain structures including


The limbic system is composed of four main structures:

- 1. Limbic cortex**
- 2. Hippocampus**
- 3. Amygdala, &**
- 4. Septal area.**

- These structures **form connections** between the hypothalamus, thalamus and cerebral cortex.

- The **hippocampus** is important in memory and learning, while the **limbic system itself** is important in the control of the emotional responses.


## The Limbic System



Legend:  
■ Limbic Gyrus   ■ Intralimbic Gyrus   ■ Fornix & Inner Arc

# CORTICAL STRUCTURES

1. **Limbic lobe.**
2. **Hippocampal formation.**
3. **Septal areas**  
**(Fornix, connecting the hippocampus with mammillary bodies and septal nuclei).**
4. **Prefrontal area(part of olfactory system)**




# LIMBIC LOBE

- C-shaped ring of grey matter on the medial side of each cerebral hemisphere, surrounding the corpus callosum.

- It includes:

1. Subcallosal area
2. Cingulate gyrus
3. Isthmus
4. Parahippocampal gyrus and the
5. Uncus.



# HIPPOCAMPUS

It is a limbic system structure that is involved in:

**Formation,**  
**Organization**, and  
**Storing** of **memories**.

It is important in **forming new memories** and **connecting emotions and senses**, such as **smell** and **sound**, to memories.


**It is a horseshoe paired structure**, one in each cerebral hemisphere.

It acts as a memory indexer by sending memories to the appropriate part of the **cerebral hemisphere** for **long-term storage** and retrieving them when necessary.



- **Site:**
- It is a scrolled (infolding) inferomedial part of temporal lobe.
- **Function:**
- Memory (file new memories as they occur).
- The hippocampus & its connections are necessary for **consolidation** of **new short-term memories.**

# HIPPOCAMPUS



- Its principal **efferent pathway** is called the: **FORNIX**:


*It is C-shaped group of fibers connecting the hippocampus with mammillary body and then to the anterior nuclei of thalamus.*

*It consists of:*

*Fimbria,  
Crus,  
Body &  
Column.*


- The **Fornix** is an important component of **PAPEZ CIRCUIT** (based on connecting the hypothalamus with limbic lobe to control emotions )

# HIPPOCAMPUS




- **Site:**
- almond shaped **mass of nuclei** that **lies near the temporal pole**, deep within the temporal lobes ,close to the **tail of the caudate nucleus.**
- **Function:**
- It is involved in **FEAR**,
- **Emotions**
- **Anger, &**
- **Hormonal secretions.**

# AMYGDALA



# CONNECTIONS OF AMYGDALA

- **Inputs:**
- **Association areas** of **visual**, **auditory** & **somatosensory** cortices.
- **Outputs:**
- **Hypothalamus** &
- **Autonomic nuclei** in the **brain stem**,
- **Lesion:**  
Lack of emotional responses & docility.



# Septal nuclei

## Site:


Located anterior to the interventricular septum

## Main connections:

1. To Hypothalamus
2. To Habenular nuclei

## Function:

It is the **pleasure** zone.



⦿ **Korsakoff's psychosis :** Korsakoff syndrome is a chronic memory disorder caused by severe deficiency of thiamine (vitamin B-1) & alcoholic intoxication.

⦿ (**Retrograde** = loss of retained old memories occurred before the injury & **anterograde amnesia**= inability to gain new memories).

⦿ **Temporal lobe epilepsy**

⦿ The **hippocampus** is a common focus site in epilepsy, and can be damaged through **chronic seizures**.


⦿ It is **sometimes** damaged in diseases such as **herpes encephalitis**,

⦿ **Alzheimer's disease:**

**hippocampus** is one of the first brain areas to show damage in Alzheimer's disease. **Anterograde amnesia** —the inability to form and retain new memories.

⦿ **Schizophrenia.** (mental disorder with inappropriate actions and feelings),

## Lesions associated with limbic lobe disorders





**THANK YOU**