


Histology of the Eye

Objectives

- By the end of this lecture, the student should be able to describe:
 - *The general structure of the eye.*
 - *The microscopic structure of:*
 - » *Cornea.*
 - » *Retina.*


EYE BULB

- Three coats (3 Tunics):
- **1- Fibrous tunic:**
 - Cornea.
 - Sclera.
- **2- Vascular tunic:**
 - Choroid.
 - Ciliary body.
 - Iris.
- **3- Neural tunic:**
 - Retina.



CORNEA

- It is the transparent, avascular and highly innervated anterior portion of the fibrous coat.
- It is composed of 5 distinct layers:
 1. Corneal epithelium.
 2. Bowman's membrane.
 3. Stroma.
 4. Descemet's membrane.
 5. Corneal endothelium.




CORNEA (Cont.)

■ Corneal epithelium:

- Non-keratinized Stratified squamous epithelium.
- Contains numerous free nerve endings.

■ Bowman's membrane:


- It is homogenous non-cellular layer containing type I collagen fibrils.



CORNEA (Cont.)

■ Stroma:

- It is the thickest layer (about 90%).
- It is composed of parallel lamellae of dense collagenous C.T.
- Each lamella is composed mainly of parallel type I collagen fibers with long fibroblasts.



CORNEA (Cont.)

■ Descemet's membrane:

- It is a thick basement membrane.

■ Corneal endothelium:

- It is simple squamous epithelium.

– Functions:



1- Formation of Descemet's membrane.

2- Keeping the stroma relatively dehydrated

(sod. pump → water withdrawal from the stroma).


LIMBUS (CORNEO SCLERAL JUNCTION)

- It is the transition region between the cornea and sclera.
- It is about 1.5 mm width.
- It is highly vascular.
- It contains:
 1. Trabecular meshwork:
Endothelium-lined spaces. It leads to canal of Schlemm.
 2. Canal of Schlemm:
It drains the aqueous humor into the venous system.



SCLERA

- It covers the posterior 5/6 of the fibrous tunic.
- **Sclera Proper:** consists of interlacing bundles of type I collagen (dense collagenous C.T., irregular type).
- Melanocytes are located in the deeper regions.





CHOROID

- It is the vascular, pigmented posterior portion of the middle vascular tunic.

■ **Structure:**


It is composed mainly of loose C.T. with melanocytes.

It is separated from the retina by its Bruch's membrane.





CILIARY BODY

- It is the anterior continuation of the choroid. It surrounds the lens.
- **Structure:**
 - It is formed of loose vascular and pigmented C.T. that contains 3 bundles of smooth muscle cells (ciliary muscle).
 - Its inner surface is lined by pars ciliaris retinae (2 rows of columnar cells; outer pigmented and inner non- pigmented layers) .
 - Its inner surface is highly folded forming the ciliary processes.



CILIARY PROCESSES



- Processes project from the inner surface of the anterior 1/3 of the ciliary body towards the lens.
- Are covered by pars ciliaris retinae (2 rows of columnar cells).
- They give attachment to the lens suspensory ligaments (zonule fibers).



IRIS

It is formed of 5 layers:

- 1- Anterior border layer:
Incomplete layer of fibroblasts and melanocytes.
- 2- Stroma:
Poorly vascularized C.T. with fibroblasts and melanocytes.
- 3- Vessel layer:
Well-vascularized loose C.T.
Centrally, it contains circularly arranged smooth muscle fibers (sphincter pupillae muscle).



IRIS


- 4- Dilator pupillae muscle layer:

Contains radially arranged myoepithelial cells.

- 5- Posterior surface layer (pigmented epithelium layer):

It is composed of 2 rows of **pigmented** epithelial cells (pars iridis retinae).



They are the continuation of pars ciliaris retinae.





RETINA

It is composed of **10 distinct layers** (from outside to inside):


- 1- Pigmented epithelium.
- 2- Rods and cones layer.
- 3- Outer limiting membrane.
- 4- Outer nuclear layer.
- 5- Outer plexiform layer.
- 6- Inner nuclear layer.
- 7- Inner plexiform layer.
- 8- Ganglion cell layer.
- 9- Optic nerve fiber layer.
- 10- Inner limiting layer.



RETINA (Cont.)



© Deltagen Inc.



RETINA (Cont.)

- **Pigmented Epithelium:**
 - Cuboidal to columnar cells (single layer).
 - Apical microvilli.
 - Abundance of melanin granules.
- Functions:
 - 1- Absorb light.
 - 2- Phagocytosis of membranous discs from tips of rods.
 - 3- Esterification of Vitamin A (in SER).

RODS AND CONES LAYER (Cont.)

- Are photoreceptor cells.

- Each has:

1. Dendrite formed of:

- Outer segment (OS): contains membranous discs containing rhodopsin (in rods) and iodopsin (in cones).

- Connecting Stalk: with modified cilium.

- Inner segment (IS).


2. Cell body.

3. Axon: synapses with dendrite of bipolar neuron of inner nuclear layer.

- **Functions:**

Rods are receptors for dim light (low intensity light).

Cones are receptors for bright light and color vision (red, green & blue).



RETINA (Cont.)

■ Outer limiting membrane:

- A region of zonulae adherentes junctions between Muller cells and the photoreceptors.

■ Outer nuclear layer:

- Contains nuclei of the rods & cones.


■ Outer plexiform layer:

- Contains axodendritic synapses between the photoreceptor cells and dendrites of bipolar and horizontal cells.

■ Inner nuclear layer:

- Contains the nuclei of:

- 1- Bipolar neurons.
- 2- Horizontal neurons.
- 3- Amacrine neurons (unipolar neurons):
- 4- Neuroglial cells (Muller cells) that extend between the vitreous body and the inner segments of rods and cones.



RETINA (Cont.)

■ Inner plexiform layer:

Contains axodendritic synapses between axons of bipolar neurons and dendrites of ganglion cells and amacrine cells.

■ Ganglion cell layer:

Contains cell bodies of large multipolar neurons of the ganglion cells.


■ Optic nerve fiber layer:

Contains unmyelinated axons of the ganglion cells.

N.B. These axons become myelinated as the nerve pierces the sclera.




■ The inner limiting membrane:

It is formed by the basal laminae of the Muller cells.



RETINA (Cont.)

- **Fovea centralis:**
- It lies in the center of macula lutea.
- Cones are highly concentrated in the fovea.
- It is responsible for visual acuity.



RETINA (Cont.)

■ Types of cells in the retina:


1- Pigmented epithelium.

2- Nerve cells:

- Photoreceptor cells (rods & cones)
- Bipolar neurons.
- Ganglion cells.
- Association neurons:
 - i. Horizontal cells.
 - ii. Amacrine cells.

3- Neuroglial cells:

- Muller's cells.
- Astrocytes.



CONJUNCTIVA

- It is the transparent mucous membrane lining the inner surfaces of the eyelids (**palpebral conjunctiva**) and reflecting onto the sclera of the anterior surface of the eye (**bulbar conjunctiva**).

- **L/M:**

- **1- Epithelium:**

Stratified columnar epithelium with numerous goblet cells.

- **2- Lamina propria:**

Loose C.T.



**GOOD
LUCK**