

OMENTUM

ANATOMY DEPARTMENT

DR.SANAA AL-SHAARAWY


Dr. Essam Eldin Salama

OBJECTIVES


- *At the end of the lecture the students must know:*
- Brief knowledge about **peritoneum** as a thin serous membrane and its main parts; **parietal** and **visceral**.
- The **peritoneal cavity** and its parts the **greater sac** and the **lesser sac** (Omental bursa).
- **The peritoneal folds** : omenta, mesenteries, and ligaments.
- The **omentum**, as one of the peritoneal folds
- The **greater omentum**, its boundaries, and contents.
- The **lesser omentum**, its boundaries, and contents.
- The **omental bursa**, its boundaries.
- The **Epiploic foramen**, its boundaries.
- **Mesentery** of the small intestine, and **ligaments** of the liver.
- **Nerve supply** of the peritoneum.
- **Clinical points**.

The peritoneum

- ❖ Is a **thin serous membrane**,
- Lining the wall of the abdominal and pelvic cavities, (the **parietal peritoneum**).
- Covering the existing organs, (the **visceral peritoneum**).
- The potential space between the two layers is the **peritoneal cavity**.



The peritoneal Cavity



❖ The **peritoneal cavity** is the largest one in the body.

❖ Divisions of the peritoneal cavity :

▪ **Greater sac**; extends from diaphragm down to the pelvis.


▪ **Lesser sac**; lies behind the stomach.

▪ Both cavities are interconnected through the **epiploic foramen**.

▪ In male : the peritoneum is a closed sac .

▪ In female : the sac is not completely closed because it communicates with the exterior through the uterine tubes, uterus and vagina.


The peritoneum



Schoenwolf et al: Larsen's Human Embryology, 4th Edition.
Copyright © 2008 by Churchill Livingstone, an imprint of Elsevier, Inc. All rights reserved

partially covered by visceral peritoneum.

Intraperitoneal viscera



Retroperitoneal viscera

Intraperitoneal organ :



Is entirely surrounded by the peritoneum and has a supporting mesentery : stomach & 1st part of duodenum, liver, gall bladder, spleen, jejunum, ileum, transverse colon, sigmoid colon, uterus, and ovaries.

Extraperitoneal or retroperitoneal organ :

Structure that lies behind the peritoneum or partially covered by the peritoneum and has no supporting mesentery.

Primary retroperitoneal organs: Aorta, Inferior vena cava, kidneys, Suprarenal glands, urinary bladder, vagina, and rectum.

Secondary retroperitoneal organs: develop in mesenteries, but get pushed against the body wall (parietal peritoneum) during growth so that only half of their surface is covered by peritoneum : pancreas, duodenum, ascending and descending colon.






Folds of the peritoneum

□ Types of peritoneal folds :


- **Omenta.**
- **Mesenteries.**
- **Ligaments.**

The peritoneal ligaments, omenta, and mesenteries permit blood, lymph vessels, and nerves to reach the viscera





Omenta

- ❖ **Two layered fold of peritoneum connecting the stomach to another viscus.**
- **The lesser omentum attaches the lesser curvature of the stomach to the liver.**
- **The greater omentum connects the greater curvature of the stomach to the transverse colon.**





Lesser omentum





- ❑ Extends between the **liver** and the **lesser curvature of the stomach+1st part of duodenum**.
- It is continuous with the two layers of **peritoneum** which cover the anterior & posterior surfaces of stomach and 1st part of the duodenum.
- Ascends as a **double fold** to the **porta hepatis of liver**, and fissure for ligamentum venosum.
- To the **left** of porta hepatis it is carried to the **diaphragm**.
- Its **right** border is a **free margin**; constitutes the anterior boundary of the epiploic foramen.
- ❑ Contents between the two layers of the lesser omentum :
 - Close to the right free margin, are the **hepatic artery, common bile duct, portal vein, lymphatics, and hepatic plexus of nerves**.
 - At the attachment to the stomach, run the **right and left gastric vessels**.

Greater omentum




Greater omentum



- **The largest peritoneal fold**, with cribriform appearance, contains some adipose tissue.
- **It consists of** a double sheet of peritoneum, folded on itself so that it is made up of four layers (anterior 2 layers + posterior 2 layers).
- **The two layers** which descend from the greater curve of the **stomach** and commencement of the duodenum, pass downward in front of the small intestines, then turn upon themselves, and ascend to the **transverse colon**, where they separate and enclose it.
- **Its left border** is continuous with the **gastrosplenic ligament**.
- **Its right border** extends as far as the commencement of the duodenum.
- **Contents** : the **anastomosis** between the **right and left gastroepiploic vessels**.

Omental bursa, (Lesser Sac)





❑ It is a part of the peritoneal cavity behind the stomach.

❑ Boundaries of the *omental bursa* ;

▪ Anterior wall, from above downward, by the **caudate lobe** of the liver, the **lesser omentum**, back of the **stomach**, and the **anterior two layers** of the **greater omentum**.

▪ Posterior wall, from below upward, by the posterior two layers of the **greater omentum**, the **transverse colon**, and the ascending layer of the **transverse mesocolon**, the upper surface of the **pancreas**, the **left suprarenal gland**, and the upper end of the **left kidney**.


Epiploic foramen



- It is the communication between the greater and lesser sacs .
- It is bounded by;
- In front by the **free border** of the lesser omentum, with its contents : **hepatic artery, common bile duct, and portal vein** between its two layers.
- Behind by the **peritoneum** covering the **inferior vena cava**.
- Above (roof) by the **peritoneum** on the **caudate process** of the liver.
- Below (floor) by the **peritoneum** covering the commencement of the **duodenum** and the **hepatic artery**, before ascending between the two layers of the lesser omentum.

Mesentery

- Two-layered fold of peritoneum suspends the small intestine from the posterior abdominal wall
- Broad and a fan-shaped
- Intestinal border—folded, 7 m long
- Root of mesentery :
 - 15 cm long
 - Directed obliquely from duodenojejunal flexure at the level of left side of L2 to the ileocecal junction in the right iliac fossa at the level of right sacroiliac joint.





Ligaments

Two-layered folds of peritoneum that attach solid viscera to the abdominal wall and diaphragm.

Ligaments of liver

- Falciform ligament of liver
- Coronary ligament
- Left and right triangular ligaments
- Ligamentum teres



Nerve Supply of the Peritoneum

- **The parietal peritoneum** is sensitive to pain, temperature, touch, and pressure.

The parietal peritoneum lining the anterior abdominal wall is supplied by :

- **lower six thoracic (lower 6 intercostal) and first lumbar nerves.**
- The central part of the diaphragmatic peritoneum is supplied by the **phrenic nerves.**

- **The visceral peritoneum** is sensitive only to stretch and tearing.

It is supplied by :

- **autonomic nerves** that supply the viscera or traveling in the mesenteries.

Clinical points

□ Peritoneal Pain

From the Parietal Peritoneum

Abdominal pain originating from the parietal peritoneum is therefore **of the somatic type**, it is **usually severe**, and can be **accurately localized**.

From the Visceral Peritoneum

The visceral peritoneum, including the mesenteries, is **innervated by autonomic nerves**.

It is due to Stretch caused by **over distension of a viscus** and pulling on a mesentery

That gives rise to the sensation of pain.


[leading to **abdominal pain** poorly localized, poorly characterized pain. (dull ,unclear, cramping)]

Clinical points

□ Peritoneal Dialysis:

Because the peritoneum is a semi permeable membrane :

- It allows transfer of substances (glucose solution) across itself to remove the waste products.
- It has been used of in patients with acute renal insufficiency.



THANK YOU

SUMMARY

- **The peritoneum** is divided into 2 layers :

- **Parietal layer**, lines the abdominal and pelvic walls.

- **Visceral layer**, covers the abdominal and pelvic organs.

Folds of the peritoneum : Omenta, Mesenteries, and Ligaments.

- **Omenta**: are folds of peritoneum.

- **Lesser omentum** connects the lesser curvature of stomach and 1st part of duodenum to the liver.

- Right border of lesser omentum is free and it forms the anterior boundary of epiploic foramen.

- **Contents of lesser omentum :**

- right & left gastric vessels.

- Hepatic artery.

- Bile duct.

- Portal vein.

- Nerves, lymph vessels & fat.

SUMMARY

- **Greater omentum** : connects the greater curvature of stomach with the transverse colon.
- **Contents of greater omentum** :
 - Right & left gastroepiploic vessels.
 - Lymph nodes, vessels & fats.
- **Lesser sac of peritoneum (Omental Bursa)** :
 - **Boundaries** :
 - **Opening to lesser sac (epiploic foramen)** :
 - **Contents in the free margin of lesser omentum.**

SUMMARY

Mesentery: two-layered fold of peritoneum

suspends the small intestine from the posterior abdominal wall.

Ligaments: two-layered folds of peritoneum that attach solid viscera to the abdominal wall.

Function of peritonuem: *The peritoneal ligaments, omenta, and mesenteries permit blood, lymph vessels, and nerves to reach the viscera.*

Nerve Supply of the Peritoneum :

parietal peritoneum: lower six thoracic and first lumbar nerves and the phrenic nerves.

visceral peritoneum: autonomic nerves that supply the viscera.

Clinical aspects :

- Peritoneal Pain.
- Peritoneal Dialysis.