

GIT BLOCK

Gastrointestinal Diseases

Dr. Maha Arafah

Pathology, 2016

- **8 LECTURES**

Gastro-esophageal reflux disease

Peptic Ulcer Disease

Diarrhea

Malabsorption

Colonic polyps and carcinoma-1

Colonic polyps and carcinoma-2

Inflammatory bowel disease-1

Inflammatory bowel disease-2


Gastro-esophageal reflux disease

OBJECTIVES


- **Describe the following aspects of reflux esophagitis:**
 - 1) Definition
 - 2) Pathogenesis
 - 3) Clinical features
 - 4) Pathology (gross and microscopic features)
 - 5) Complications
- **Describe the following aspects of Barrett esophagus:**
 - 1) Definition
 - 2) Main cause
 - 3) Pathology (gross and microscopic features)
 - 4) Complications (dysplasia and adenocarcinoma)

Definition

- American College of Gastroenterology (ACG)
 - Symptoms OR mucosal damage produced by the abnormal reflux of gastric contents into the esophagus
 - Often chronic and relapsing
 - May see complications of GERD in patients who lack typical symptoms



Reflux Esophagitis



Gastroesophageal Reflux Disease (GERD)

- Gastroesophageal reflux is a normal physiologic phenomenon experienced intermittently by most people, particularly after a meal.
- Gastroesophageal reflux disease (GERD) occurs when the amount of gastric juice that refluxes into the esophagus exceeds the normal limit, causing symptoms with or without associated esophageal mucosal injury.

Physiologic vs Pathologic

- Physiologic GER
 - Postprandial
 - Short lived
 - Asymptomatic
 - No nocturnal symptoms

- Pathologic GERD
 - Symptoms
 - Mucosal injury
 - Nocturnal symptoms

Reflux Esophagitis

- **Esophagitis is rarely caused by agents other than reflux**
- Acute esophagitis may be caused by:

Infective agents:

- Fungal infection (mainly by *Candida albicans*) is common
- Viral infections of the esophagus (particularly by herpes simplex and cytomegalovirus) are seen in AIDS patient
- Bacterial infection is very rare

or

Physical agents: irradiation

Chemical: Ingestion of caustic agent

Epidemiology

- About 44% of the US adult population have heartburn at least once a month
- 14% of Americans have symptoms weekly
- 7% have symptoms daily

Epidemiology

- Prevalence of Symptoms of Gastroesophageal Reflux in a Cohort of Saudi Arabians: A Study of 1265 Subjects
- The mean age was 29.97 ± 11.58 years. Females formed 67.81% of the respondents and 62.73% had one or more episodes of heartburn per week.
- The prevalence of GERD in the surveyed population was 45.4%. GERD was more prevalent in older individuals (mean age 31.9 vs. 30.0 years) and in those with a higher BMI

– Saudi J Gastroenterol. 2014 Jul-Aug; 20(4): 248–254.

Pathophysiology

- Abnormal lower esophageal sphincter
- or
- Increase abdominal pressure

GERD

Pathophysiology

A. Abnormal lower esophageal sphincter

1. Functional (frequent transient LES relaxation)
2. Mechanical (hypotensive LES)
3. Foods (eg, coffee, alcohol, smoking)
4. Medications (eg, calcium channel blockers),
5. Location hiatal hernia

The most common cause of (GERD).


decrease the pressure of the LES.

Hiatal hernia present in ~70% of people with GERD

• or

B. Increase abdominal pressure

Obesity
Pregnancy present in ~80%
Increased gastric volume




Clinical Manifestations

- Most common symptoms
 - Heartburn—retrosternal burning discomfort
 - Regurgitation—effortless return of gastric contents into the pharynx without nausea, retching, or abdominal contractions



Atypical symptoms....coughing, chest pain, and wheezing.





Diagnostic Evaluation

- If classic symptoms of heartburn and regurgitation exist, the diagnosis of GERD can be made clinically and treatment can be initiated

Esophagogastroduodenoscopy

- Endoscopy (with biopsy if needed)
 - In patients with unusual signs/symptoms
 - Those who fail a medication trial
 - Those who require long-term tx


Esophagogastroduodenoscopy
(EGD or upper endoscopy)



pH


- 24-hour pH monitoring
 - Accepted standard for establishing or excluding presence of GERD for those patients who do not have mucosal changes
 - Trans-nasal catheter or a wireless capsule shaped device

Morphology



Simple hyperemia

Morphology of GERD



Eosinophils and neutrophils

basal zone hyperplasia
>20% of total epithelial
thickness

Elongation of lamina propria papillae

Reflux Esophagitis

- The severity of symptoms is not closely related to the degree of histologic damage
- The degree of histologic damage tends to increase with disease duration.

Treatment

- H₂ receptor Blockers
- Proton pump inhibitors

reduce gastric acidity
typically provides
symptomatic relief



Antireflux surgery

Complications

- Erosive esophagitis
- Stricture
- Barrett's esophagus


Complications

- Erosive esophagitis
 - Responsible for 40-60% of GERD symptoms
 - Severity of symptoms often fail to match severity of erosive esophagitis
 - Red mucosa with erosions leading to hematemesis and melena



Complications

- Esophageal stricture
 - Result of healing of erosive esophagitis and lead to dysphagia
 - May need dilation



Complications








8-15%

- Barrett's Esophagus
- Definition:
 - Intestinal metaplasia within the esophageal squamous mucosa

Main cause (Pathophysiology)

- Barrett's Esophagus
 - Acid damages lining of esophagus and causes chronic esophagitis
 - Damaged area heals in a metaplastic process and abnormal columnar cells replace squamous cells
 - Associated with the development of adenocarcinoma




Many patients with Barrett's are asymptomatic

Morphology





- **Endoscopic image of Barrett's esophagus: An area of red mucosa**




- **Barrett's esophagus is marked by the presence of columnar epithelia in the lower esophagus, replacing the normal squamous cell epithelium**


Barrett esophagus



Summary



Barrett esophagus



- Complications: Dysplasia and adenocarcinoma

- **The most common malignant tumors of the esophagus are squamous carcinomas and adenocarcinomas**
- The prognosis for both types of carcinoma is poor

Squamous carcinomas

are most common in the middle and lower esophagus. They mostly develop in men who are heavy alcohol drinkers or heavy smokers, and may be preceded by epithelial dysplastic change.

- Not related to GERD



Case scenario: A man with retrosternal pain

- A 57-year-old presents with a history of a retrosternal burning sensation, particularly after large meals, and often on retiring to bed at night. Treatment with antacids has had little effect and he has been referred by his GP for endoscopy.
- Upper gastrointestinal tract endoscopy reveals reddening of the lower esophageal mucosa from the level of the gastroesophageal junction to a point 32 cm from the incisors. There is no evidence of a hiatus hernia. The proximal border of the reddened area is irregular, and this area is biopsied. The biopsy shows gastric and intestinal-type glandular mucosa.

1. What is the likely cause of the symptoms?

- The symptoms of 'heartburn' are suggestive of gastroesophageal reflux disease (GERD), with or without the presence of a hiatus hernia.
- Other important causes of retrosternal pain should not be overlooked, including cardiovascular causes, especially myocardial ischaemia, as well as other rarer causes including pneumothorax and musculoskeletal pain.

2. What is the final diagnosis?


- The endoscopic and biopsy appearances confirm a Barrett's oesophagus. This is a metaplastic process which develops as a result of persistent reflux of gastric contents into the esophagus, the normal squamous mucosa being replaced by glandular mucosa of gastric or intestinal type

3. What further information do you require from the biopsy report?

It is important to look for dysplastic change in the biopsy which may herald the development of adenocarcinoma.


4. What are the major causes of reflux esophagitis?

- Reflux of gastric contents is the major cause of reflux esophagitis. Many factors play a role:
 - (a) the presence of a sliding hiatal hernia is the most common
 - (b) heavy alcohol use
 - (c) heavy tobacco use
 - (d) increased gastric volume
 - (e) decreased efficacy of LES
 - (f) pregnancy
 - (g) CNS depressants
 - (h) hypothyroidism



hiatus hernia

An X-ray film of the chest shows a retrocardiac, gas-filled structure.





CT scan of the abdomen with contrast was done which showed marked distention and distortion of the stomach with herniation of a portion of the stomach into the lower thorax, representing a combination of hernia through the esophageal hiatus hernia


5. What are other causes of esophagitis?

Ingestion of irritants (eg, alcohol, corrosive acids); infections in immunosuppressed hosts by fungi (eg, *Candida*) or viruses (eg, CMV, herpes); uremia; radiation therapy; graft-versus-host disease; and cytotoxic anticancer therapy.

6. What are the gross and microscopic features of reflux esophagitis?


Gross features:

- Simple **hyperemia**
- Erosion
- Ulceration
- Stricture
- Development of Barrett esophagus
- Development of mass:
adenocarcinoma




Microscopic features of reflux esophagitis


1. basal zone hyperplasia,




2. Eosinophils and neutrophils




3. Elongation of lamina propria papillae






Barrett esophagus without dysplasia



Barrett esophagus with dysplasia



Adenocarcinoma in Barrett Esophagus



Mass , need biopsy
adenocarcinoma

7. What are the major complications of reflux esophagitis?

The potential complications of severe reflux esophagitis are

- (a) Ulcer
- (b) Bleeding
- (c) Development of stricture
- (d) Development of Barrett esophagus and adenocarcinoma.

complications of reflux esophagitis

-development of Barrett esophagus and adenocarcinoma

