



PLACENTA



Dr Jamila ELmedany






- **It is a Fetomaternal structure.**
- Formed by the beginning of the **4th month.**
- It is the primary site for exchange of gases and nutrients between the mother and the fetus.

Formation of Placenta



- **Fetal Part:**
- **Villous Chorion.**
- It is the bushy area at the embryonic pole
- Its villi are more in number, enlarged and branch profusely.
- **Maternal Part:**
- **Decidua Basalis** (part of the decidua deep to the conceptus).



- **Decidua (Gravid Endometrium) :**
- **It is the functional layer of the endometrium during pregnancy which is shed after parturition.**

FULL TERM PLACENTA


Discoid in shape.

Weighs (500 – 600)g.



Has two surfaces:

Fetal

Maternal




Fetal Surface




- 1. **Smooth** because it is covered with the amnion.
- 2. **The umbilical cord** is attached to its center.
- 3. The **chorionic vessels** are radiating from the umbilical cord.

Maternal Surface




- **Rough.**
- Formed of (15 –20) irregular convex areas (**Cotyledons**)
- which are separated by grooves (**placental septa**).
- Each cotyledon is covered by a thin layer of **decidua basalis**.

Structure of a Cotyledon




- It consists of two or more stem villi with their many branch villi.
- It receives (80-100) maternal spiral arteries that enter the intervillous spaces at regular intervals.

INTERVILLOUS SPACE




- large blood filled spaces which are freely communicating
- They receive spiral arteries from the lacunae in the syncytiotrophoblast.
- The spaces are drained through endometrial veins.
- Both arteries and veins pass through pores in the cytotrophoblastic shell.

Fetal Placental Circulation




- **Two Umbilical Arteries:**
- Carry poorly oxygenated blood from the fetus to the placenta.
- within the branch chorionic villi, they form:
- **Arterio-capillary venous network:**
- It brings the fetal blood extremely close to the maternal blood.
- The well oxygenated fetal blood in the capillaries passes into veins accompanying the chorionic arteries.
- At the umbilical cord, they form the **One Umbilical Vein.**

Maternal Placental Circulation




- **80 –100 spiral endometrial arteries** discharge into the intervillous space.
- The blood is propelled in jet like fountains by the maternal blood pressure.
- The pressure of this entering blood is higher than that in the intervillous space.
- It forms a roof of the space.
- As the pressure dissipates, the blood flows slowly around the branch villi.




- Exchange of metabolites and gases with the fetal blood.
- As the pressure decreases, the blood flows back from the chorionic plate and enter the **endometrial veins** to the maternal circulation.

PLACENTAL MEMBRANE




- It is a composite thin membrane of extra fetal tissues which **separates** the fetal and maternal bloods.
- **Up to (20) weeks, it is composed of (4) layers:**
- **Syncytiotrophoblast.**
- **Cytotrophoblast.**
- **Connective tissue of the villus.**
- **Endothelium of fetal capillaries.**

PLACENTAL MEMBRANE




- At full term
- It becomes thinner and composed of (3) layers only:
- 1. Syncytiotrophoblast.
- 2. Connective tissue.
- 3. Endothelium of the capillaries.
- At some sites, the syncytio comes in direct contact with the endothelium of the capillaries and forms **Vasculosyncytial** placental membrane.


FUNCTIONS OF THE PLACENTA



- **1. Metabolic :**
- **Synthesis of: Glycogen, Cholesterol and Fatty Acids.**
- They supply the fetus with nutrients and energy.
- **2. Transportation of:**
- **(A) Gases:**
- Exchange of O₂, CO₂ and CO is through **simple diffusion.**
- The fetus extracts (20 –30) ml of O₂/minute from the maternal blood.



- **(B) Nutrients and Electrolytes:**
- Water, Amino acids, Carbohydrates, Vitamins and Free Fatty Acids are rapidly transferred to the fetus.
- **C. Maternal Antibodies:**
- Maternal immunoglobulin G gives the fetus passive immunity to some infectious diseases (**measles, small box**) and not to others (**chicken box**).



- **(D) Drugs and Drug metabolites:**
- They cross the placenta by simple diffusion.
- They can affect the fetus directly or indirectly by interfering with placental metabolism.
- **(E) Hormones:**
- **Protein hormones** do not reach the embryo in sufficient amounts.
- some of these hormones
- **(Thyroxine &**
- **Testosterone** which may cause masculinization of a female fetus) can cross the placental membrane.
- **F) Waste products:**
- Urea and uric acid pass through the placental membrane by simple diffusion.

- (3) Endocrine Synthesis:

- (1) Progesterone :

- Maintains pregnancy if the corpus luteum is not functioning well.

- (2) Estrogen

- Stimulates uterine growth and development of the mammary glands.

- 3) hCS or Hpl:


- A growth hormone that gives the fetus the priority on maternal blood glucose.
- It promotes breast development for milk production.

- (4) hCG:

- Maintains the corpus luteum and used as **indicator of pregnancy**.

- **Fetal drug addiction can be due to some drugs as Heroin.**
- **All sedatives and analgesics can affect the fetus to some degree.**
- **Drugs used for management of labor can cause respiratory distress to the newborn.**


ANOMALIES OF PLACENTA



- **1. Placenta Accreta:**
- Abnormal absence of chorionic villi with partial or complete absence of the decidua basalis.
- **2. Placenta Percreta:**
- Chorionic villi penetrate the myometrium to the perimetrium.
- The most common presenting sign of these two anomalies is trimester bleeding.
- **3. Placenta Previa:**
- The blastocyst is implanted close to or overlying the internal uterine os.
- It is associated with late pregnancy bleeding.
- Delivery is through Cesarean section.

Fate of Placenta

- The strong uterine contractions that continue after birth compress uterine blood vessels to limit bleeding & cause the placenta to **detach** from the uterine wall (within 15 minutes after birth of the infant).





THANK YOU