



The Uterine Corpus

Sufia Husain.
Pathology Department
KSU, Riyadh
April 2017

Reference: Robbins & Cotran Pathology and Rubin's Pathology



Uterus with bilateral tube and ovaries, posterior view




<http://pixgood.com>







http://www.humpath.com/IMG/jpg/uterus_03_1.jpg

VAGINA AND UTERUS



http://www.wikilectures.eu/images/thumb/8/82/Vagina_uterus.png/720px-Vagina_uterus.png



Lecture Outline

Lecture: Endometrial hyperplasia, uterine cancer and fibroids (leiomyomas).

At the end of this lecture, the student should know:

Lesions of endometrium of uterus: know the risk factors, clinical presentation, macroscopic and histological features of

- **Endometrial hyperplasia**
- **Endometrial carcinoma**

Lesions of myometrium of uterus:

- **Leiomyoma** : understand the pathology and clinical features of uterine leiomyomas and is aware that leiomyoma (fibroid) is the commonest neoplasm arising in the female genital tract.
- **Leiomyosarcoma**


Endometrial Hyperplasia



Endometrial Hyperplasia



- Endometrial hyperplasia is a process in which there is a proliferation of endometrial glands resulting in an increase in gland/stroma ratio when compared to normal.
- It is induced by persistent, prolonged estrogenic stimulation of the endometrium.
- The endometrial hyperplasia may progress to endometrial carcinoma.
- The development of cancer is based on the level and duration of the estrogen excess.
- The risk is of developing carcinoma depends on the severity of the hyperplastic changes and associated cellular atypia.



Endometrial Hyperplasia: causes



Causes of Endometrial Hyperplasia: **any condition in which there is high estrogen level can lead to endometrial hyperplasia e.g.**

- a. Anovulatory menstrual cycles (failure of ovulation).
- b. Excessive endogenous production of estrogen (by the body)
e.g. in
 - polycystic ovary syndrome (Stein-Leventhal syndrome),
 - granulosa cell tumors of the ovary
 - excessive ovarian cortical function (cortical stromal hyperplasia)
- c. Exogenous administration of estrogenic steroids without counter balancing progestins, over a long period of time.

Endometrial Hyperplasia: clinical



- Mild type of hyperplasia tends to occur in younger patients
- The great majority of mild hyperplasia regress, either spontaneously or after treatment.
- The more severe type of hyperplasia occur mainly in peri- and postmenopausal women. This form has a significant premalignant potential.
- Patients with endometrial hyperplasia usually present with abnormal uterine bleeding.

Endometrial hyperplasia



Endometrial Hyperplasia: classification

In endometrial hyperplasia there is proliferation of both glands and stroma but the proliferation of the glandular component is much more. Therefore over crowding of glands occur. Endometrial hyperplasia is histologically classified based on:

- 1) **Gland architecture:** into → **simple or complex**, depending on the degree of glandular complexity and crowding.
- 2) **Cytologic features:** into → **with or without atypia**.
(Atypia/pleomorphism = loss of polarity, vesicular nuclei, prominent nucleoli, rounded cells.)


Therefore the classification is as follows:

I. Simple hyperplasia

- Without atypia
- With atypia


II. Complex hyperplasia

- Without atypia
- With atypia






No hyperplasia and no atypia



quizlet.com


Hyperplasia with atypia



<http://www.indiacancersurgerysite.com/>

Simple hyperplasia without atypia

- ☞ Simple hyperplasia (cystic hyperplasia): glands are variably shaped and sized and cystically dilated with abundant cellular stroma and give a "Swiss Cheese" appearance.
- ☞ There is a mild increase in the gland-to-stroma ratio
- ☞ These lesions rarely progress to adenocarcinoma
- ☞ Simple hyperplasia may progress to cystic atrophy



Simple hyperplasia with atypia




- ❧ Uncommon
- ❧ It has the Architecture of simple hyperplasia, but there is cytologic atypia within the glandular epithelial cells
- ❧ 10% of such lesions progress to carcinoma

Complex hyperplasia without atypia




- ☞ Proliferation of endometrial glands resulting in complex crowded glands with papillary infoldings and irregular shapes. The crowded glands are back-to-back with very little intervening stroma.
- ☞ The epithelial cells remain cytologically normal.
- ☞ 3% progression to carcinoma



Complex hyperplasia with atypia



- ❧ Complex proliferation of endometrial glands (back-to-back irregular glands) with atypia.
- ❧ The nuclei show loss of polarity and are enlarged and rounded and may have irregular nuclear membranes
- ❧ Commonly about 30% of women with this diagnosis have carcinoma somewhere in the uterus when a hysterectomy is performed
- ❧ About 30% progress to carcinoma



Endometrial Hyperplasia: Clinical behavior and premalignant potential



- Some endometrial hyperplasia revert to normal spontaneously or with medical treatment, others persist as hyperplasia, and a few progresses to endometrial adenocarcinoma.
- Generally, patients who have hyperplasia with atypia are more likely to develop carcinoma than those without atypia. The risks for developing adenocarcinoma in each are as follows:
 - Simple hyperplasia without atypia — 1%
 - Complex hyperplasia without atypia — 3%
 - Simple hyperplasia with atypia (simple atypical hyperplasia) — 10%
 - Complex hyperplasia with atypia (complex atypical hyperplasia) — 30%
- Atypical hyperplasia in postmenopausal women appears to have a higher rate of progression to adenocarcinoma.

Endometrial Hyperplasia, Risk Factors



-
- ❧ Obesity
 - ❧ Western diet
 - ❧ Nulliparity
 - ❧ Diabetes Mellitus
 - ❧ Hypertension
 - ❧ Hyperestrinism




Endometrial adenocarcinoma


Endometrial adenocarcinoma

This is a common neoplasm in women. Overall it is the fifth commonest cancer in women.

- Endometrial cancers arise mainly in postmenopausal women
- They cause postmenopausal bleeding
- Early detection and cures are possible
- These tumors are classified into two broad categories:
 - **Type I carcinomas (also known as endometrioid carcinoma):** accounts for 80% of endometrial cancers. It is the most common type. e.g. → endometrioid adenocarcinoma and its variants.
 - **Type II carcinomas:** they are papillary serous carcinoma and clear cell carcinoma. Papillary serous is the more common form of type II carcinoma.



Endometrial Cancer




Type I endometrial carcinoma/ endometrioid carcinoma



- *Endometrioid carcinoma* is associated with estrogen excess and endometrial hyperplasia. The majority of the carcinomas are well differentiated.
- Risk factors for type I are the same as that of endometrial hyperplasia and include:
 - Endometrial hyperplasia is a precursor to endometrioid carcinoma
 - obesity (women with upper body fat have 3 times the risk of women with lower body fat),
 - estrogen therapy,
 - nulliparity (as a result of infertility due to chronic anovulation),
 - chronic anovulation,
 - late menopause,
 - hypertension,
 - diabetes,
 - tamoxifen therapy,
 - and high socioeconomic status.
 - The disease may follow atypical hyperplasia but may occur independently of it especially in older patients.


Usual sequence of events in Type I endometrioid carcinoma



Type I endometrioid carcinoma: genetics




- ❧ Mutations in the PTEN gene have been identified in 30% to 80% of endometrioid carcinomas
- ❧ There may be inactivation of DNA mismatch repair genes
- ❧ In the more poorly differentiated endometrioid carcinomas, mutations in *p53* can be found in up to 50% of cases




http://www.proteinatlas.org/images_dictionary/endometrial_cancer__1__figo_1__overview.jpg

Endometrial carcinoma



Type II endometrial carcinomas: Serous carcinoma

- ❧ *Serous carcinoma* arises in older women, with endometrial atrophy (small atrophic uterus).
- ❧ They occur in late in life, about one decade later than type I carcinoma
- ❧ There is no association with hyperestrinism or preexisting hyperplasia
- ❧ They represent 15% of cases of all endometrial carcinoma
- ❧ Mutations in *p53* are present in at least 90% of serous endometrial carcinoma
- ❧ The precursor of serous carcinoma is **endometrial intraepithelial carcinoma** (like carcinoma in situ)
- ❧ These tumors are large bulky poorly differentiated tumors which invade early into the myometrium and have a poor prognosis. Extrauterine extension is common.



CHARACTERISTICS OF TYPE I AND TYPE II ENDOMETRIAL CARCINOMAS

FEATURES	TYPE I	TYPE II
HISTOLOGIC TYPE	Endometrioid adenocarcinoma	Serous or clear cell carcinoma
AGE	Premenopausal and perimenopausal (50-60 yrs)	Post menopausal (~ 70 yrs)
UNOPPOSED ESTROGEN	Present	Absent
PRECURSOR LESION	Hyperplasia with atypia	Endometrial intraepithelial carcinoma
GROWTH	Slow growing	Rapidly progressing
GRADE	Low	High
MYOMETRIAL INVASION	Usually superficial	Usually deep
PROGNOSIS	Favorable	Poor
GENETIC ALTERATIONS NOTED	PTEN, microsatellite instability	P53 mutations

Endometrial adenocarcinoma: clinical features



- Most patients are between 50 and 60 years.
- Many of the patients tend to be nulliparous and obese.
- Endometrial adenocarcinoma manifests as abnormal vaginal bleeding and excessive leucorrhea.
- Elderly women present with postmenopausal bleeding.
- The diagnosis of endometrial cancer must be confirmed by biopsy or curettage and histologic examination of the tissue.



http://upload.wikimedia.org/wikipedia/commons/0/00/Endometrial_hyperplasia.jpg

Endometrial carcinoma: basic morphology



☞ Grossly:

- May look close to normal or exophytic or infiltrative

☞ Microscopy:

- Both type I and II are adenocarcinomas.
- In both cases tumors originate in the endometrium and can eventually infiltrate the underlying myometrium, enter vascular spaces and metastasize to lymph nodes.
- Serous carcinoma has much greater cytologic atypia and are more poorly differentiated and is therefore more aggressive

☞ Tumor spreads by:

- Direct myometrial invasion with extension to the periuterine structures
- Through lymphatics to lymph nodes
- In the late stages, metastasize to the lungs, liver, bones, others




Endometrial adenocarcinoma: prognosis



-
- Clinical behavior of endometrial adenocarcinoma depends on the histologic type, the grade (degree of differentiation) and the stage (extent of spread).
 - Endometrioid carcinoma (type I) has a better prognosis than the other histologic types.
 - Serous carcinomas (type II) have poorer prognosis
 - Stage is the major determinant of survival.

How endometrial carcinoma can spread

(stages 1, 2 and 3 of endometrial carcinoma)



"Diagram showing stage 1A and 1B, 2 and 3A to 3C cancer of the womb CRUK 196, 206 and 224" by Cancer Research UK - Original email from CRUK. Licensed under CC BY-SA 4.0 via Wikimedia Commons - http://commons.wikimedia.org/wiki/File:Diagram_showing_stage_1A_and_1B,_2_and3A_to_3C_cancer_of_the_womb_CRUK_206.svg#/media/File:Diagram_showing_stage_1A_and_1B,_2_and_3A_to_3C_cancer_of_the_womb_CRUK_196_and_206_and_224.svg

LEIOMYOMA



Leiomyoma (fibroid) of uterus



- Leiomyoma is a benign tumor of smooth muscle origin.
- It is the most common neoplasm of the female genital tract and probably the most common neoplasm in women.
- The tumor is estrogen responsive. Estrogen stimulates their growth. Leiomyomas often increases in size during pregnancy and decrease in size after menopause.
- About 40% of leiomyomas have an associated chromosomal abnormality
- They are benign tumors with no appreciable malignant potential (incidence of malignant transformation to sarcoma is 0.1-0.5%).

Leiomyoma (fibroid) of uterus



Clinical features



- It can be single or multiple. Mostly it is multiple.
- Irregular abnormal bleeding and sometimes pelvic pain.
- It may cause anemia from heavy bleeding.
- Can have urinary frequency if the fibroid is compressing the urinary bladder.
- It may interfere with implantation and therefore cause infertility.
- In pregnant women it may cause abortion, obstructed labor, post partum hemorrhage etc.
- *Alternatively it maybe* entirely asymptomatic

Uterine Leiomyoma/ Fibroid




Leiomyoma may be located anywhere in the myometrium.

- **Submucosal** tumors are present immediately below the endometrium.
- **Intramural** tumors, the most common, lie within the myometrium.
- **Subserosal** fibroids lie beneath the serosal surface of the uterus or are pedunculated and attached to the serosa.
- Pedunculated ones may lose their connection to the uterus forming a "parasitic leiomyoma".



Leiomyoma gross:

- Well circumscribed, spherical, dense and firm-to-hard masses.
- Cut section shows whorled, tan-white cut surfaces.



Leiomyoma: Microscopically, there are interlacing bundles of smooth muscle cells with collagenous stroma between bundles. The individual muscle cells are uniform in size and shape. They have the characteristic oval to elongated nucleus. Mitotic figures are scarce.




Leiomyoma - histology

A histological micrograph of a leiomyoma stained with hematoxylin and eosin (H&E). The image shows a dense field of spindle-shaped smooth muscle cells arranged in interlacing fascicles. The nuclei are uniform in size and shape, and the overall appearance is that of a benign smooth muscle tumor.

**Uniform, bland spindled cells
Fascicular arrangement**

<http://pixgood.com/leiomyoma-histology.html>



<http://pixgood.com/myometrium.html>

Leiomyosarcoma



- ❧ It is the malignant tumor of the smooth muscle.
- ❧ It is rare.
- ❧ Sites include the uterus and soft tissue
- ❧ Poor prognosis.

