



Drugs In OVULATION INDUCTION




Drugs In OVULATION INDUCTION



ILOs

By the end of this lecture you will be able to:

- ② Recall how ovulation occurs and specify its hormonal regulation
- ② Classify ovulation inducing drugs in relevance to the existing deficits
- ② Expand on the pharmacology of each group with respect to mechanism of action, protocol of administration, indication, efficacy rate and adverse effects.



Ovulation Induction

1. Antiestrogens

SERMs;
Clomiphene
Tamoxifen

2. GnRH

GnRH-agonists
Leuprolin
Goserelin

4. Rx Hyperprolactinaemia

D₂ R agonists

Bromocriptine


3. Gonadotrophins

HMGs; Menotropin
HCGs; Pregnyl

5. Rx POLYCYSTIC OVARIAN SYNDROME (PCOS)

Most common cause of infertility
Insulin resistance may play a role ???

Metformin



Normogonadotrophic



ANTIESTROGENS


1. CLOMIPHENE

Pharmacological effects

- Compete with estrogen on the hypothalamus and anterior pituitary gland; ↓ negative feed back of endogenous estrogen → ↑GnRH → ↑production of FSH & LH → **OVULATION**


Indication

- Female infertility; due to anovulation or oligoovulation . not due to ovarian or pituitary failure → **Normogonadotrophic**
- The success rate for ovulation → 80% & pregnancy → 40% .



Method of administration

- Clomiphene given → 50 mg/d for 5 days from 5th day of the cycle to the 10th day.
- If no response give 100 mg for 5 days again from 5th to 10th day
- Each dose can be repeated not more than 3 cycles .




ADRs

1. Hot Flashes & breast tenderness
2. Gastric upset (nausea and vomiting)
3. Visual disturbances (reversible)
4. ↑ nervous tension & depression
5. Skin rashes
6. Fatigue
7. Weight gain
8. Hair loss (reversible)
9. Hyperstimulation of the ovaries & high incidence of multiple birth.

2. TAMOXIFEN

Is similar & alternative to clomiphene
But differ in being **Non Steroidal**

- Tamoxifen is a good alternative to clomiphene in women with PCOS and clomiphene-resistant cases
- Used in palliative treatment of estrogen receptor- positive breast cancer.




Ovulation Induction

1. Antiestrogens

SERMs;
Clomiphene
Tamoxifen

2. GnRH

GnRH-agonists
Leuprolin
Goserelin



2. GONADOTROPIN RELEASING HORMONE (GnRH)

Uses:


Induction of ovulation in patients with hypothalamic amenorrhea (GnRH deficient)

Analogues with agonist activity:

Leuprolin, Goserelin


- GnRH and agonists, given S.C. in a **pulsatile (drip)** to stimulate gonadotropin release (1 – 10 μg / 60 – 120 min)
Start from day 2-3 of cycle up to day 10

- Given **continuously**, when gonadal suppression is desirable e.g. precocious puberty and advanced breast cancer in women and prostatic cancer in men



ADRS OF GnRH Agonists

- **GIT disturbances, abdominal pain, nausea....etc**
- **Headache**
- **Hypogonadism *on long term use* ➔**
 - ◆ **Hot flashes**
 - ◆ **↓Libido**
 - ◆ **Osteoporosis**
 - ◆ **Rarely ovarian hyperstimulation ➔ (ovaries swell & enlarge)**



Ovulation Induction

1. Antiestrogens


SERMs;
Clomiphene
Tamoxifen

2. GnRH

GnRH-agonists
Leuprolin
Goserelin

3. Gonadotrophins

HMGs; Menotropin
HCGs; Pregnyl



3. GONADOTROPHINS

[FSH & LH]

Are naturally produced by the pituitary gland

For therapeutic use, extracted forms are available as;

1. Human Menopausal Gonadotrophin(hMG)→ extracted from postmenopausal urine → contains LH & FSH → **MENOTROPIN**
2. Human Chorionic Gonadotrophin(hCG) extracted from urine of pregnant women → contains mainly LH) → **PREGNYL**

Indication

- Stimulation & induction of ovulation in infertility 2ndry to gonadotropin deficiency (**pituitary insufficiency**)

Success rate for inducing ovulation is usually $\geq 75\%$

GONADOTROPHINS

Method of administration

hMG is given i.m every day starting at day 2-3 of cycle for 10 days followed by **hCG** on (10th - 12th day) for OVUM RETRIEVAL .

ADRs

FSH containing preparations; Fever
Ovarian enlargement (hyper stimulation)
Multiple Pregnancy (approx. 20%)

LH containing preparations; Headache & edema




Figure 81-2. Plasma concentrations of the gonadotropins and ovarian hormones during the normal female sexual cycle.

4. Hyperprolactinaemia

D₂ R Agonists

BROMOCREPTINE

Is an ergot derivative (not a hormone)

Mechanism

D₂ R Agonists binds to dopamine receptors in the anterior pituitary gland & inhibits prolactin secretion


Indications

➤ Female infertility 2ndry to hyperprolactinaemia

ADRs

- GIT disturbances; nausea, vomiting, constipation
- Headache dizziness & orthostatic hypotension
- Dry mouth & nasal congestion
- Insomnia

Bromocriptine



Hyperprolactinaemia

No Ovulation

Drugs In OVULATION INDUCTION

G
L
U
O
C
O
K
D

