# **ANATOMY OF THE BRAIN STEM**

# (EXTERNAL FEATURES)

By

Prof. Ahmed Fathalla& Dr.Sanaa Alshaarawy

#### **OBJECTIVES**

#### At the end of the lecture, students should:

- List the components of brain stem.
- Describe the site of brain stem.
- Describe the relations between components of brain stem & their relations to cerebellum.
- Describe the external features of both ventral & dorsal surfaces of brain stem.
- List cranial nerves emerging from brain stem.
- Describe the site of emergence of each cranial nerve.

## **DEVELOPMENT OF BRAIN**

- The brain develops from the cranial part of neural tube.
- The cranial part divides into 3 parts:




Figure 18 - 20. Diagrammatic sketches of the brain vesicles, indicating the adult derivatives of their walls and cavities. \*The rostral part of the third ventricle forms from the cavity of the telencephalon; most of this ventricle is derived from the cavity of the diencephalon.

#### **FOREBRAIN:** subdivides into:

- 1-Two cerebral hemispheres (cavities: 2 lateral ventricles).
- 2-Diencephalon (cavity: 3rd ventricle):
- thalamus, hypothalamus, epithalamus & subthalamus


MIDBRAIN (cavity: cerebral aqueduct).


HINDBRAIN (cavity: 4th ventricle): subdivides into


- 1-Pons.
- 2-Cerebellum.
- 3- Medulla oblongata.

#### **BRAIN STEM**

- ☐ The brainstem is the region of the brain that connects the cerebrum with the spinal cord
- ☐ SITE:
- It lies on the basilar part of occipital bone (clivus).
- ☐ PARTS: From above downwards:
- Mid brain, pons & medulla oblongata
- ☐ CONNECTIONS WITH CEREBELLUM:
- Each part of brain stem is connected to cerebellum by cerebellar peduncles (superior, middle & inferior).










### **FUNCTIONS OF BRAIN STEM**


- 1. Pathway of tracts between cerebral cortex & spinal cord.
- 2. Site of origin of <u>nuclei</u> of cranial nerves (from 3<sup>rd</sup> to 12<sup>th</sup>).
- Site of emergence of <u>cranial</u> nerves (from 3<sup>rd</sup> to 12<sup>th</sup>).
- 4. Contains groups of nuclei & related fibers known as reticular formation responsible for: control of level of consciousness, perception of pain, regulation of cardiovascular & respiratory systems.








# **BRAIN – VENTRAL SURFACE**




#### **MEDULLA – VENTRAL SURFACE**

#### ■ Ventral median fissure:\*

- Continuation of <u>ventral</u> <u>median fissure</u> of spinal cord
- Divides the medulla into 2 halves
- Its lower part is marked by decussation of most of pyramidal (corticospinal) \* fibers (75%-90%).
- □ Pyramid: \*
- An elevation, lies on either side of ventral median fissure
- Produced by corticospinal tract.




- ☐ Olive: \*
- An elevation, lies lateral to the pyramid.
- Produced by inferior olivary nucleus (important in control of movement).
- □ Nerves emerging from Medulla (4 nerves):
- Hypoglossal (12<sup>th</sup>): from sulcus between pyramid & olive
- Glossopharyngeal (9<sup>th</sup>), vagus (10<sup>th</sup>) & cranial part of accessory (11<sup>th</sup>): from sulcus dorsolateral to olive (from above downwards)




#### **PONS – VENTRAL SURFACE**

- Basilar sulcus: \*
- Divides the pons into 2 halves, occupied by basilar artery.
- ☐ Transverse pontine (pontocerebellar) fibers: \*
- Originate from pontine nuclei, cross the midline & pass through the contralateral middle cerebellar peduncle to enter the opposite cerebellar hemisphere.




- ☐ Nerves emerging from Pons (4 nerves):
- Trigeminal (5th): from the middle of ventrolateral aspect of pons, as 2 roots: a small medial motor root & a large lateral sensory root.
- Abducent (6<sup>th</sup>): from sulcus between pons & pyramid.
- vestibulocochlear (8<sup>th</sup>): at cerebellopontine angle (junction between medulla, pons & cerebellum). Both nerves emerge as 2 roots: from medial to lateral: motor root of 7<sup>th</sup>, sensory root of 7<sup>th</sup> vestibular part of 8<sup>th</sup> & cochlear part of 8<sup>th</sup>




### MID BRAIN - VENTRAL SURFACE

- large column of descending fibers (crus cerebri or basis pedunculi), on either side, separated by a depression called the interpeduncular fossa\*
- Nerve emerging from Midbrain (one):
- Occulomotor (3<sup>rd</sup>): from medial aspect of crus cerebri.



## **MEDULLA – DORSAL SURFACE**

The features differ in the caudal part **(closed** medulla) and the cranial part (open medulla).




open medulla

closed medulla




- Dorsal median sulcus: divides the closed medulla into 2 halves.
- Fasciculus gracilis: on either side of dorsal median sulcus.
- Gracile tubercle: an elevation produced at the upper part of fasciculus gracilis, marks the site of gracile nucleus.
- Fasciculus cuneatus: on either side of fasciculus gracilis.
- Cuneate tubercle: an elevation produced at the upper part of fasciculus cuneatus, marks the site of cuneate nucleus.

# CLOSED MEDULLA




- Cavity: 4<sup>th</sup> ventricle
- On either side, an inverted V-shaped sulcus divides the area into 3 parts (from medial to lateral):
- 1. Hypoglossal triangle\*: overlies hypoglossal nucleus.
- 2. Vagal triangle\*: overlies dorsal vagal nucleus.
- 3. Vestibular area\*: overlies vestibular nuclei.

#### **OPEN MEDULLA**





### **PONS – DORSAL SURFACE**

- Separated from open medulla by an imaginary line passing between the margins of middle cerebellar peduncle.
- On either side of median sulcus, it divides into 2 parts (from medial to lateral):
- Medial eminence & facial colliculus\*: overlies abducent nucleus.
- Vestibular area:\*
  overlies vestibular
  nuclei.




The dorsal surfaces of open medulla and pons <u>lie</u> in the caudal 1/3<sup>rd</sup> and the rostral 2/3<sup>rd</sup> of the <u>floor of the 4<sup>th</sup> ventricle</u> respectively.





## MID BRAIN - DORSAL SURFACE

- Marked by 4 elevations:
- Two superior colliculi: concerned with visual reflexes.
- 2. Two inferior colliculi: forms part of auditory pathway.
- □ Nerve emerging from Midbrain (one):
- Trochlear (4<sup>th</sup>): just
  <u>caudal</u> to <u>inferior</u>
  <u>colliculus</u> (The only
  <u>cranial nerve</u>
  <u>emerging from dorsal</u>
  <u>surface of brain stem</u>).



# THANK YOU

### **SUMMARY**

☐ The brain stem is composed (from above downwards) of: midbrain, pons & medulla oblongata which are continuous with each other, with diencephalon above & with spinal cord below. ☐ The brain stem is connected with cerebellum through three pair of cerebellar peduncles. ☐ The brain stem is the site of cranial nuclei, the pathway of important ascending & descending tracts & the site of emergence of cranial nerves (from 3<sup>rd</sup> to 12<sup>th</sup>). ☐ Cranial nerves (with the exception of 4<sup>th</sup>) emerge from ventral surface of brain stem.

## **QUESTION 1**

- ☐ The cranial nerve that emerges from dorsal surface of midbrain is:
- 1. Occulomotor (3<sup>rd</sup>).
- 2. Trochlear (4<sup>th</sup>).
- 3. Abducent (6th).
- 4. Facial (7<sup>th</sup>).

# **QUESTION 2**

- ☐ Regarding the medulla oblongata:
- 1. The pyramid is lateral to olive.
- 2. The hypoglossal nerve is the most lateral nerve emerging from it.
- 3. The cuneate tubercle is lateral to gracile tubercle.
- 4. The cerebellum is connected to it by middle cerebellar peduncle.