


Objectives

- By the end of the lecture the student should be able to:
- List the parts of the ear: External, Middle (tympanic cavity) and Internal (labyrinth).
- Describe the parts of the external ear: auricle and external auditory meatus.
- Identify the boundaries of the middle ear: roof, floor and four walls (anterior, posterior, medial and lateral).

Objectives


- Define the contents of the tympanic cavity:
- I. Ear ossicles,: (malleus, incus and stapes)
- II. Muscles, (tensor tympani and stapedius).
- III. Nerves (branches of facial and glossopharyngeal).
- List the parts of the inner ear, bony part filled with perilymph (Cochlea, vestibule and semicircular canals), in which is suspended the membranous part that filled with endolymph).
- List the organs of hearing and equilibrium.


EXTERNAL EAR



It is formed of the auricle, & the external auditory meatus.

- The Auricle has a characteristic shape and collects air vibrations.
- It consists of a thin plate of elastic <u>cartilage</u> covered by a double layer of skin.
- It receives the insertion of extrinsic muscles, which are supplied by the facial nerve. Sensation is carried by great auricular & auriculotemporal nerves.







- The external auditory canal is a curved S-shaped tube about 2.5 cm, that conducts & collects sound waves from the auricle to the tympanic membrane. Its outer 1/3rd is elastic cartilage, while its inner 2/3^{rds} are boney.
- It is lined by skin, and its outer 1/3rd is provided with hairs, sebaceous and Ceruminous Glands: (modified sweat glands that secrete a yellowish brownish substance called the ear wax).

5


MIDDLE EAR (TYMPANIC CAVITY)





- Middle ear is a narrow, oblique, slit- like cavity (air-filled) in the petrous temporal bone & lined with mucous membrane.
- It contains the <u>auditory ossicles</u>, which transmit the vibrations of the tympanic membrane (eardrum) to the internal ear.

MIDDLE EAR (TYMPANIC CAVITY)





Communicates Anteriorly

- with the Nasopharynx through the Auditory Tube, which extends from the anterior wall downward, forward, and medially to the nasopharynx).
- The posterior 1/3rd of the canal is bony, and its anterior 2/3rds are cartilaginous.
- Its function is to equalize the pressure on both sides of the ear drum.

The middle ear has:

- Roof,
- Floor,
- and 4 walls:
- Anterior,
- Posterior,
- Lateral, and
- Medial.





The Floor is formed by a thin plate of bone, which separates the middle ear from the bulb of the internal jugular vein.


The anterior wall is formed below by a thin plate of bone that separates tympanic cavity from the internal carotid artery.

There are 2 canals at the upper part of the anterior wall.

The upper smaller is the canal for the tensor tympani muscle.

The lower larger is for the auditory tube.

Anterior wall





has in its Upper part a large, irregular opening, the aditus to the mastoid antrum.

Below: a small,
hollow, conical
projection, the
pyramid, which
houses the
stapedius muscle
and its tendon.

The tendon
emerges from the
apex of the
pyramid.

Posterior wall






The lateral wall:

- Is largely formed by the tympanic membrane.
- The membrane is obliquely placed, facing downward, forward, & laterally.
- It is extremely sensitive to pain.
- Nerve supply of ear drum:
- Outer surface:
- 1- Auriculotemporal nerve.
- 2- Auricular branch of vagus.
- Inner surface:
- Tympanic branch of the glossopharyngeal nerve.


TYMPANIC MEBRANE



- It is <u>concave</u> laterally, and at the depth of its concavity there is a small depression, "the <u>Umbo</u>" produced by the tip of the handle of the malleus.
- When the membrane is illuminated through an otoscope, the concavity produces a "Cone of Light," which radiates anteriorly and inferiorly from the umbo.
- Most of the of the membrane is tense and is called the Pars Tensa.
- A small triangular area on its upper part is slack and called the Pars Flaccida


- Greater part of the medial wall shows a rounded projection, called promontory, that results from the underlying 1st turn of the cochlea.
- Above and behind the promontory lies the oval window (Fenestra Vestibuli), which is closed by the base of the stapes. Below and behind the promontory lies the round window (Fenestra Cochleae), which is closed by the secondary tympanic membrane.

Medial wall



It is formed by the lateral wall of the inner ear.


Auditory Ossicles




The auditory ossicles are (3) malleus, incus, and stapes.

They transmit sound waves from tympanic membrane to the perilymph of the internal ear.

They are covered by mucous membrane & articulated by synovial joints.




- Origin: Cartilage of the auditory tube and the bony walls of its own canal.
- Insertion: into the handle of the malleus.
- Nerve supply: Mandibular nerve.
- Action: Contracts reflexly in response to loud sounds to limit the excursion of the tympanic membrane.



- Origin: Internal walls of the hollow pyramid.
- Insertion: The tendon emerges from the apex of the pyramid and is inserted into the neck of the stapes.
- Nerve supply: Facial nerve.
- Action: Reflexly damps down the vibrations of the stapes by pulling on the neck of that bone.

NERVES IN MIDDLE EAR



Tympanic nerve

- It is a branch of the glossopharyngeal nerve.
- It gives:
- Tympanic plexus on the promontory
- The tympanic plexus gives the,
- Lesser petrosal
 nerve which relays
 in the otic
 ganglion.
- It gives
 secretomotor
 supply to the
 parotid gland

Enters through the

Internal acoustic meatus with the 8th nerve.


It expands to form Geniculate ganglion.

It passes vertical behind the pyramid.

It leaves the middle ear

through the stylomastoid foramen.

FACIAL NERVE



BRANCHES OF FACIAL NERVE

1. <u>Greater Petrosal</u> nerve.

Arises from Geniculate Ganglion.

Carries preganglionic parasympathetic to supply:

Lacrimal,


Nasal

Palatine glands.

2. Nerve to Stapedius.

3. Chorda Tympani:


Arises just before the facial nerve exits.




Anterior semicircular canal Cochlea Vestibulocochlear nerve [VIII] Posterior semicircular canal Lateral semicircular canal


INTERNAL EAR, OR LABYRINTH

Labyrinth is situated in the petrous part of the temporal bone, medial to the middle ear.




Labyrinth






It consists of:


- Bony labyrinth: a series of bony chambers lined by endosteum.
- They contain a clear fluid, the perilymph, in which is suspended the membranous labyrinth.
- Membranous labyrinth: consists of a series of membranous sacs and ducts within the bony labyrinth, It is filled with endolymph.




- Its first turn
 produces the
 promontory
 on the medial
 wall of the
 tympanic
 cavity.
- It contains
 the cochlear
 duct (part of
 the
 membranous
 labyrinth).


Cochlea









In the lateral wall of the vestibule are the fenestra vestibuli, which is closed by the base of the stapes, and the fenestra cochleae, which is closed by the secondary tympanic membrane.






<u>The Membranous Labyrinth</u> consists of (**Four ducts & Two sacs**) Which are freely communicate with one another:

- Sacs: Utricle & Saccule (lodged in the bony vestibule).
- Ducts: Three semicircular Ducts, (lie within the bony semicircular canals),
- Cochlear Duct: (lies within the bony cochlea). The cochlear duct divides the bony cavity into Scala Vestibuli and Scala tympani.




The perilymph within the scala vestibuli is separated from the middle ear by the base of the stapes at the fenestra vestibuli.

The perilymph in the scala tympani is separated from the middle ear by the secondary tympanic membrane at the fenestra cochleae.



Located on the walls of the **utricle and saccule** are specialized sensory receptors, which are sensitive to the orientation of the head to gravity or other acceleration forces.

The utricle, saccule and semicircular ducts are concerned with maintenance of Equilibrium



THANK YOU