



•DEVELOPMENT OF VERTEBRAL COLUMN & SPINAL CORD

***Prof. Ahmed Fathalla Ibrahim
Professor of Anatomy
College of Medicine
King Saud University
E-mail: ahmedfathala@gmail.com***

OBJECTIVES

At the end of the lecture, students should be able to:

- List the layers of the spinal cord and its contents.
- List subdivisions of mantle & marginal zones.
- List meningeal layers and describe positional changes of spinal cord.
- Describe development of vertebral column from sclerotomic portion of paraxial mesoderm.
- Describe chondrification & ossification stages in vertebral development.
- Describe spina bifida and its types.




DEVELOPMENT OF SPINAL CORD

DEVELOPMENT OF NEURAL TUBE


□ Ectodermal cells dorsal to notochord thicken to form the neural plate.

□ A longitudinal groove develops in the neural plate (neural groove).


□ The margins of the neural plate (neural folds) approach to each other and fuse to form the neural tube.




A




B




C



D




E



F

DEVELOPMENT OF SPINAL CORD




□ The spinal cord develops from the caudal 2/3 of the neural tube



DEVELOPMENT OF SPINAL CORD

The cells of neural tube form:

- ❑ An inner **ventricular zone** of undifferentiated cells
- ❑ A middle **mantle zone** of cell bodies of neurons (**future grey matter**)
- ❑ An outer **marginal zone** of nerve fibers or axons of neurons (**future white matter**)




MANTLE LAYER OF SPINAL CORD



- ❑ Neurons of mantle layer (future grey matter) differentiate into:
 1. A dorsal alar plate (future dorsal horn): containing **sensory** neurons
 2. A ventral basal plate (future ventral horn): containing **motor** neurons
- ❑ The 2 areas are separated by a longitudinal groove (sulcus limitans).


MANTLE LAYER OF SPINAL CORD



Proliferation and bulging of both alar & basal plates cause:

- ❑ Formation of longitudinal dorsal & ventral median septa
- ❑ Narrowing of the lumen to form a small central canal

MARGINAL LAYER OF SPINAL CORD




- ❑ Marginal layer increases in size due to addition of ascending, descending & intersegmental nerve fibers.
- ❑ Myelination of nerve fibers starts at **4th month** & continues **during the 1st postnatal period**. Motor fibers myelinate before sensory fibers.
- ❑ Marginal layer (future white matter) is divided into: dorsal, lateral and ventral funiculus (white column)


MENINGES

- ❑ They are 3 membranes covering the neural tube:
 1. Outer thick **dura** matter: mesodermal in origin
 2. Middle **arachnoid** matter: ectodermal in origin
 3. Inner thin **pia** matter: ectodermal in origin
- ❑ A cavity appears between arachnoid & pia (subarachnoid space) & becomes filled with cerebrospinal fluid.


POSITIONAL CHANGES OF SPINAL CORD




- ❑ Initially, the spinal cord occupies the whole length of the vertebral canal.
- ❑ As a result a faster growth of vertebral column, the caudal end of spinal cord (**conus medullaris**) shift gradually to a higher level.




A




B




C



D



E



F


Notochord stimulates :

- Neural tube formation
- Vertebral column formation

Neural tube

INTRAEMBRYONIC MESODERM


- ❑ Proliferates between Ectoderm & Endoderm
EXCEPT in the central axis of embryo where **NOTOCHORD** is found.
- ❑ Differentiates into 3 parts:
 1. **Paraxial mesoderm**
 2. Intermediate mesoderm
 3. Lateral mesoderm
- ❑ Paraxial mesoderm divides into units (somites).
- ❑ Each somite divides into 3 parts:
 1. **Sclerotome**
 2. Myotome
 3. Dermatome




A

1- Loosely arranged cells


2- densely packed cells



B



C



D




FORMATION OF BODY OF VERTEBRA

- ❑ **At 4th week**, each sclerotome is formed of:
 1. A cranial part of loosely arranged cells
 2. A caudal part of densely packed cells
- ❑ The caudal part of each sclerotome fuses with the cranial part of succeeding sclerotome to form the centrum (body primordium)
- ❑ Each centrum develops from 2 adjacent sclerotomes.




FATE OF NOTOCHORD

- ❑ **In the region of the bodies of vertebrae: It degenerates .**
 - ❑ **Between bodies of vertebrae: It forms the intervertebral discs (nucleus pulposus).**
- N.B.:** Annulus fibrosus part of the intervertebral discs are formed by the mesoderm surrounding the notochord.

VERTEBRAL DEVELOPMENT




Fusion of bony halves of vertebral arch occurs at 3-5 years



All centers unite around 25 years

CURVATURES OF VERTEBRAL COLUMN



□ Primary curves
(thoracic & pelvic or sacral): develop prenatally


□ Secondary curves:
develop postnatally

1. **Cervical:** as a result of lifting the head
2. **Lumbar:** as a result of walking


SPINA BIFIDA

- ❑ **Cause:** Failure of fusion of the halves of vertebral arches
- ❑ **Incidence:** 0.04-0.15%
- ❑ **Sex:** more frequent in females
- ❑ **Types:**
 1. Spina bifida occulta (20%)
 2. Spina bifida cystica (80%)


Spina bifida occulta




Spina bifida with meningocele



Spina bifida with meningoencephalocele



Spina bifida with myeloschisis



SPINA BIFIDA OCCULTA




- The closed type
- Only one vertebra is affected
- No clinical symptoms
- Skin overlying it is intact
- Sometimes covered by a tuft of hair

SPINA BIFIDA CYSTICA

With meningocele



With myeloschisis



SPINA BIFIDA CYSTICA

- ❑ The open type
- ❑ Neurological symptoms are present
- ❑ Subdivided into:
 1. **Spina bifida with meningocele:** protrusion of sac containing meninges & cerebrospinal fluid
 2. **Spina bifida with meningomyelocele:** protrusion of sac containing meninges with spinal cord and/or nerve roots
 3. **Spina bifida with myeloschisis:** spinal cord is open due to failure of fusion of neural folds

SUMMARY OF DEVELOPMENT OF SPINAL CORD

- ❑ The spinal cord develops from the caudal 2/3 of the ectodermal neural tube.**
- ❑ Layers of spinal cord are (from inside outward): ventricular, mantle (future grey matter) and marginal (future white matter).**
- ❑ Mantle layer differentiates into dorsal alar plate (with sensory neurons) & ventral basal plate (with motor neurons) separated by sulcus limitans.**
- ❑ Marginal layer is divided into dorsal, lateral & ventral funiculus.**

SUMMARY OF DEVELOPMENT OF SPINAL CORD

- ❑ Myelination of nerve fibers starts at 4th month & continues during the 1st postnatal period. Motor fibers myelinate before sensory fibers.**
- ❑ Meninges are 3 membranous sac covering the neural tube (from outside inward): dura (mesodermal in origin), arachnoid and pia (both are ectodermal in origin).**
- ❑ A cavity between arachnoid & pia matters (subarachnoid space) contains cerebrospinal fluid.**
- ❑ During development the end of spinal cord shifts its position: at 24 weeks (level of S1), at birth (level of L3), adult position (level of L1-L2).**

SUMMARY OF DEVELOPMENT OF VERTEBRAL COLUMN


- ❑ Vertebral column develops from sclerotomic portion of paraxial mesoderm.
- ❑ Sclerotome around neural tube forms vertebral (neural) arch.
- ❑ Sclerotome around notochord forms body of vertebra. Each body develops from 2 adjacent sclerotomes.
- ❑ Notochord forms nucleus pulposus portion of the intervertebral discs.
- ❑ Chondrification centers appear at 6th week.
- ❑ Three primary ossification centers appear at 8th week.

SUMMARY OF DEVELOPMENT OF VERTEBRAL COLUMN

- ❑ Fusion between halves of neural arch occurs at 3-5 years, between neural arch & body at 3-6 years.
- ❑ Five secondary ossification centers appear at puberty and fuse around 25 years.
- ❑ Spina bifida is due to failure of fusion of the halves of the neural (vertebral) arch. It may be **occulta** (20%, closed type, no symptoms) or **cystica** (80%, open type, with symptoms).

QUESTION 1

Which one of the following regions of spinal cord contains cell bodies of sensory neurons?

1. Alar plate 
2. Ventricular zone
3. Basal plate
4. Dorsal funiculus


QUESTION 2

At which one of the following periods of life fusion between vertebral arch & body of vertebra occurs?

1. 8th week
2. Puberty
3. 3-6 years ←
4. Around 25 years

QUESTION 3

Regarding spina bifida which one of the following statements is correct?

1. The closed type is more frequent than the open type.
2. The closed type presents with clinical symptoms.
3. Spina bifida is due to failure of fusion between the halves of vertebral arch. 
4. In cases of spina bifida with meningocele, the spinal cord is open.

A stylized illustration of two yellow roses with green leaves on a dark teal background. The roses are rendered in a low-poly, geometric style. The text 'THANK YOU' is overlaid in the center in a bold, blue, sans-serif font with a slight drop shadow.

THANK YOU