

Vision - 4

# Color Vision

*Dr. Salah Elmalik*

# Objectives

- Define color vision Identify and describe the mechanism of color vision and the three types of cones, including the range of spectral sensitivity and color blindness
- Identify color vision theory
- Describe the items needed for any color perception Compare different types of color blindness

Red Rose



WEBSHOIS

# Color Vision

- It is the ability to discriminate between different colors.
- 1- there are 3 primary colors( blue- red- green) sensed by cones in fovea & appreciated within photopic vision.
- 2- sensation of extraspectral colors as white, yellow, orange, purple, can be produced by mixing properties of the blue & red & green in different combinations.
- 3- black means absence of light ( not darkness because in dark we do not see black only)

# Color (Photopic) Vision

'Young - Helmholtz theory'

'The Trichromatic theory'

# Color vision theory :\_( Young-Helmholtz theory )

- 1- we have 3 kinds of cones each has a specific photopigment (iodopsin)& is sensitive to one of the 3 primary colors
- a- Blue cone system:- has S pigment ( blue sensation pigment) which respond to short wave length ( 440 nm senses the blue color)
- b- Green cone system:- has M pigment ( green sensation pigment) which respond to middle wave length ( 535 nm senses the green color & less to yellow) & absorb light at the green portion.

- Red cone system:- has L pigment ( red sensation pigment) which respond to large wave length at or  $> 535$  nm so senses the red & yellow color & absorb light at the red portion.








# History of color vision

Newton (1704) used a prism to show that sunlight was composed of light with all colors in the rainbow. He defined it as the **spectrum**.



# Mixing colors




# Photopic vision (CONES)

**Helmholtz ..1860:**

The three primary colors are perceived by three photoreceptor pigments (with broad absorption curves)

White light is produced by mixing three colours

# Cone wavelength ranges




# Photopic vision (CONES)

Cone pigments: three kinds



# Cone wavelength ranges



# Photopic vision

Sensation of any color determined by:

a-wavelength of light

b-amount of light absorbed by each type of  
cones

c-frequency of impulses from each cone system  
to ganglion cells which is determined by  
wave length of light.



# Photopic vision





1994 The Exploratorium

perception of white is due to:  
equal stimulation of

blue & red & green cones.

(white is a combination of all wave  
lengths)






# Color vision is coded by :-

- # different responses in ganglion cells that depends upon the wave length of stimulus which determine frequency of impulses in ganglion cells
- # the color perception in the brain depends on the amount of activity in each of the 3 cone systems as mentioned above.

# *Color Perception*


- Perception of orange is due to stimulation of 99% of red cones & 42% of green cones & 0% of blue cones( so ratio is 99:42: 0)
- For perception of yellow the ratio is 83:83: 0.
- Perception of blue is due to stimulation of 0% of red cones & 0% of green cones & 97% of blue cones( so ratio is 0:0: 97 )

# Test for Color Blindness



The above has been reproduced from Ishihara's Tests for Colour Blindness published by KANEHARA & CO., LTD., Tokyo, Japan, but tests for colour blindness cannot be conducted with this material. For accurate testing, the original plates should be used.

# Ishihara Charts



# Color Blindness

Weakness or total blindness in detecting a primary color:

## Definitions:

1. **Trichromats**: see the 3 1ry colors
2. **Dichromats**: blind to one 1ry color
3. **Monochromats**: have only one color pigment

# Color Blindness -cont.

- ▣ Prot ..... Red
  - ▣ Deuter .... Green
  - ▣ Trit ..... Blue
  - ▣ Anamoly ...weakness
- 
- ▣ Protanamoly
  - ▣ Deuteranamoly
  - ▣ Tritanamoly
- Trichromats



# Color Blindness -cont.



- ▣ Anamoly ...weakness
  - ▣ Anopia .... Total loss
  
  - ▣ Protanopia
  - ▣ Deuteranopia
  - ▣ Tritanopia
- Dichromats
- 
- ```
graph LR; A[Protanopia] --- B[Dichromats]; C[Deuteranopia] --- B; D[Tritanopia] --- B;
```

# COLOR BLINDNESS

## Red - Green Blindness:-



- Green & red cones see different colors between wave length 525-675 nm & distinguish them.
- If either of these cones are absent, the person can not distinguish 4 colors ( red - green- yellow- orange)& he can not distinguish red from green (primary colors) so called ( red - green blindness).

## Trichromatic Vision



## Dichromatic Vision

Protanope – severe red/green color deficiency



# Color Blindness -cont.

- It is x- linked disease transmitted from females to their male sons, never occur in females as they have 2 x chromosomes -
- Males have one x & one y chromosome so if this one x chromosome miss the gene for color vision , he will get red-green color blindness(their gene is on x chromosome). -
- Females show the disease only if both x chromosomes lack the gene -
- Females from color blind fathers are carriers transmit the disease to  $\frac{1}{2}$  of their sons.

