

Metabolic Syndrome

Endocrine Block

Dr. Sumbul Fatma

Objectives

- Define metabolic syndrome, insulin resistance and dyslipidemia
- Discuss the risk factors for metabolic syndrome and other medical conditions associated with it
- Define the diagnostic criteria for Metabolic syndrome
- Discuss the management of metabolic syndrome and current treatment options

Overview

- Introduction
- Features of metabolic syndrome
- Insulin resistance
- Dyslipidemia
- Risk factors
- Markers
- Diagnosis
- Management and treatment

Metabolic Syndrome

- A combination of metabolic abnormalities which increase the risk of heart disease, diabetes and other diseases
- Obesity is a component of met. synd.
- Signals from adipocytes in obesity cause metabolic abnormalities such as:
 - Dyslipidemia
 - Glucose intolerance
 - Insulin resistance
 - Hypertension

Features of metabolic syndrome


- Obesity
- High serum triglycerides (TGs)
- Low HDL cholesterol
- Hypertension
- Hyperglycemia
- Insulin resistance (hyperinsulinemia)

Insulin Resistance

- Cells become less responsive to insulin → high plasma insulin → hyperglycemia
- Hydrolysis of stored fats → high plasma FFAs
- Reduction of glucose uptake/use by cells
- Reduction of glycogenesis → hyperglycemia
- Compensatory hyperinsulinemia causes down regulation of insulin receptor
- Defects in insulin receptor

Dyslipidemia

- Insulin resistance in adipocytes → increased activity of hormone-sensitive lipase → high plasma FFAs
- FFAs → TGs/cholesterol in the liver
- Excess TGs/cholesterol are released as VLDL in the blood
- HDL levels are decreased



Copyright © 2008 Wolters Kluwer Health | Lippincott Williams & Wilkins

Dyslipidemia and met. synd. are strongly related

- Dyslipidemia is an early indicator of insulin resistance
- Liver fat plays a major role in dyslipidemia due to insulin resistance

Risk factors for metabolic syndrome

- Obesity
- Alcoholism
- Sedentary Lifestyle
- Smoking
- Hypercortisolism (Steroid use/Cushing disease)
- Drugs (rifampicin, isoniazid, etc.)
- Mutations in insulin receptor

Metabolic syndrome is linked to:

- Heart disease
 - 1.5-3 fold increase in atherosclerosis
- Type-2 diabetes mellitus
 - 5-fold increase
- Kidney disease
- Reproductive abnormalities in women
 - Polycystic ovarian syndrome
 - Impaired ovulation and fertility
 - Irregular menstruation

Metabolic syndrome is linked to:

- Nonalcoholic steatohepatitis (fatty liver disease)
 - Related to impaired lipid metabolism
- Cancer
 - Obesity is a major risk factor for cancer of esophagus, colon and rectum, liver, gall bladder
 - Being overweight and obese accounts for 14% of all cancer deaths in men and 20% of those in women

Diagnosis – WHO criteria (1999)

- Impaired glucose tolerance
- Diabetes mellitus
- Insulin resistance

PLUS any of these two:

Component	Criterion
Hypertension	BP >140/90 mmHg
Dyslipidemia	High plasma TGs (>1.7mmol/L) Low HDL cholesterol (men <0.9, women <1.0 mmol/L)
Central or General obesity	Waist to hip ratio >0.9 in men, >0.85 in women And/or BMI >30
Microalbuminuria	Urinary albumin excretion rate \geq 20ug/min or albumin:creatinine ratio \geq 30mg/g

NCEP* ATP** III Guideline (2002)

Diagnosis: If any 3 or more of these risk factors are present

- Waist circumference:
 - Men >102 cm (>40 in)
 - Women >88 cm (>35 in)
- Triglycerides >150 mg/dL
- HDL cholesterol:
 - Men <40 mg/dL
 - Women <50 mg/dL
- Blood pressure 130/ 85 mm Hg
- Fasting glucose >100 mg/dL

*National Cholesterol Education Program

**Adult Treatment Panel

Markers of metabolic syndrome

- Lipoproteins (LDL, HDL)
- Adipokines (Leptin, adiponectin)
- Inflammatory markers
 - c-reactive protein, TNF- α , IL-6, IL-8
- Hemostatic marker
 - Plasminogen activator inhibitor-1

Managing Metabolic Syndrome

- Primary intervention: Lifestyle changes
 - Weight reduction
 - Target BMI < 25
 - Reduced intake of calories and fats
 - More physical activity
 - Smoking cessation

Managing metabolic syndrome

- Secondary intervention: Medication to treat existing risk factors
 - Management of
 - Blood pressure (anti-hypertensive drugs)
 - Lipids (statins, fibrates)
 - Blood glucose (metformin, TZDs)
 - Aspirin for CVD prevention

Lowering blood pressure

Modification	Recommendation	Average drop in SBP
Weight loss	Maintain normal body weight	5-10 for every 22lbs loss
Healthy eating plan	Meals rich in fruits, vegetables; low fat dairy; low saturated fats and cholesterol	8-14
Sodium restriction	< 2400 mg/day	2-8
Regular physical activity	30 min. most of the week	4-9

Hypertension and clotting disorders

- Treat hypertension to goal (< 130/80 mmHg)
- Low dose diuretics
- ACE inhibitor
- Aspirin:
 - To treat clotting disorders
 - Daily low dose aspirin (81-325mg) for:
 - Men > 45
 - Postmenopausal women

Current Treatment

- Statins
- Metformin
- Fibrates
- Thiazolidinediones (TZDs)
- Aspirin

Metformin

- Reduces blood glucose levels by inhibiting hepatic gluconeogenesis
 - Hepatic gluconeogenesis is active in patients due to liver's resistance to the effects of insulin
- Reduces lipid synthesis in the liver
- Helps reducing blood lipids

Fibrates

- Reduce blood lipid levels
- Activate transcription factor:
 - Peroxisome proliferator activated receptor- α (PPAR- α)
- Activated PPAR- α \rightarrow transcription of genes of lipid degradation / uptake by the cells:
 - Carnitine:palmitoyl transferase I (enhances FA uptake into mitochondria)
 - Lipoprotein Lipase
 - Stimulates apoA1 and apoAII protein synthesis (major proteins in HDL)

Thiazolidinediones (TZDs)

- Used for the treatment of insulin resistance and type-2 diabetes mellitus
- TZDs activate PPAR- γ class of transcription factors expressed primarily in the adipose tissue
- Activates the transcription of adiponectin
- Adiponectin reduces the fat content of the liver and enhances insulin sensitivity

Take home message

- Metabolic syndrome is a combination of metabolic abnormalities that increase the risk of heart disease, diabetes and other diseases
- The features of metabolic syndrome include obesity, high serum triglycerides (TGs), low HDL cholesterol, hypertension, hyperglycemia and insulin resistance
- Obesity, alcoholism, sedentary lifestyle and smoking are some of the risk factors for metabolic syndrome
- Management of the syndrome includes lifestyle modifications to reduce weight and medications

References

- Textbook of Biochemistry with Clinical Correlations by Thomas M. Devlin, 6th Edition, pp 862-863.
- Lippincott's Biochemistry. 5th Edition, pp 353-355, Lippincott Williams & Wilkins, New York, USA.