

*Use of Insulin in the treatment
of diabetes mellitus*

Prof. Hanan Hagar



Objectives:

by the end of this lecture, students should be able to:

- *Define diabetes and mention different types of diabetes*
- *Differentiate between difference in treating type I and type II diabetes.*
- *Understand mechanism of action, secretion, and actions of insulin.*
- *Describe different types of insulin analogues*
- *Be able to recognize the difference in pharmacokinetic profile between different types of insulin analogues.*
- *Know uses of different insulin analogues*

Diabetes mellitus

- Is a **chronic metabolic disorder** characterized by high blood glucose level caused by caused by deficiency of insulin or by increased insulin resistance.

Diabetes mellitus

Fasting plasma glucose (no food for 8 hrs)

Normal <100 mg/dl (5.6 mmol/l).

Pre-diabetes 100-125 mg/dl (5.6-6.9 mmol/L).

Diabetes if

Fasting >126 mg/dl (7 mmol/L)

or 2h after a meal > 200 mg/dl (11.1 mmol/L).

Complications of diabetes

- **Cardiovascular problems**
 - **Micro- and macro-vascular disease**
- **Renal failure (nephropathy).**
- **Blindness (retinopathy).**
- **Neuropathy.**
- **Risk of foot amputation**

Types of diabetes

- **Type I diabetes (IDDM)**
due to autoimmune or viral diseases
- **Type II diabetes (NIDDM)**
due to genetic susceptibility and other factors (age, obesity).

Type I Diabetes

IDDM

- 10-20% occurrence.
- During childhood or puberty
- β -cells are **completely destroyed.**
- Absolute deficiency of insulin secretion
- Treated **by insulin.**

Type II Diabetes

NIDDM



- **80-90% occurrence**
- **Over age 35**
- **Pancreatic β -cells are not producing enough insulin**
- **Obesity is an important factor.**
- **Insulin resistance in peripheral tissues.**
- **Treated by oral hypoglycemic drugs.**

Characteristic	Type 1	Type 2
Onset (Age)	Usually during childhood or puberty	Usually over age 40
Type of onset	Abrupt	Gradual
Prevalence	10-20%	80-90 %
Genetic predisposition	Moderate	Very strong
Defects	β -cells are destroyed	β -cells produce inadequate quantity of insulin
Endogenous insulin	Absent	Present (not enough)
Insulin resistance	absent	present
Nutritional status	Usually thin	Usually obese
Ketosis	Frequent	Usually absent
Clinical symptoms	Polydipsia, polyphagia, polyuria, weight loss	Often asymptomatic
Related lipid abnormalities	Hypercholesterolemia frequent	Cholesterol & triglycerides often elevated
Treatment	Insulin injection	Oral hypoglycemic drugs

A decorative graphic in the top-left corner consisting of a light green square and a white rounded rectangle. A thick, dark blue horizontal bar spans across the top of the page.

INSULIN

Human Insulin



Insulin receptors

- **Present on cell membranes of most tissues.**
- **Liver, muscle and adipose tissue**

A decorative graphic in the top-left corner consisting of a light green square partially overlapping a white rounded rectangle. A thick, dark blue horizontal bar spans across the white area below the rounded rectangle.

Effects of insulin

I. Carbohydrate Metabolism:

- ↑ glucose uptake & utilization by peripheral tissues.
- ↑ Glycogen synthesis (**glycogen synthase**)
- ↑ Conversion of carbohydrate to fats.
- ↓ Gluconeogenesis.
- ↓ Glycogenolysis (liver).
- ↑ Glycolysis (muscle).

II. Fat Metabolism:

- **Liver:**

- ↑ **Lipogenesis.**
- ↓ **Lipolysis.**
- **Inhibits conversion of fatty acids to keto acids.**

- **Adipose Tissue:**

- ↑ **Triglycerides storage.**
- ↑ **Fatty acids synthesis.**
- ↓ **Lipolysis**

III. Protein Metabolism:

Liver:

- ↓ protein catabolism.

Muscle:

- ↑ amino acids uptake.
- ↑ protein synthesis.
- ↑ glycogen synthesis (**glycogenesis**).

IV. potassium

- **↑ potassium uptake into cells.**

Routes of administrations of exogenous insulin

- **Can not be given orally (why ?)**
- **Insulin syringes (s.c., arms, abdomen, thighs).**
- **Portable pin injector (pre-filled).**
- **Continuous S.C. infusion (insulin pump).**
 - **More convenient**
 - **Eliminate multiple daily injection**
 - **Programmed to deliver basal rate of insulin.**

Routes of administrations of exogenous insulin

- **Intravenously (in a hyperglycemic emergency)**

Under Clinical Trials

- **Inhaled aerosols, transdermal, intranasal.**

Pin injector



Insulin pump



Insulin degradation

1. Basal level of endogenous insulin is 5-15 $\mu\text{U/ml}$.
2. Half life of circulating insulin is 3-5 min.
3. 60% liver & 40% kidney (**endogenous insulin**)
4. 60% kidney & 40% liver (**exogenous insulin**)

Sources of Exogenous Insulin

- **Beef Insulin**
 - Differs from human insulin by 3 amino acids (antigenic).
- **Porcine Insulin**
 - Differs by one amino acid (antigenic).

Human Insulin analogues

- **Prepared by recombinant DNA techniques.**
- **Less immunogenic.**
- **Modifications of amino acid sequence of human insulin can change pharmacokinetics.**

Types of insulin preparations

Differ in pharmacokinetic properties mainly

- Rate of absorption (Onset of action).
- Duration of action.

Variation is due to:

- Change of amino acid sequence.
- Size and composition of insulin crystals in preparations (monomers, dimers, hexamers).

Types of insulin preparations

Insulin Analogues

- **Ultra-short acting insulins**
e.g. Lispro, aspart
very fast onset of action and short duration
- **Short acting insulins**
e.g. regular insulin
fast onset of action and short duration.

Types of insulin preparations

- **Intermediate acting insulins**
 - e.g. NPH, lente
 - **Slow onset, intermediate duration of action.**
- **Long acting insulins**
 - e.g. glargine, detemir
 - **Slow onset and long duration of action.**

Ultra-short acting insulins

Insulin lispro, insulin aspart

- **Clear** solutions at neutral pH.
- Do not aggregate or form dimers or hexamers (**monomeric analogue**).
- **Fast onset** of action (5-15 min)
- S.C. (5 -15 min before meal).
- **Short duration** of action (3-5 h)


Ultra-short acting insulins

Insulin Lispro, insulin aspart


- **Reach peak level 30-90 min after injection.**
- **3 times/day.**
- **Mimic the prandial mealtime insulin release.**
- **I.V. in emergency.**

Insulin aspart

Structure



Ultra-short acting insulins



Ultra-short acting insulins

insulin lispro, insulin aspart

- Preferred for external insulin pump (**Lispro does not form hexamers**)
- used to control post-prandial hyperglycemia (s.c.) and emergency diabetic ketoacidosis (i.v).

Short acting insulins (Regular insulin)

- **Soluble** crystalline zinc insulin
- **Clear** solutions at neutral pH.
- **Forms hexamers.**
- **Onset of action 30-45 min (s.c.).**
- **I.V.** in emergency situations.
- **Peak 2-4 h.**
- **Duration 6-8 h.**

Short acting insulins (regular insulin)

- **2-3 times/day.**
- **Control postprandial hyperglycemia (s.c.) & emergency diabetic ketoacidosis (i.v.).**
- **Can be used in pregnancy**

	Ultra-Short acting insulins e.g. Lispro, aspart, glulisine
Physical characteristics	Clear solution at neutral pH
chemistry	Monomeric analogue
Route & time of administration	S.C. 5 min (no more than 15 min) before meal I.V. in emergency (e.g. diabetic ketoacidosis)
Onset of action	Fast 5 – 15 min (S.C)
Peak level	30 – 90 min
Duration	3 – 5 hr Shorter
Usual administration	2 – 3 times/day
	postprandial hyperglycemia & emergency diabetic ketoacidosis

Short-acting (regular) insulins e.g. Humulin R, Novolin R
Clear solution at neutral pH
Hexameric analogue
S.C. 30 – 45 min before meal I.V. in emergency (e.g. diabetic ketoacidosis)
rapid 30 – 45 min (S.C)
2 – 4 hr
6 – 8 hr longer
2 – 3 times / day
postprandial hyperglycemia & emergency diabetic ketoacidosis

Advantages of Insulin Lispro vs Regular Insulin

- **Rapid onset of action (due to rapid absorption)**
- **Reduced risk of postprandial hypoglycemia and hyperinsulinemia (due to shorter duration of action, no more than 3-4 hrs regardless of dose).**

Intermediate acting insulins

Isophane (NPH) insulin

Lente insulin

Isophane (NPH) Insulin

- **NPH, is a Neutral Protamine Hagedorn insulin in phosphate buffer.**
- **NPH insulin is combination of protamine & crystalline zinc insulin (1: 6 molecules). proteolysis release insulin.**


Isophane (NPH) Insulin

- **Turbid suspension** at neutral pH.
- **Given S.C. only not i.v.**
- **Can not be used in ketoacidosis or emergency**
- **Onset of action 1-2 h.**
- **Peak serum level 5-7 h.**
- **Duration of action 13-18 h.**


Isophane (NPH) Insulin

Insulin mixtures

- **NPH/regular insulin**
 - **75/25 , 70/30 , 50/50**
- **(NPL= NPH / lispro) (NPA= NPH / aspart)**
- **NPL & NPA have the same duration as NPH**
- **Have two peaks.**

A

Prandial and basal insulin replacement

B

Lente insulin

- **Mixture of:**
 - **30% semilente insulin (amorphous precipitate of zinc insulin in acetate buffer)**
 - **70% ultralente insulin (poorly soluble crystal of zinc insulin)**
- **Turbid suspension at neutral pH**
- **Given S.C., not intravenously**

Lente insulin (Humulin L, Novolin L)

- **Delayed onset of action (1-3 h)**
- **Peak serum level 4-8 h.**
- **Duration of action 13-20 h.**
- **Lente and NPH insulins are equivalent in activity.**
- **Lente is not used in diabetic ketoacidosis or emergency.**

Long acting insulins

Insulin glargine (lantus), Insulin detemir (Levemir)

Insulin glargine (Lantus)

- Clear solution **BUT** forms precipitate (hexamer) at injection site.
- Slow onset of action 2 h.
- absorbed less rapidly than NPH & Lente insulin.
- Given s.c., **not intravenously**
- Should not be mixed with other insulins in the same syringe.


Insulin glargine (Lantus)

- **Maximum effect after 4-5 h**
- **Prolonged duration of action (24 h).**
- **Once daily**
- **produce broad plasma concentration plateau (low continuous insulin level).**
- **Glargine must be used in regimens with rapid or short acting insulins.**


Advantages over intermediate-acting insulins:

- **Constant circulating insulin over 24 hr with no peak (**peakless profile**).**
- **Produce flat prolonged hypoglycemic effect.**
- **Safer than NPH & Lente insulins
(reduced risk of nocturnal hypoglycemia).**

NPH vs Glargine



Insulin preparations



Insulin Dosing considerations

- **Blood glucose monitoring is required in all patients receiving insulin**
- **Rotate injection sites within the same region.**
- **Insulin should be stored in refrigerator and warm up to room temp before use.**

Complications of Insulin Therapy:

- Hypoglycemia
- Hypersensitivity reactions.
- Lipodystrophy at injection site
- Weight gain (due to anabolic effects of insulin)
- Insulin resistance
- Hypokalemia

Summary

- Insulin analogues are used to treat type I diabetes.
- *Fast acting insulins (lispro, aspart)*, given s.c. or i.v., produce fast action, used to mimic postprandial insulin.
- *Short acting insulin (Regular insulin)*, given s.c. or i.v. produce rapid action, used to mimic postprandial insulin.
- *Intermediate acting insulin (lente, Isophane)* produce slower action, than regular insulin, given s.c. not i.v.
- *Long acting insulins (glargine, detemir)* produce constant circulating insulin over 24 hr with no peak (peakless profile), s.c. not i.v.