

# Physiology of Consciousness



- Is the brain state in which a person is being aware of the self and surroundings
- It is a product of electrical activity of the brain



# 4 levels of consciousness

- 1- Normal consciousness
- 2- Clouded consciousness
- 3- Sleep
- 4- Coma

# Level of consciousness

- (1) Normal Consciousness

State of normal arousal

Being fully awake

And

Aware of the self and surroundings

- (2) Clouded consciousness : person conscious but mentally confused
- e.g., in cases of drug or alcohol intoxication
- High fever associated (malaria or septicemia , dementia , etc)



- (3) Sleep : person unconscious ( in relation to the external world & surroundings)
- but is arousable ( can be aroused ) .



- (4) Coma : person unconscious and not arousable



# brain Structures involved in the conscious state:

- 1- Brain stem Reticular formation
- 2- Thalamus
- 3- Hypothalamus
- 4- Ascending projection pathways
- 5- Wide spread area in the cerebral cortex




# 1 - Reticular formation

Set of interconnected nuclei that are located throughout the brainstem (Pons, Midbrain, Upper medulla), and the thalamus

Role in behavioral arousal


Role in consciousness (sleep/awake cycle)

Connect the brain stem to the CC



# Reticular Activating System

Buzzle.com





# consists of 3 parts:

Lateral Reticular Formation

Paramedian Reticular Formation

Raphe nuclei (Median RF)





# ○ Lateral Reticular Formation

Has small neurones


Receives information from ascending tracts for touch and pain.


Receives vestibular information from median vestibular nerve.

Receives auditory information from superior olivary nucleus.

Visual information from superior colliculus.

Olfactory information via medial forebrain bundle






# Paramedian Reticular Formation


Has large cells.

Receives signals from lateral reticular formation

Contains noradrenergic (NA) & Dopaminergic (DA) neurones, projects onto cerebral hemispheres.

Cholinergic (Ch1) neurones project onto the thalamus









# Raphe nuclei (Median RF)

In the midline of the reticular formation

Contain serotonergic projections to the brain and spinal cord.





# Functions of reticular formation:

- **1. Somatic motor control** (Reticulospinal tracts)
- **2. Cardiovascular control**
  - Through cardiac and vasomotor centers of the medulla oblongata
- **3. Pain modulation**
- Pain signals from the lower body >> >> RF >> >> cerebral cortex
- RF is origin of the descending analgesic pathways
- (act on the spinal cord to block the transmission of some pain signals to the brain)



# 2- Thalamus:


Located in the mid-part of the diencephalon

Cholinergic projections from the thalamus are responsible for:


- ⊙ Activation of the cerebral cortex.
- ⊙ Regulation of flow of information through other thalamic nuclei to the cortex via projections into reticular nuclei.

# 3 - Hypothalamus

- Tuberomammillary nucleus in the hypothalamus projects to the cortex and is involved in maintaining the awake state








- **Bulboreticular Facilitory ( Excitatory ) = Reticular Excitatory Area of the Brain Stem**
- **Sends excitatory signals into Thalamus**
- **>>>> thalamus excites almost all areas of the cortex .**
- **The Bulboreticular Facilitory ( Excitatory ) Area + Thalamus =Reticular Activating System ( RAS)**
- **The RAS is the system which keeps our cortex awake and conscious**






# Anatomical components of RAS

- The RAS is composed of several neuronal circuits connecting the brainstem to the cortex
- Originate in the upper brainstem reticular core and project through synaptic relays in the thalamic nuclei to the cerebral cortex
- As a result, individuals with bilateral lesions of thalamic intralaminar nuclei are lethargic or drowsy



# RAS

- Lesion in the mid-pons  
>>>>>>unconsciousness
- Pons (uppers & middle) and midbrain are essential for wakefulness .





# Sensory inputs to RAS

## Control Loop

Feed-Back Differential

Ascending Neural Radiations to Cortex

Descending Neural Radiations to the Hippocampus/Thalamus/hypothalamus

Cerebral Cortex

Anterior Thalamic Nucleus

Cerebral Hemisphere

Olfactory Bulb

Visual Impulses

Hypothalamus

Pituitary Gland

Mamillary Body of Hypothalamus

Amygdaloid Nucleus

Corpus Callosum

Thalamus

Pineal Gland


Hippocampus

Cerebellum

Auditory Impulses

Projection to Spinal Cord

Ascending Sensory Tracts



# Functions of RAS:

- 1- Regulating sleep-wake transitions
- If inhibitory area activity increase >>>>  
reduce the activity of RAS >>>>> less  
afferent signal to the CC >>>>> sleep

# 2-Attention

- RAS mediate transitions from relaxed wakefulness to of high attention.

# 3-RAS and learning

- The RAS is the center of balance for the other systems involved in learning, self-control or inhibition, and motivation.
- Provides the neural connections for processing and learning of information,
- Selective attention (to the correct task)

# RAS dysfunction

If RAS is depressed:

An under-aroused cortex

Difficulty in learning

Poor memory

Little self-control

lack of consciousness or even coma.

If the RAS is too excited,

Over aroused cortex

**Hyper-vigilance (sensory sensitivity )**

Touching everything

Talking too much

Restless


Hyperactive




# Indices of Level of Consciousness

- Appearance & Behavior :
  - Posture ( sitting , standing ? )
  - Open eyes ?
  - Facial expression ?
  - Responds to stimuli ( including the examiner's questions about name , orientation in time & place ? & other general Qs like who is the president ? )
- Vital signs :
  - Pulse , BP , respiration , pupils , reflexes , particularly brainstem reflexes , etc )
  - EEG → Each of these states ( wakefulness , sleep , coma and death ) has specific EEG patterns
  - Evoked potentials ( in cases of Brain Death ).


# Electroencephalogram



- Alpha waves:
  - Recorded from the parietal & occipital regions
  - Awake and relaxed + eyes closed
  - 10 to 12 cycles/second.
  
- Beta waves:
  - Frontal lobes
  - Produced by visual stimuli and mental activity
  - 13 to 25 cycles per second .




- **Theta:**
  - Temporal and occipital
  - 5 to 8 cycles/second
  - Normal in newborn
  - Theta waves in adults indicates severe emotional stress
- **Delta:**
  - From the cerebral cortex
  - 1 to 5 cycles/second
  - Sleep (adults) and in an awake infant
  - In an awake adult indicates brain damage.







Normal EEG ( at  
normal  
magnification )



Brain Death ( Flat EEG ,at very high  
magnification )

