

PHYSIOLOGY OF SYMPATHETIC AND PARASYMPATHETIC NERVOUS SYSTEM




Prof. Sultan Ayoub Meo


MBBS, Ph.D, M Med Ed (Scotland), FRCP (London), FRCP
(Dublin), FRCP (Glasgow), FRCP (Edinburgh)

Professor, Department of Physiology, College of Medicine
King Saud University, Riyadh, KSA


THE NERVOUS SYSTEM



THE NERVOUS SYSTEM



THE AUTONOMIC NERVOUS SYSTEM



THE AUTONOMIC NERVOUS SYSTEM



SYMPATHETIC (GAS PEDAL)

- Fight or flight response
- Protection and survival
- Stress response
- Adrenal (stress) glands activated

PARASYMPATHETIC (BRAKE PEDAL)

- Rest
- Digest
- Relax
- Growth & development




The Human Nervous System



"You can't be in growth and protection at the same time."

— Dr. Bruce Lipton

DISTRIBUTION OF THE SYMPATHETIC AND PARASYMPATHETIC NERVOUS SYSTEM





Parasympathetic: Craniosacral: Originate from cranial nerves (3rd, 7th, 9th, 10th), and sacral spinal nerves S2,3,4
Sympathetic: Thoracolumbar: Originate in the thoracic & lumbar regions of the spinal cord (T1-T12; L1-L2,3)

DISTRIBUTION OF THE SYMPATHETIC AND PARASYMPATHETIC NERVOUS SYSTEM

Three divisions of the ANS:

*sympathetic nervous system,
parasympathetic nervous system
and enteric nervous system*



THE AUTONOMIC NERVOUS SYSTEM

Somatic nervous system: Controls organs under voluntary control (mainly skeletal muscles)


Autonomic Nervous System (ANS): Not under voluntary control. It regulates individual organ, visceral functions and homeostasis, known as the visceral or automatic system. Effectors includes cardiac, smooth muscles and glands.

Helps to adapt the changes in environment. Adjusts or modifies functions in response to stress such as blood pressure, body temperature, sweating etc.

COMPARISON OF AUTONOMIC AND SOMATIC SYSTEMS

Somatic system

- One motor neuron extends from the CNS to skeletal muscle
- Axons are well myelinated
- Conduct impulses rapidly



Cause of demyelination

- Inflammatory processes
- Viral demyelination
- Metabolic derangements
- Hypoxic–ischemic demyelination
- Focal compression.
- Multiple sclerosis
- Acute encephalomyelitis


COMPARISON OF AUTONOMIC AND SOMATIC MOTOR SYSTEMS

Autonomic nervous system


- Chain of two motor neurons
 - Preganglionic neuron
 - Postganglionic neuron
- Conduction is slower due to thin or unmyelinated axons




COMPARISON OF AUTONOMIC AND SOMATIC MOTOR SYSTEMS



DIFFERENCES IN SYMPATHETIC AND PARASYMPATHETIC DIVISIONS



- Sympathetic – long postganglionic fibers
- Parasympathetic – short postganglionic fibers
- Sympathetic axons – highly branched
- Parasympathetic axons – few branches



Sympathetic and parasympathetic systems are consists of myelinated pre-ganglionic fibers which make synaptic connections with un-myelinated postganglionic fibers and then innervate the effector organ. These synapses usually occur in clusters called ganglia.

DIFFERENCES IN SYMPATHETIC AND PARASYMPATHETIC DIVISIONS

Preganglionic neuron:

- ❑ Cell body in brain or spinal cord
- ❑ Axon is myelinated type fiber that extends to autonomic ganglion

Postganglionic neuron:

- ❑ Cell body lies outside the CNS in an autonomic ganglion
- ❑ Axon is unmyelinated type fiber that terminates in a visceral effector

THE AUTONOMIC NERVOUS SYSTEM

The ANS is predominantly an efferent system transmitting impulses from the Central Nervous System (CNS) to peripheral organ systems.

Its effects include:









- Control of heart rate and force of contraction
- Constriction and dilatation of blood vessels
- Contraction and relaxation of smooth muscle
- Visual accommodation
- Secretions from exocrine and endocrine glands.

THE AUTONOMIC NERVOUS SYSTEM

ANS activated by centers located in the **spinal cord, brain stem, hypothalamus** and also **cerebral cortex** especially the **limbic cortex** can transmit signals to the lower centers, influence autonomic control.

ANS operates by visceral reflexes. *Subconscious sensory signals from a visceral organ enter the autonomic ganglia, brain stem or hypothalamus and then return subconscious reflex responses directly back to the visceral organ to control its activities.*

DIFFERENCES IN SYMPATHETIC AND PARASYMPATHETIC DIVISIONS

	central nervous system	peripheral nervous system		target organs
somatic nervous system		 myelin sheath ACh		skeletal muscle
sympathetic nervous system		 ACh paravertebral or prevertebral ganglion	 NE	smooth muscle glands
para-sympathetic nervous system		 myelin sheath	 ACh terminal ganglion	
	ACh = acetylcholine NE = norepinephrine	pre-ganglionic axon	ganglion	post-ganglionic axon

NEUROTRANSMITTERS OF AUTONOMIC NERVOUS SYSTEM



Neurotransmitter released by pre-ganglionic axons

- Acetylcholine for both branches (cholinergic)

Neurotransmitter released by postganglionic axons

- Sympathetic – most release norepinephrine (adrenergic)
- Parasympathetic – release acetylcholine

Feature	Sympathetic	Parasympathetic
Origin of pre-ganglionic fibers	Thoracolumbar nerves	Craniosacral nerves
Location of ganglia	Far from visceral effector organs; in sympathetic chain or collateral ganglia	Near or within viscera effector organs
Neurotransmitter	In ganglia, acetylcholine; in effector organs, norepinephrine	In ganglia, acetylcholine; in effector organs, acetylcholine



THE AUTONOMIC NERVOUS SYSTEM

Subdivision	Nerves Employed	Location of Ganglia	Chemical Messenger	General Function
Sympathetic	Thoracolumbar	Alongside vertebral column	Norepinephrine	Fight or flight
Parasympathetic	Craniosacral	On or near an effector organ	Acetylcholine	Conservation of body energy

THE AUTONOMIC NERVOUS SYSTEM


Organization of the Autonomic Nervous System

Characteristics	Sympathetic Division	Parasympathetic Division	Somatic Nervous System*
Origin of preganglionic neurons	Spinal cord segments T1–L3 (thoracolumbar)	Nuclei of CN III, VII, IX, and X; spinal cord segments S2–S4 (craniosacral)	—
Location of autonomic ganglia	Paravertebral and prevertebral	In or near effector organs	—
Length of preganglionic axons	Short	Long	—
Length of postganglionic axons	Long	Short	—
Effector organs	Smooth muscle; cardiac muscle; glands	Smooth muscle; cardiac muscle; glands	Skeletal muscle
Neurotransmitter and receptor type in ganglion	ACh/nicotinic receptor	ACh/nicotinic receptor	—
Neurotransmitter in effector organs	Norepinephrine (except sweat glands)	ACh	ACh
Receptor types in effector organs	α_1 , α_2 , β_1 , β_2	Muscarinic	Nicotinic


ACh, Acetylcholine; CN, cranial nerve.

DISTRIBUTION OF THE SYMPATHETIC AND PARASYMPATHETIC NERVOUS SYSTEM

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.



FUNCTIONS OF THE SYMPATHETIC AND PARASYMPATHETIC NERVOUS SYSTEM



FUNCTIONS OF THE SYMPATHETIC AND PARASYMPATHETIC NERVOUS SYSTEM


Structure	Sympathetic Stimulation	Parasympathetic Stimulation
Iris (eye muscle)	Pupil dilation	Pupil constriction
Salivary Glands	Saliva production reduced	Saliva production increased
Oral/Nasal Mucosa	Mucus production reduced	Mucus production increased
Heart	Heart rate and force increased	Heart rate and force decreased
Lung	Bronchial muscle relaxed	Bronchial muscle contracted
Stomach	Peristalsis reduced	Gastric juice secreted; motility increased
Small Intes	Motility reduced	Digestion increased
Large Intes	Motility reduced	Secretions and motility increased
Liver	Increased conversion of glycogen to glucose	
Kidney	Decreased urine secretion	Increased urine secretion
Bladder	Wall relaxed Sphincter closed	Wall contracted Sphincter relaxed

THE SYMPATHETIC NERVOUS SYSTEM



THE SYMPATHETIC NERVOUS SYSTEM

Fight-or-flight Response



THE SYMPATHETIC NERVOUS SYSTEM

FEAR, FIGHT- FLIGHT RESPONSE

- ❑ The sympathetic system enables the body to be prepared for fear, flight or fight
- ❑ Sympathetic responses include an increase in heart rate, blood pressure and cardiac output
- ❑ Diversion of blood flow from the skin and splanchnic vessels to those supplying skeletal muscle
- ❑ Increased pupil size, bronchiolar dilation, contraction of sphincters and metabolic changes such as the mobilisation of fat and glycogen.

THE SYMPATHETIC NERVOUS SYSTEM

Frequently referred to as the **fear, fight or flight response**

It has a stimulatory effect on organs and physiological systems, responsible for rapid sensory activity (pupils in the eye) and movement (skeletal muscle).

It diverts blood flow away from the GIT and skin via vasoconstriction.

Blood flow to skeletal muscles, lungs is not only maintained, but enhanced (by as much as 1200%), in case of skeletal muscles.

THE SYMPATHETIC NERVOUS SYSTEM

Dominance by the sympathetic system is caused by physical or emotional stress “E situations”

Emergency, Embarrassment, Excitement, Exercise

Alarm reaction = flight or fight response:

- Dilation of pupils
- Increase heart rate, force of contraction & BP
- Decrease in blood flow to nonessential organs
- Increase in blood flow to skeletal & cardiac muscle
- Airways dilate & respiratory rate increases
- Blood glucose level increase

THE PARASYMPATHETIC DIVISION

- ❑ The parasympathetic nervous system has "rest and digest" activity.
- ❑ Concerned with conservation and restoration of energy, as it causes a reduction in heart rate and blood pressure, and facilitates digestion and absorption of nutrients, and consequently the excretion of waste products
- ❑ The chemical transmitter at both pre and postganglionic synapses in the parasympathetic system is Acetylcholine (Ach).

THE PARASYMPATHETIC DIVISION

Enhance “rest-and-digest” activities

Normally dominate over sympathetic impulses

SLUDD type responses: salivation, lacrimation, urination, digestion & defecation

3 “Decreases” decreased HR, diameter of airways and diameter of pupil

- Paradoxical fear when there is no escape route or no way to win causes massive activation of parasympathetic division
loss of control over urination and defecation

THE AUTONOMIC NERVOUS SYSTEM

Acetylcholine activates mainly two types of *receptors*. They are called *muscarinic* and *nicotinic* receptors.

Muscarine activates only muscarinic receptors whereas nicotine activates only nicotinic receptors; **acetylcholine activates both of them.**

Muscarinic receptors are found on all effector cells that are stimulated by the postganglionic cholinergic neurons of either the parasympathetic nervous system or the sympathetic system.

Nicotinic receptors are found in the autonomic ganglia at the synapses between the preganglionic and postganglionic neurons of both the sympathetic and parasympathetic systems.

THE AUTONOMIC NERVOUS SYSTEM

	<u>Sympathetic</u> (exceptions)	(adrenergic, with	<u>Parasympathetic</u> (<u>muscarinic</u>)
<u>circulatory system</u>			
<u>cardiac output</u>	increases		M2: decreases
<u>SA node: heart rate</u> (<u>chronotropic</u>)	$\beta 1, \beta 2$: increases		M2: decreases
<u>cardiac muscle:</u> <u>contractility</u> (<u>inotropic</u>)	$\beta 1, \beta 2$: increases		M2: decreases (<u>atria</u> only)
<u>conduction at AV node</u>	$\beta 1$: increases		M2: decreases
<u>vascular smooth</u> <u>muscle</u>	M3: contracts; $\alpha =$ contracts; $\beta 2 =$ relaxes		---
<u>platelets</u>	$\alpha 2$: aggregates		---
<u>mast cells - histamine</u>	$\beta 2$: inhibits		---

THE AUTONOMIC NERVOUS SYSTEM

Sympathetic (adrenergic)

Parasympathetic (muscarinic)

respiratory system

smooth muscles of bronchioles

β 2: relaxes (major contribution); α 1: contracts (minor contribution)

M3: contracts

nervous system

pupil of eye

α 1: relaxes

M3: contracts

ciliary muscle

β 2: relaxes

M3: contracts

THE AUTONOMIC NERVOUS SYSTEM

	<u>Sympathetic</u> (<u>adrenergic</u> , with exceptions)	<u>Parasympathetic</u> (<u>muscarinic</u>)
<u>digestive system</u>		
<u>salivary glands</u> : secretions	β : stimulates viscous, <u>amylase</u> secretions; $\alpha 1$ = stimulates <u>potassium</u> cation	stimulates watery secretions
<u>lacrimal glands</u> (tears)	decreases	M3: increases
<u>kidney</u> (<u>renin</u>)	secretes	---
<u>parietal cells</u>	---	M1: secretion
<u>liver</u>	$\alpha 1, \beta 2$: <u>glycogenolysis</u> , <u>gluconeogenesis</u>	---
<u>GI tract</u> motility	decreases	M1, M3: increases
<u>smooth muscles</u> of <u>GI tract</u>	$\alpha, \beta 2$: relaxes	M3: contracts
<u>sphincters</u> of <u>GI tract</u>	$\alpha 1$: contracts	M3: relaxes

THE AUTONOMIC NERVOUS SYSTEM

Sympathetic (adrenergic)

Parasympathetic (muscarinic)

ENDOCRINE

pancreas (islets)

$\alpha 2$: decreases secretion

adrenal medulla

N: secretes epinephrine

urinary system

bladder wall

$\beta 2$: relaxes

contracts

ureter

$\alpha 1$: contracts

relaxes

sphincter

$\alpha 1$: contracts; $\beta 2$ relaxes

relaxes

sweat gland

M: stimulates (major contribution);

secretions

$\alpha 1$: stimulates (minor contribution)

arrector pili

$\alpha 1$: stimulates

THANK YOU

