



MED437
KING SAUD UNIVERSITY



Cranial Nerves 9th & 10th

Lecture (8)

Please check our [Editing File](#)

هذا العمل مبني بشكل أساسي على عمل دفعة ٤٣٦ مع المراجعة والتدقيق وإضافة الملاحظات ولا يعني عن المصدر الأساسي للمذاكرة


- **Important**
 - **Doctors Notes**
 - Notes/Extra explanation
- {وَمَنْ يَتَوَكَّلْ عَلَى اللَّهِ فَهُوَ حَسْبُهُ}

Objectives

By the end of the lecture, the student will be able to:

- ✓ Define the deep origin of both Glossopharyngeal and Vagus Nerves.
- ✓ Locate the exit of each nerve from the brain stem.
- ✓ Describe the course and distribution of each nerve .
- ✓ List the branches of both nerves.

Extra Slide: Mnemonics And Pictures To Help Memorise The Cranial Nerves



I Olfactory Smell	II Optic Vision	III Oculomotor Upward Medial Downward Up and In
IV Trochlear Down and In	V Trigeminal Touch Forehead and Cheek Clench Teeth	VI Abducens Look Side to Side
VII Facial Taste for the Anterior 2/3 of Tongue Smile	VIII Acoustic Hearing Equilibrium	IX Glossopharyngeal Posterior 1/3 of the Tongue Speech
X Vagus Defecation Slowed Heart Rate	XI Spinal Accessory Shoulder Shrug	XII Hypoglossal Tongue Movement

CRANIAL NERVE MNEMONIC


S = Sensory M = Motor B = Both

O Olfactory	O On	S Some
O Optic	O Old	S Say
O Oculomotor	O Olympus	M Marry
T Trochlear	T Towering	M Money
T Trigeminal	T Tops	B But
A Abducens	A A	M My
F Facial	F Finn	B Brother
A Acoustic	A And	S Says
G Glossopharyngeal	G German	B Big
V Vagus Nerve	V Viewed	B Brains
S Spinal	S Some	M Matter
H Hypoglossal	H Hops	M More



Glossopharyngeal (IX) 9th Cranial Nerve

- It is principally a **Sensory** nerve with preganglionic parasympathetic and few motor fibers (it is a mixed nerve but most of the fibers are sensory).
- It has no real nucleus to itself. Instead it shares nuclei with VII (facial) and X (vagus).




Each cranial nerve has superficial attachment (fibers in ventral surface of brain stem) and deep attachment (origin of nuclei)




Glossopharyngeal (IX) 9th Cranial Nerve

Superficial Attachment (مكان خروجه من البيرين ستيم)

- It arises from the ventral aspect of the medulla by a linear series of small rootlets, in groove between olive and inferior cerebellar peduncle.
- It leaves the cranial cavity by passing through the jugular foramen in company with the Vagus (10), Acessory (11) nerves and the Internal jugular vein.




Extra



Lateral view of Jugular Foramen




Extra



Extra

Glossopharyngeal (IX) 9th Cranial Nerve Course (extracranial)

- It Passes forwards between **Internal jugular vein** and **External carotid artery**.
- Lies Deep to **Styloid process**.
- Passes between external and internal carotid arteries at the posterior border of **Stylopharyngeus** then lateral to it.
- It reaches the pharynx by passing between middle and inferior constrictors, deep to **Hyoglossus**, where it breaks into terminal branches.




Glossopharyngeal (IX) 9th Cranial Nerve


Component of fibers & Deep origin (Deep origin = nuclei)

Recall: **S**A**M**E
Sensory → Afferent
Motor → Efferent

<u>SVE</u> fibers: Special Visceral Efferent	originate from nucleus ambiguus (NA) motor fiber	supply <i>stylopharyngeus</i> muscle
<u>GVE</u> fibers: General Visceral Efferent	arise from inferior salivatory nucleus (ISN) or another name is (dorsal motor nucleus) Parasympathetic	relay in otic ganglion , the postganglionic fibers supply <i>parotid gland</i>
<u>SVA</u> fibers: Special Visceral Afferent	arise from the cells of inferior ganglion, their central processes terminate in nucleus of solitary tract (NST) sensory	the peripheral processes supply the taste buds on posterior third of tongue.
<u>GVA</u> fibers: General Visceral Afferent	end in nucleus of solitary tract.	visceral sensation (pain and temp.) from mucosa of <i>posterior third of tongue, pharynx, auditory tube and tympanic cavity, carotid sinus</i>



IX. GLOSSOPHARYNGEAL
 GSA GVA GVE SVA SVE



Glossopharyngeal (IX) 9th Cranial Nerve Ganglia & Communications


It has two ganglia: Same dorsal root ganglia (termination)

○ **Superior ganglion:** Superior → Small

- Small, with no branches.
- It is connected to the **Superior Cervical sympathetic ganglion**.



○ **Inferior ganglion:**

- Large and carries general sensations from pharynx, soft palate and tonsil.
- It is connected to Auricular Branch of Vagus.
- The Trunk of the nerve is connected to the Facial nerve at the **stylomastoid foramen**




Glossopharyngeal (IX) 9th Cranial Nerve Branches

1. **Tympanic:** *parasympathetic* relays in the otic ganglion and gives secretomotor to the *parotid gland*
 2. **Nerve to Stylopharyngeus** muscle. (*SVE fiber*)
 3. **Pharyngeal:** to the mucosa of pharynx.
 4. **Tonsillar.**
 5. **Lingual :**
- } (*GVA fiber*)
6. carries sensory branches, general and special (taste) from the posterior third of the tongue. (*SVA fiber*)
- **Sensory branches** from the carotid sinus and body (**pressoreceptors and chemoreceptors**).




Glossopharyngeal (IX) 9th Cranial Nerve Nerve Lesions



It produces:

- Difficulty of swallowing; Impairment of **taste** and **sensation** over the *posterior one-third* of the tongue ,palate and pharynx.
- **Absent gag reflex**. Dysfunction of the parotid gland (*it is salivary gland, dysfunction will lead to dry mouth*).




How to test for IX injury?

- Have the patient open the mouth and inspect the palatal arch on each side for asymmetry.
- Use a tongue blade to depress the base of the tongue gently if necessary.
- Ask the patient to say "**ahhh**" as long as possible. Observe the palatal arches as they contract and the soft palate as it swings up and back in order to close off the nasopharynx from the oropharynx.
- Normal palatal arches will constrict and elevate, and the uvula will remain in the midline as it is elevated. With paralysis there is no elevation or constriction of the affected side.
- warn the patient that you are going to test the gag reflex. Gently touch first one and then the other palatal arch with a tongue blade, waiting each time for gagging.

Glossopharyngeal (IX) 9th Cranial Nerve

Summary


Glossopharyngeal Nerve and Branches



A mixed nerve, it carries motor fibers to pharyngeal muscles for swallowing and parasympathetic motor fibers to salivary glands. Sensory fibers carry messages from the pharynx, tonsils, posterior of tongue (taste). Glossopharyngeal fibers also carry afferent messages from the carotid sinus baroreceptors.

Vagus (X) 10th Cranial Nerve




- It is a **Mixed** nerve.
- Its name means wandering حائر (it goes all the way to the abdomen)
- So it is the *longest* and most *widely distributed* cranial nerve.
- The principal role of the vagus is to provide parasympathetic supply to organs throughout the thorax and upper abdomen.
- It also gives sensory and motor supply to the *pharynx* and *larynx*.



Vagus (X) 10th Cranial Nerve Superficial Attachment & Course

- Its rootlets exit from medulla between olive and inferior cerebellar peduncle.
- Leaves the skull through **jugular foramen**.
- It occupies the posterior aspect of the carotid sheath between the internal jugular vein laterally and the internal and common carotid arteries medially.
- It has two ganglia:
 - Superior ganglion in the jugular foramen
 - Inferior ganglion, just below the jugular foramen

The sensory fibers end in the ganglion before continuing just like the dorsal root ganglion




Vagus (X) 10th Cranial Nerve Communications

○ Superior ganglion with:

- Inferior ganglion of glossopharyngeal nerve,
- Superior cervical sympathetic ganglion &
- Facial nerve.



○ Inferior ganglion with:

- Cranial part of accessory nerve,
- Hypoglossal nerve,
- Superior cervical sympathetic ganglion.
- 1st cervical nerve.



Vagus (X) 10th Cranial Nerve Course (**extracranial**)

- The vagus runs down the neck on the prevertebral muscles and fascia.
- The **internal jugular vein** lies behind it, and the **internal** and **common carotid** arteries are in front of it, all the way down to the superior thoracic aperture.
- It lies on the prevertebral muscles and fascia.
- Enters thorax through its **inlet**:
 - *Right Vagus* descends in front of the **right subclavian artery**.
 - *Left Vagus* descends between the **left common carotid** and **subclavian arteries**.




Vagus (X) 10th Cranial Nerve

Components of fibers & Deep origin

Recall: **S**AME
Sensory → Afferent
Motor → Efferent

<p><u>SVE</u> fibers: Special Visceral Efferent (Motor)</p>	originate from Nucleus Ambiguus,	to muscles of pharynx and larynx (and soft palate)
<p><u>GVE</u> fibers: General Visceral Efferent (Preganglionic parasympathetic)</p>	originate from Dorsal Nucleus of Vagus synapses in parasympathetic ganglia,	short postganglionic fibers innervate cardiac muscle, smooth muscles and glands of viscera.
<p><u>SVA</u> fibers: Special Visceral Afferent</p>	to Spinal Tract & Nucleus of Trigeminal	sensation from auricle, external acoustic meatus and cerebral dura mater*,
<p><u>GVA</u> fibers: General Visceral Afferent (sensory)</p>	to Nucleus of Solitary Tract	carry impulse from viscera in neck, thoracic and abdominal cavities




When stretches causes headache*

Vagus (X) 10th Cranial Nerve

Branches (The fibers will give branches)

stimulation of auricular will also stimulate cardiac branch to heart so trauma to the ear may lead to tachycardia and cardiac arrest

1. Meningeal sensory	Dura (SVA)
2. Auricular nerve Sensory + parasympathetic	external acoustic meatus, and tympanic membrane.
3. Pharyngeal (enters the wall of the pharynx) Sensory + Motor + para	mucous membrane of the pharynx, superior and middle constrictor muscles, all the muscles of the palate except the tensor palate .
4. To carotid body + sinus (sensory)	
5. Superior Laryngeal: It divides into: (1) Internal Laryngeal : sensory	provides sensation to the hypopharynx(back), the epiglottis, and the part of the larynx that lies above the vocal folds
(2) External Laryngeal : motor	supplies the cricothyroid muscle




Vagus (X) 10th Cranial Nerve Branches

6. Recurrent Laryngeal : (motor + sensory)

- it goes round the subclavian artery on the right, and round the arch of the aorta on the left
- Below the vocal cords
- It runs upwards and medially alongside the trachea, and passes behind the lower pole of the thyroid gland.

motor supply to **all the muscles of the larynx**, except the **cricothyroid**. It also provides sensation *to the larynx below the vocal folds*.


Injury will in **thyroidectomy will lead to hoarseness of voice or complete loss of voice**



Vagus (X) 10th Cranial Nerve Lesions



- Vagus nerve lesions produce palatal and pharyngeal and laryngeal paralysis.
- Abnormalities of esophageal motility, gastric acid secretion, gallbladder emptying, and heart rate; and other autonomic dysfunction.
- Hoarseness of voice




How to diagnose X nerve Injury?

- Listen to the patient talk as you are taking the history.
- Hoarseness, whispering, nasal speech, or the complaint of aspiration or regurgitation of liquids through the nose (since soft palate is not working) should make you especially mindful of abnormality.
- Give the patient a glass of water to see if there is choking or any complaints as it is swallowed.
- Laryngoscopy is necessary to evaluate the vocal cord (to assess movement).

Vagus (X) 10th Cranial Nerve Summary

- X is a **mixed** nerve.
- It contains afferent, motor, and parasympathetic fibers.
- The afferent fibers convey information from:
esophagus, tympanic membrane, external auditory meatus and part of cochlea of the middle ear.
End in trigeminal sensory nucleus.
- Chemoreceptors in aortic bodies and baroreceptors in aortic arch.
- Receptors from thoracic & abdominal viscera, end in **nucleus solitarius**.
- The motor fibers arise from **nucleus ambiguus** of medulla to innervate muscles of soft palate, pharynx, larynx, and upper part of esophagus.
- The parasympathetic fibers originate from dorsal motor nucleus of vagus in medulla distributed to cardiovascular, respiratory, and gastrointestinal systems (up to right 2/3 the rest is from sacral fibers s2-s4).



Causes of both IX & X nerve lesions:

1. Lateral medullary syndrome:

A degenerative disorder seen over age of 50 mostly due to ***Thrombosis of the Inferior Cerebellar Artery.***

2. Tumors compressing the cranial nerves in their exiting foramina from the cranium via the skull base

Manifested by:

- Ipsilateral* paralysis of the muscles of the Palate, Pharynx and Larynx.
- Ipsilateral loss of Taste from the Posterior Third of tongue.

*the same side of the body

Pharyngeal plexus: 9th,10th,11th CNs and C1 sympathetic ganglia





Figure X-12 Tumor of the glomus cells of the jugular bulb compressing cranial nerves IX, X, and XI (lateral view showing cut jugular foramen).

Extra

Nerve	Glossopharyngeal (9th)	Vagus (10th)
Type	Sensory (mainly)	Mixed
Exits from	Ventral medulla (b/w olive and inferior cerebellar peduncle)	Ventral medulla (b/w olive and inferior cerebellar peduncle)
Leaves cranial cavity via	Jugular foramen	Jugular foramen
Nuclei	<ol style="list-style-type: none"> 1. Nucleus ambiguus 2. Inferior salivatory nucleus 3. Nucleus of solitary tract 	<ol style="list-style-type: none"> 1. Nucleus ambiguus 2. Dorsal nucleus of vagus 3. Spinal tract & nucleus of trigeminal 4. Nucleus of solitary tract
Branches	<ol style="list-style-type: none"> 1. Tympanic 2. Nerve to stylopharyngeus muscle 3. Pharyngeal 4. Tonsillar 5. Lingual 6. Sensory (from carotid sinus and body) 	<ol style="list-style-type: none"> 1. Meningeal 2. Auricular nerve 3. Pharyngeal 4. To carotid body 5. Superior laryngeal (internal and external laryngeal) 6. Recurrent laryngeal
Nerve lesion manifestation	<ol style="list-style-type: none"> 1. Difficulty swallowing 2. Impairment of taste and sensation of posterior 1/3 of the tongue, palate and pharynx. 3. Absent gag reflex 4. Dysfunction of parotid gland 	<ol style="list-style-type: none"> 1. Palatal, pharyngeal, and laryngeal paralysis. 2. Abnormalities of: <ul style="list-style-type: none"> • esophageal motility gastric acid secretion • gall bladder emptying • heart rate 3. Other autonomic dysfunction
Cause of nerve lesion	<ol style="list-style-type: none"> 1. Lateral medullary syndrome 2. Tumors compressing the cranial nerves in their exit 	

MCQs

1. Glossopharyngeal shares its nuclei with:

- A- CN 7 & 8
- B- CN 7 & 10
- C- CN 8 & 10
- D- CN 11

Answer: B

2. Cranial nerves 9 and 10 exit the cranial cavity through:

- A- foramen magnum
- B- foramen ovale
- C- jugular foramen
- D- carotid foramen

Answer: C

3. The tympanic branch of glossopharyngeal supplies:

- A- parotid gland
- B- pineal gland
- C- pituitary gland
- D- sublingual gland

Answer: A

4. A patient was shown to have absent gag reflex. Which nerve is most likely affected:

- A- facial
- B- vagus
- C- glossopharyngeal

Answer: C

5. The right vagus nerve descends:

- A- in front right subclavian artery.
- B- in front of right subclavian vein .
- C- between left common carotid and subclavian arteries.
- D- between right common carotid and subclavian arteries.

Answer: A

6. Which of the following nuclei give fibers to muscles of pharynx and larynx?

- A- nucleus solitarius
- B- nucleus ambiguus
- C- nucleus of trigeminal

Answer: B

7. Cricothyroid is supplied by which branch of the vagus nerve:

- A- internal laryngeal
- B- external laryngeal
- C- recurrent laryngeal

Answer: B

8. CN 9 & 10 lesions can be caused by:

- A- lateral medullary syndrome
- B- medial medullary syndrome
- C- anterior medullary syndrome
- D- posterior medullary syndrome

Answer: A

9. Is a branch from the vagus nerve:

A- tympanic

B- meningeal

C- tonsillar

D- nerve to stylopharyngeus

Answer: B

2. The vagus nerve is a :

A- sensory nerve

B- motor nerve

C- mixed nerve

Answer: C

SAQ:

1- list three branches from the vagus?

Meningeal, Auricular nerve and Pharyngeal

2-list the three nuclei of **Glossopharyngeal (9th)**

Nucleus ambiguous, Inferior salivatory nucleus and Nucleus of solitary tract



Good luck
Special thank for team436 ❤️

Team Leaders:

**Faisal Fahad Alsaif
Rawan Mohammad Alharbi**

Team Members:

**Abdulaziz Aldukhayel
Abdulrahman Alduhayyim
Rinad Alghoraiby
Rawan Mishal**

- References:
 1. Girls' & Boys' Slides
 2. Greys Anatomy for Students
 3. TeachMeAnatomy.com

 [Twitter.com/Anatomy437](https://twitter.com/Anatomy437)

 Anatomyteam.437@gmail.com