



# Esophageal motility

## Objectives :

- ❖ Mastication and chewing reflex
- ❖ Swallowing process and its stages
- ❖ Types of esophageal peristalsis
- ❖ Esophageal sphincter
- ❖ Function of lower esophageal sphincter
- ❖ Achalasia
- ❖ Incompetence of lower esophageal sphincter

## Done by :

- Team leader: Rahaf AlShammari
- Team members:
  - ◆ Majd AlBarrak, Shahad AlZahrani
  - ◆ Khalid AlMutairi, Mohammed AlHassan
  - ◆ Abdullah AlSargani, Abdullah AlZaid
  - ◆ Wejdan AlBadrani, Maha AlAmri
  - ◆ Hadeel AlMukainzi, Shahad AlTayash
  - ◆ Norah AlKadi




## Colour index:

- Important
- Numbers
- Extra

# Mastication (Chewing)

## Functions:

1. To lubricate the bolus with salivary secretion.
2. To breakdown the bolus to small particles.
3. To begin digestion of carbohydrate (by amylase).




## Teeth organization:

- Anterior teeth (incisors) for cutting.
- Posterior teeth (molars) for grinding.

## Chewing muscles are innervated by

- Masseter
- Temporalis
- Pterygoid (lateral & medial)




## nerve V (trigeminal nerve):

**Taste center (Hypothalamus + Brainstem)** → rhythmical chewing movements (Chewing reflex).

## Chewing (stretch) reflex:

The presence of a bolus of food in the mouth initiates reflex inhibition of the muscles of mastication → the lower jaw drops → stretch reflex of the jaw muscles & contraction → jaw closure & compression of the bolus again against the linings of the mouth → inhibits the jaw muscles once again, allowing the jaw to drop and repeat.




Very useful video


# Swallowing (Deglutition)

**Swallowing** is The ordered sequence of events that propel food from the mouth to the stomach.

- Initiatarion is voluntary, thereafter it is entirely under reflex control.
- Inhibits respiration and prevents the entrance of food into the airway passages.
- The reflex portion of swallowing is controlled by the **swallowing center** in the medulla.




## Stages of Swallowing



### I- The Oral Stage:

- This stage involves the voluntary rolling of the chewed food (bolus) posteriorly into the pharynx by the upward and backward pressure applied by the tongue against the palate.
- The activation of somatosensory receptors in the pharynx initiates the involuntary swallowing reflex in the medulla. From here on, swallowing becomes entirely automatic and can not be stopped.



## II- The Pharyngeal Stage:



At the pharynx, the bolus stimulates epithelial swallowing receptor areas all around the pharynx opening and impulses pass to swallowing center and accordingly initiate a series of autonomic pharyngeal muscle contractions as follows:

The soft palate is pulled upward to close the posterior nares which prevents the food from entering the nasal cavities.

The palatopharyngeal folds pull medially to approximate and form a sagittal slit through which food must pass into the posterior pharynx.

Form a passage for the food.



The vocal cords are strongly approximated and the larynx is pulled upward and anteriorly by the neck muscles. These actions cause the epiglottis to swing backward over the opening of the larynx.


The upward movement of the larynx pulls up and enlarges the opening to the esophagus. The upper esophageal sphincter relaxes and food move into the upper esophagus.

Once the larynx is raised and the pharyngoesophageal sphincter relaxes, the entire muscular wall of the pharynx contracts propelling the food by peristalsis into the esophagus. Swallowing center inhibits the respiratory center.



عشان كذا لو تكلمنا واحنا ناكل  
مممكن نشرق

## Summary of the Pharyngeal stage:

- The trachea is closed.
- The esophagus is opened. "Sphincter relaxed"
- A fast peristaltic wave initiated by the nervous system of the pharynx forces the bolus of food into the upper esophagus .
- Time of process is < 2 second.



## Nervous Initiation of the Pharyngeal Stage of Swallowing



### III- The Esophageal Stage :

- The esophagus is a conduit to move food rapidly from the pharynx to the stomach.
- The esophageal stage is controlled partly by the swallowing reflex and partly by the enteric nervous system (ENS). In case of vagotomy enteric nervous system takes over. Vagotomy: cutting of vagus nerve
- Physiologically, esophagus is divided into three functionally distinct regions:
  - Upper esophageal sphincter (UES)
  - Esophageal Body
  - Lower esophageal sphincter (UES)

The upper 1/3  
of the  
esophagus:

Striated muscle (peristaltic waves are controlled by impulses from IX and X nerves)

The lower 2/3  
of the  
esophagus:

Smooth muscle (controlled by the X nerve through connections with esophageal myenteric nervous system).



The  
swallowing  
reflex closes  
UES:

A peristaltic wave begins just below UES when bolus pass through it and traverse the entire esophagus.

The lower  
esophageal  
sphincter:

Relaxes and the bolus enters the stomach.  
"Receptive relaxation"

# Esophageal peristalsis:



## Types of Esophageal Peristalsis:

### Primary

- A continuation of peristaltic wave that begins in pharynx and spreads into esophagus during the pharyngeal stage of swallowing.
- This wave passes from the pharynx to the stomach in 8-10 sec.
- Coordinated by swallowing center.


### Secondary

- Bolus very big or sticky.
- Occurs in response to distention and continue until all the food is emptied into the stomach.
- ENS and swallowing center are both involved




# Types of Esophageal Peristalsis:

## Primary





## Secondary





# Esophageal Sphincters



## The Upper Esophageal Sphincter (UES)

### Where

It is formed of striated muscle but is not under voluntary control, located at the lower end of pharynx and guards the entrance into the esophagus.

### Function

- It prevents esophageal air insufflation during negative intrathoracic pressure events, as during inspiration.
- It prevents esophagopharyngeal/laryngeal reflux during esophageal peristalsis.

### When Relax

It relaxes during swallowing for about 1 second allowing the bolus to be forced through the relaxed UES

## The lower Esophageal Sphincter (LES)

### Where

It is formed by circular smooth muscle located in an area of ~ 3 cm upward of the junction with the stomach.

### Function

In between swallowing, it normally remains tonically constricted to prevent reflux of gastric contents into esophagus. During swallowing, the peristaltic wave passes down the esophagus and causes a "receptive relaxation" of LES for 7-10 sec. (mediated by vagal impulses) to allow easy propulsion of the swallowed food into the stomach.

### Why

Pressure in the esophagus is mostly -ve (except for a short intra-abdominal segment). So that pressure in the stomach is always higher than the esophagus.

# Causes of Competence and the antireflux functions of the LES

1	Its resting pressure (15- 30 mmHg).
2	The <b>diaphragm</b> wraps around the esophagus at the level of LES. Contraction of the diaphragm during inspiration helps to increase the pressure at the LES.
3	The flutter-valve closure of the lower esophagus by the <b>increased intra-abdominal pressure</b> prevents the high pressure in the stomach from forcing its contents into the esophagus.
4	<b>Valve-like closure</b> of the distal end of the esophagus. This involves a short portion of the esophagus that extends slightly into the stomach and caves the esophagus inward in response to increased intraabdominal P. "Caving of esophagus inside the stomach"


Doctor said: 3 mechanisms behind competency of LES:

- 1-Tonic contraction of smooth muscles.
- 2- Diaphragm contraction and wrapping.
- 3-Valve-like closure.

## Competence and the Antireflux Functions of the LES

### Control of LES Function

Circular musculature of LES is regulated by nerves, (extrinsic & intrinsic), hormones and neuromodulators.



### Contraction of LES


- Between swallows (tonic vagal cholinergic impulses)
- Stimulation of sympathetic nerves
- Gastrin hormone.

### Relaxation of LES


- During swallowing (efferent vagal impulses).
- The transmitter is NO or VIP.
- Secretin and cholecystikinin (CCK)

# Achalasia


Doctor said: Failure of sphincter to relax due to absence of inhibitory neuron (vagus nerve)



Normal



Achalasia



It is due to **pathology of or absence of the myenteric plexus containing VIP & NO** in the lower third of esophagus (may be due to autoimmune destruction of the nerves in esophageal wall).



The musculature of the lower esophagus instead remains contracted and **the myenteric plexus has lost the ability to transmit a signal to cause relaxation of the LES** (high resting pressure).



**Food becomes putridly infected during the long periods of esophageal stasis** causing ulceration of the esophageal mucosa, severe substernal pain or even rupture & death.



**Food transmission from the esophagus into the stomach is impeded or prevented.** As achalasia gets worse, the esophagus gradually enlarged as food collects within it.

## Diagnosis:

- I. Esophageal manometry. "Measure the pressure in the esophagus"
- II. X-rays of upper digestive system (barium swallow).
- III. Upper endoscopy.

## Treatment:

### Surgical

**Myotomy**  
cutting the muscle at the lower end of the esophageal sphincter.



**Pneumatic dilation**  
by balloon insertion into the esophageal sphincter.

### Non-surgical

**Injecting muscle relaxant** (Botox) directly into the esophageal sphincter.

**Medication**  
as muscle relaxants, nitroglycerin before eating.

## Gastroesophageal Reflux Disease (GERD) Incompetence of the LES:



## What is it

## Cause

## Contributing factors

Incompetence of the lower esophageal sphincter allows reflux of gastric contents into the esophagus.

It may result from a generalized loss of intrinsic sphincter tone or from recurrent inappropriate transient relaxations triggered by gastric secretions.

Weight gain, fatty foods, caffeinated or carbonated beverages, alcohol, tobacco smoking, and drugs (anticholinergics, antihistamines, calcium channel blockers, progesterone and nitrates).

## It can lead to

Reflux esophagitis, heartburn, esophageal stricture, ulcer, dysplastic changes (cancerous), or Barrett esophagus (normal tissue lining the esophagus resembles tissue lining of the intestine).


## Diagnosis

Is clinical, sometimes with endoscopy, with or without 24-h pH testing.

## Treatment


Involves lifestyle modification, acid suppression using proton pump inhibitors, and sometimes surgical repair.

# Summary



## Causes of Competence and the antireflux functions of the LES

1. Its resting pressure (15- 30 mmHg).
2. The diaphragm wraps around the esophagus at the level of LES
3. The flutter-valve closure of the lower esophagus by the increased intra-abdominal pressure.
4. Valve-like closure of the distal end of the esophagus.





## MCQs

1- Stages of swallowing:

- A) Oral, pharyngeal, esophageal.
- B) Mastication, oral, pharyngeal, esophageal.
- C) Pharyngeal, esophageal.
- D) Pharyngeal, esophageal, gastric.

2- In the pharyngeal stage:

- A) palatopharyngeal folds pull medially to approximate and form a horizontal slit.
- B) The larynx is pulled upward and anteriorly by the face muscles.
- C) The soft palate is pulled downward to close the posterior nares.
- D) The epiglottis swings backwards over the opening of the larynx.

3-Enlargement of the esophagus due to collection of food within it, is a complication of:

- A) Achalasia
- B) GERD
- C) H Pylori infection
- D) Heavy meal

4- One of the UES functions is:

- A) Prevention of the reflux of gastric contents into the esophagus
- B) Prevention of esophageal air insufflation.
- C) Prevention of the esophagopharyngeal/laryngeal reflux
- D) (C+B)

5- One method to diagnose GERD is

- A) Endoscopy
- B) Xray
- C) Esophageal manometry

6- The esophageal stage is controlled by:

- A) Enteric nervous system
- B) Swallowing reflex
- C) Swallowing reflex+ENS

7- The UES relaxes during swallowing for about:

- A) 7-10 sec
- B) 1 sec
- C) 2 sec
- D) 5 sec

8- The surgical treatment for Achalasia is:

- A) Injecting muscle relaxant (Botox)
- B) Pneumatic dilation by balloon
- C) Nitroglycerin
- D) Myotomy

9- Which of these causes LES contraction:

- A) Secretin
- B) CCK
- C) Gastrin
- D) Nitroglycerin

Answers

- 1- A
- 2- D
- 3- A
- 4- D
- 5- A
- 6- C
- 7- B
- 8- D
- 9- C