


## Endocrine Physiology

# The Endocrine Pancreas

*Dr. Khalid Al-Regaiey*




# Pancreas

---

- A triangular gland, which has both exocrine and endocrine cells, located behind the stomach
- Strategic location
- Acinar cells produce an enzyme-rich juice used for digestion (exocrine product)
- Pancreatic islets (**islets of Langerhans**) produce hormones involved in regulating fuel storage and use.

# The Endocrine Pancreas




# Islets of Langerhans


---

- 1-2 million islets
- Beta ( $\beta$ ) cells produce insulin (60%)
- Alpha ( $\alpha$ ) cells produce glucagon (25%)
- Delta ( $\delta$ ) cells produce somatostatin (10%)
- F cells produce pancreatic polypeptide (5%)

# Islets of Langerhans



CELL	SECRETES:
Alpha cells 	Glucagon
D cells 	Somatostatin
Beta cells	Insulin, amylin



# Insulin


---

- Hormone of nutrient abundance
- A protein hormone consisting of two amino acid chains linked by disulfide bonds
- Synthesized as part of proinsulin (86 AA) and then excised by enzymes, releasing functional insulin (51 AA) and C peptide (29 AA).
- Has a plasma half-life of 6 minutes.



# Insulin Structure

## PROINSULIN



# Insulin Synthesis

---

DNA (chromosome 11) in  $\beta$  cells



mRNA



Preproinsulin (signal peptide, A chain,  
B chain, and peptide C)



proinsulin



insulin




# Insulin Synthesis


---

- Insulin synthesis is stimulated by glucose or feeding and decreased by fasting
- Threshold of glucose-stimulated insulin secretion is 100 mg/dl.
- Glucose rapidly increase the translation of the insulin mRNA and slowly increases transcription of the insulin gene

# Glucose is the primary stimulator of insulin secretion





**(a) Beta cell at rest.** The  $K_{ATP}$  channel is open and the cell is at its resting membrane potential.



**(b) Beta cell secretes insulin.**

Closure of  $K_{ATP}$  channel depolarizes cell, triggering exocytosis of insulin.







# Regulation of Insulin Secretion

## Regulators of insulin secretion

### Stimulators of insulin secretion

↑ Serum glucose

↑ Serum amino acids

↑ Serum free fatty acids

↑ Serum ketone bodies

#### Hormones

Gastroinhibitory peptide (GIP)

Glucagon

Gastrin

Cholecystokinin (CCK)

Secretin

Vasoactive intestinal peptide (VIP)

Epinephrine ( $\beta$ -receptor)

Parasympathetic nervous system

### Inhibitors of insulin secretion

↓ Glucose

↓ Amino acids

↓ Free fatty acids

#### Hormones

Somatostatin

Epinephrine ( $\alpha$ -receptor)


Sympathetic nervous system stimulation



# Insulin Receptor

---

- the insulin receptor is a transmembrane receptor
- belongs to the large class of **tyrosine kinase receptors**
- Made of two alpha subunits and two beta subunits



---

# **Actions of insulin**

# Glucose regulation and metabolism terms

- **Gluconeogenesis** - Synthesis of glucose from noncarbohydrate precursors, Lactic acid, glycerol, amino acids, liver cells synthesis glucose when carbohydrates are depleted.
- **Glycogenesis** - Formation of glycogen, glucose stored in liver and skeletal muscle as glycogen, important energy reserve.
- **Glycogenolysis** – breakdown of glycogen (polysaccharide) into glucose molecules (monosaccharide)
- **Glycolysis** - the breakdown of glucose into pyruvate by cells for the production of ATP

- **Rapid (seconds)**
- (+) transport of glucose, amino acids,  $K^+$  into insulin-sensitive cells
- **Intermediate (minutes)**
- (+) protein synthesis
- (-) protein degradation
- (+) of glycolytic enzymes and glycogen synthase
- (-) phosphorylase and gluconeogenic enzymes
- **Delayed (hours)**
- (+) mRNAs for lipogenic and other enzymes

# Action of insulin on Adipose tissue

---

- (+) glucose entry
- (+) fatty acid synthesis
- (+) glycerol phosphate synthesis
- (+) triglyceride deposition
- (+) lipoprotein lipase
- (-) of hormone-sensitive lipase
- (+) K uptake



# Action of insulin on Muscle:

---

- (+) glucose entry
- (+) glycogen synthesis
- (+) amino acid uptake
- (+) protein synthesis in ribosomes
- (-) protein catabolism
- (-) release of gluconeogenic amino acids
- (+) ketone uptake
- (+) K uptake

# Action of insulin on Liver:

---

- (-) ketogenesis
- (+) protein synthesis
- (+) lipid synthesis
- (-) gluconogenesis, (+) glycogen synthesis, (+) glycolysis.

# General

---


- (+) cell growth

# Glucose Transport

---

- GLUT1 (erythrocytes, brain)
- GLUT2 (liver, pancreas, small intestines, kidney)
- GLUT3 (brain)
- **GLUT4**, insulin sensitive transporter (muscle, adipose tissue)

# Insulin: Summary



**Table 22-3: Insulin**

Cell of origin	Beta cells of pancreas
Chemical nature	51-amino acid peptide
Biosynthesis	Typical peptide
Transport in the circulation	Dissolved in plasma
Half-life	5 minutes
Factors affecting release	Plasma [glucose] > 100 mg/dL; ↑ blood amino acids; GI hormones (feedforward reflex) and parasympathetic amplify. Sympathetic inhibits.
Target cells or tissues	Liver, muscle, and adipose tissue primarily; brain, kidney, and intestine not insulin-dependent
Target receptor	Membrane receptor with tyrosine kinase activity; pathway with insulin-receptor substrates
Whole body or tissue action	↓ Plasma [glucose] by ↑ transport into cells or ↑ metabolic use of glucose
Action at cellular level	↑ Glycogen synthesis; ↑ aerobic metabolism of glucose; ↑ protein and triglyceride synthesis



# Glucagon

---

- A 29-amino-acid polypeptide hormone that is a potent hyperglycemic agent
- Produced by  $\alpha$  cells in the pancreas

# SYNTHESIS

---

DNA in  $\alpha$  cells (chromosome 2)



mRNA



Preproglucagon



proglucagon



glucagon

# Factors Affecting Glucagon Secretion:

## ***Effects on Glucagon Secretion***

### **Stimuli for Glucagon Secretion**

↓ Blood glucose

↑ Serum amino acids (arginine, alanine)

Sympathetic nervous system stimulation

Stress

Exercise

### **Inhibitors of Glucagon Secretion**

Somatostatin

Insulin


↑ Blood glucose

# Glucagon Actions

---

- Its major target is **liver**:
  - **Glycogenolysis**
  - **Gluconeogenesis**
  - **Lipid oxidation** (fully to CO<sub>2</sub> or partially to produce keto acids “**ketone bodies**”).
  - Release of glucose to the blood from liver cells

# Glucagon Action on Cells:




**Table 22-5: Glucagon**

Cell of origin	Alpha cells of pancreas
Chemical nature	29-amino acid peptide
Biosynthesis	Typical peptide
Transport in the circulation	Dissolved in plasma
Half-life	4–6 minutes
Factors affecting release	Stimulated by plasma [glucose] < 200 mg/dL, with maximum secretion below 50 mg/dL; ↑ blood amino acids.
Target cells or tissues	Liver primarily
Target receptor/second messenger	G protein-coupled receptor linked to cAMP
Whole body or tissue action	↑ Plasma [glucose] by glycogenolysis and gluconeogenesis; ↑ lipolysis leads to ketogenesis in liver
Action at molecular level	Alters existing enzymes and stimulates synthesis of new enzymes
Feedback regulation	↑ Plasma [glucose] shuts off glucagon secretion
Other information	Member of secretin family along with VIP, GIP, and GLP-1



# The Regulation of Blood Glucose Concentrations





## *Diabetes*

Olivia has a lot of baby fat on her leg so it's a good place to give her a shot.

~Olivia's Mom



# Diabetes

---

- Diabetes is probably the most important metabolic disease.
- It affects every cell in the body and affects carbohydrate, lipid, and protein metabolism.
- characterized by the polytriad:
  - **Polyuria** (excessive urination)
  - **Polydypsia** (excessive thirst)
  - **Polyphagia** (excessive hunger).

# Types of Diabetes

---

## **Type 1** Diabetes

Affects children

### **Cause:**

inadequate  
insulin secretion

### **Treatment :**

insulin injection

## **Type 2** diabetes


Affects adults

**Cause** defect in  
insulin action


### **Treatment :**

diet or OHA

# ***Diabetes Mellitus***



# Type 1 diabetes



# Diabetes Mellitus Type I

- Caused by an immune-mediated selective destruction of  $\beta$  cells
- $\beta$  cells are destroyed while  $\alpha$  cells are preserved:  
No insulin :::: high glucagon  $\Rightarrow$  high production of glucose and ketones by liver  
glucose & ketones  $\uparrow$   $\Rightarrow$  osmotic diuresis  
keto acids  $\uparrow$   $\Rightarrow$  diabetic ketoacidosis

# Diabetes Mellitus: Type II

---

- More common in some ethnic groups
- Insulin resistance keeps blood glucose too high
- Chronic complications: atherosclerosis, renal failure & blindness

# Long Term Complications of Uncontrolled Diabetes

---

- **MICROVASCULAR DISEASE**

- Hyperglycemia damages small blood vessels:

- diabetic **retinopathy** → vision loss.

- diabetic **neuropathy** → damage to nerves → most common cause of amputation in Western world.

- diabetic **nephropathy** → kidney damage → chronic renal failure.





Oral glucose tolerance test

# *Glucose Tolerance Test*

---


- Both the FPG and OGTT tests require that the patient fast for at least 8 hours (ideally 12 hr) prior to the test.
- The oral glucose tolerance test (OGTT):
  - FPG test
  - Blood is then taken 2 hours after drinking a special glucose solution

# *Glucose Tolerance Test (GTT)*

---

- Following the oral administration of a standard dose of glucose, the plasma glucose concentration normally rises but returns to the fasting level within 2 hours.
- If insulin activity is reduced, the plasma glucose concentration takes longer than 2 hours to return to normal and often rises above 200 mg/dl.
- Measurement of urine glucose allows determination of the renal threshold for glucose.

# GTT



# *Glucose Tolerance Test*

The following results suggest different conditions:

- **Normal values:**
  - FPG < 100 mg/dl
  - 2hr PPG < 140 mg/dL
- **Impaired glucose tolerance**
  - 2hr PPG = 140 - 199 mg/dL
- **Impaired fasting glucose**
  - FPG = 100-125
- **Diabetes**
  - FPG  $\geq$  126 mg/dl
  - 2hr PPG levels  $\geq$  200 mg/dL

# Symptoms of Diabetes Mellitus

## *Symptoms of Diabetes Mellitus*

Hyperglycemia

Polyuria

Polydipsia

Polyphagia

Ketoacidosis (IDDM)








Hyperlipidemia

Muscle wasting

Electrolyte depletion



# Diabetes Mellitus (DM)

Organs/tissue involved	Organ/tissue responses to insulin deficiency	Resulting condition of:		Signs and symptoms
		Blood	Urine	
	Decreased glucose uptake and utilization	Hyperglycemia	Glycosuria	<b>Polyuria</b> - dehydration - soft eyeballs  <b>Polydipsia</b> Fatigue Weight loss <b>Polyphagia</b>
	Glycogenolysis		Osmotic diuresis	
	Protein catabolism and gluconeogenesis			
	Lipolysis and ketogenesis	Lipidemia and ketoacidosis	Ketonuria  Loss of Na <sup>+</sup> , K <sup>+</sup> ; electrolyte and acid-base imbalances	Acetone breath Hyperpnea Nausea/vomiting/abdominal pain Cardiac irregularities Central nervous system depression; coma
 = Muscle  = Adipose tissue  = Liver				









---

*The End*

*Thank You*