



LECTURE 1

HYPOTHALAMIC AND PITUITARY

GONADAL AXIS

Dr. Laila Al Dokhi
Assistant Professor
Department of Physiology






Objectives

By the end of this lecture, you should be able to:

1. Characterize hypothalamic pituitary relationship
2. Name the hypophysiotropic hormones and outline the effects that each has on anterior pituitary function
3. Name anterior pituitary gonadotropic hormones and outline the effects that each has on the gonads
4. Describe the negative and positive feedback mechanisms in the hypothalamic-pituitary-gonadal axis and their importance in the control of reproductive function




Keywords: hypophysiotropic hormones, gonadotropic hormones, androgens, estrogens

- What is a hormone?



Definition of hormone





Small group activity


- How hypothalamus controls anterior pituitary?
- What are the hormones secreted by anterior pituitary?
- How hypothalamus controls posterior pituitary?
- What are the hormones secreted by posterior pituitary?

Connection between Hypothalamus and anterior pituitary gland

Hypothalamic-hypophysial portal vessels


Figure 16.5b The hypothalamus controls release of hormones from the pituitary gland in two different ways (2 of 2).

Slide 1





The anterior pituitary secretes six hormones:

1. Adrenocorticotrophic hormone (corticotrophin, ACTH)
 2. Thyroid-stimulating hormone (thyrotropin, TSH)
 3. Growth hormone (GH)
 4. Follicle-stimulating hormone (FSH)
 5. Luteinizing hormone (LH)
 6. Prolactin (PRL)
- 


Control of male sexual functions by hypothalamic & anterior pituitary hormones :

GnRH :

- A peptide secreted by the arcuate nuclei of the hypothalamus
- Stimulate anterior pituitary gland to release gonadotropins (LH and FSH).
- GnRH is secreted intermittently for few minutes every 1 to 3 hrs.
- Secretion of LH by the anterior pituitary is also cyclical following the **pulsatile** release of **GnRH**.




Regulation of Testosterone production by LH:

- Testosterone is secreted by **leydig cells**, in the interstitium of the testis, by **LH** stimulation from the AP and
 - Its release is directly proportional to the amount of LH.
 - Mature leydig cells are found in a child's testis few weeks after birth & then disappear until puberty when it appear again.
- 


Negative feedback control of testosterone

- LH stimulate Testosterone secretion by the testis
- Testosterone inhibit the secretion of LH.
- Most of the inhibitory effect result from direct inhibition of GnRH release from the hypothalamus
- Inhibition of GnRH leads to decrease secretion of both LH & FSH.



Regulation of spermatogenesis by FSH and testosterone


- **FSH** binds with specific FSH receptors attached to the sertoli cell in the seminiferous tubules, which causes these cells to grow & secrete spermatogenic substances.
- Also testosterone & dihydrotestosterone diffuses into the seminiferous tubules from the leydig cells affect the spermatogenesis,
- So both FSH & testosterone are necessary to initiate spermatogenesis.



Negative feedback control of seminiferous tubule activity

Role of the hormone inhibin:

- When the seminiferous tubules fail to produce sperm the secretion of FSH from the AP increases.
- Conversely, when spermatogenesis proceeds rapidly pituitary secretion of FSH diminishes.
- This is due to the secretion of inhibin hormone from the sertoli cells which strongly inhibit the AP- FSH
- Inhibin has slight inhibitory effect on the hypothalamus to inhibit GnRH secretion.



Regulation of the female monthly rhythm


Interplay between the ovarian and hypothalamic-pituitary hormones:


Hypothalamic centers for release of GnRH:

- The neural activity that causes release of GnRH occurs in the mediobasal hypothalamus, in the arcuate nuclei which regulate most of the female sexual activity.
- Secretion of AP hormone is controlled by “releasing hormones”
- GnRH is secreted in pulses lasting 5-25 minutes every 1-2 hrs.
- The pulsatile release of GnRH cause intermittent output of LH secretion about every 90 minutes.

Follicular phase

- After menstruation the level of FSH & LH increases
- Mainly FSH accelerates growth of few follicles (6-12 follicles).
- The growing follicle secretes increasing amounts of estrogen





Negative feedback effects of estrogen and progesterone

- Estrogen in small amounts has strong effect to inhibit the production of LH & FSH.
- This inhibitory effect of estrogen is increased when progesterone is available.
- This inhibitory effects more on the AP directly & to lesser extent on the hypothalamus to inhibit the secretion of GnRH.

Hormone inhibin from the corpus luteum inhibits FSH & LH secretion:

- The hormone inhibin secreted by the granulosa cells of the ovarian corpus luteum inhibit the secretion of FSH & to lesser extent LH.

Positive feedback effect of estrogen before ovulation – the pre-ovulatory LH surge:

- AP secretes increased amount of LH for 1-2 days before ovulation.
- FSH surge is much smaller in the pre-ovulatory than LH surge.

The possible causes of LH secretion could be:

- Estrogen has special **positive feedback** effect of stimulating pituitary secretion of LH & to a lesser extent FSH
- The granulosa cells of the follicle begin to secrete small increasing amount of progesterone about 1 day before ovulation which stimulate LH secretion

Feedback oscillation of the hypothalamic-pituitary-ovarian system:

Postovulatory secretion of the ovarian hormones, and depression of the pituitary gonadotropins:

- During the postovulatory phase the corpus luteum secrete large quantities of both progesterone ,estrogen & inhibin
- Which all together cause negative feedback effect on AP & hypothalamus to inhibit both FSH & LH secretion.

(lowest level 3-4 days before the onset of menstruation)

Follicular growth phase :-

- 2 -3 days before menstruation, corpus luteum regress & secretion of estrogen, progesterone & inhibin decrease.
- This decrease remove the negative feedback effect on AP hormones.
- Therefore a day after menstruation FSH secretion begins to increase (2 folds) while LH secretion is low.
- These hormones causes growth of a new follicle.
- During the first 11 to 12 days of the follicular growth the rate of secretion of FSH & LH decrease due to the negative feedback effect of estrogen on the AP.

- 
- A vertical bar on the left side of the slide, consisting of several colored segments: a black segment at the top, followed by a white segment, a grey segment, a yellow segment, and a long pink segment at the bottom.
- https://youtu.be/2_owp8kNMus