
# CEREBRAL BLOOD CIRCULATION

#### **OBJECTIVES**

- At the end of the lecture, students should be able to:
- List the cerebral arteries.
- Describe the cerebral arterial supply regarding the origin, distribution and branches.
- Describe the arterial Circle of Willis.
- Describe the cerebral venous drainage and its termination.
- Describe arterial & venous vascular disorders and their clinical manifestations.

#### **CEREBRAL ARTERIAL SUPPLY**

- It is composed of two arterial systems:
  - A. Carotid System
  - B. Vertebro BasilarSystem



#### **CAROTID SYSTEM**

It is composed of:

Internal carotid artery


and its branches:

Anterior cerebral

artery&


Middle cerebral

artery




#### **VERTEBRO BASILAR SYSTEM**


- The two Vertebral
   arteries (from Subclavian
   artery) unite to form
   Basilar artery.
- It divides at the upper border of the pons into two Posterior Cerebral arteries.



Distribution of the cerebral arteries on the superolateral surface of the cerebral H





Distribution of the cerebral arteries on the medial surface of the cerebral H



#### **Anterior Cerebral Artery**


- Supplies: orbital and medial surfaces of the frontal and parietal lobes
- A narrow part on the superolateral surface.






#### **Middle Cerebral Artery**


- Supplies entire
   Superolateral surface:
  - SomatosensoryCortex
  - Motor Cortex
  - Language areas:
  - Broca's Area
  - Wernicke's Area)
  - Auditory areas:
  - Primary auditory area
  - (Heschl's Gyrus)
  - Auditory association




## **Posterior Cerebral Artery**

- Supplies:
- Anterior and inferior parts of temporal lobe, Uncus, Inferior temporal gyrus,
- Inferior and Medial parts of Occipital lobe (visual areas)









## Circulus Arteriosus (of Willis)

It joins the Carotid & Vertebrobasilar systems




- <u>located</u> on the base of the brain
- It encircles:
- Optic Chiasma, Hypothalamus Pituitary gland Midbrain.





#### Composed of:

- 2 Anterior cerebral arteries
- 2 Internal carotid arteries
- 2 Posterior cerebral arteries
- Posterior communicating arteries
- 1 Anterior communicating artery



- Branches:
- Perforating arteries (Anterior& Posterior):
- Numerous small vessels that penetrate the surface of the brain through the anterior and posterior perforating substances.
- APA supply:
- Large part of Basal Ganglia,
- Optic chiasma,
- Internal capsule & Hypothalamus
- PPA supply:
- Ventral portion of Midbrain, parts of Subthalamus and Hypothalamus






#### **Arterial Disorders**


- A. Stroke (Sudden occlusion
- of the blood supply):
- It can be:
  - 1. Hemorrhagic
  - 2. Ischemaic
  - **B. Aneurysm**
- **C. Angloma**

Hemorrhagic Stroke



Hemorrhage/blood leaks into brain tissue






Clot stops blood supply to an area of the brain




E



#### **EFFECT OF OCCLUSION of Cerebral arteries**




#### ACA



- 1. Motor & sensory disturbances in the contralateral distal leg
- 2. Difficulty in the Prefrontal lobe functions:
- Cognitive thinking, Judgment,
- Motor initiation and
- Self monitoring


#### MCA




- 1. Contralateral weakness of:
  - Face, Arm, Hand & leg
  - 2.Contralateral sensory loss of:
- Face, Arm & Hand &leg
  - 3. Visual field cut (damage to optic radiation)
- 4. Aphasia (language disturbances )
  - Broca's: production
  - Wernicke's: comprehension


#### PGA

- 1. Visual disturbances
  - Contralateral homonymous hemianopia
  - In Bilateral lesions:
     Cortical Blindness
    - patients unaware they cannot see
       (Anton's syndrome)
- 2. Memory impairment
- If the temporal lobe is affected





## **Cerebral Venous Drainage**




- Cortical Voins:
- (A) Superficial
- found in the
   Subarchnoid space
   Drain the cortical surfaces
- (B) Deep veins:
- Drain the deeper structures
- These veins are thin walled and devoid of valves.
- They ultimately drain into the
- Dural Venous Sinuses

#### **Superficial Cortical Veins**

## 1. <u>Superior</u> <u>cerebral veins</u> (6 to 12)

- Drain lateral surface of brain above the lateral sulcus
- Terminate mainly into the Superior Sagittal sinus, and partly into Superficial middle cerebral vein.




#### 2. Inferior cerebral veins:

- Run below the lateral sulcus
- Drain the lateral surface of the temporal lobe
- Terminate partly into superficial middle cerebral vein & partly into Transverse sinus.


#### 3.Superficial middle cerebral vein:

- Runs along the lateral sulcus
- Terminates into the Cavernous sinus
- It is connected posteriorly through Superior & Inferior anastomotic veins to Superior Sagittal & Transverse sinuses.



## **Deep Cerebral Veins**


- Drain the internal structures
   (basal ganglia, internal capsule, thalamus)
- They merge to form two Internal Cerebral Veins.
- The two veins unite in the midline to form the Great Cerebral vein.
- This short vessel joins the Inferior Sagittal sinus to form the Straight S



#### **Dural Venous Sinuses**

#### **Paired**


Transverse.
Sigmoid.
Cavernous.
Petrosal
(Sup & Inf)




#### **Single**

Superior sagittal. Inferior sagittal. Straight. Occipital.

Blood flows from transverse &sigmoid sinuses into IJV





#### **Venous Disorders**

- Infarcation.
- Sinus thrombosis:
- (SSS thrombosis) can complicates ear infection .
- Cavernous S thrombosis (as a complication of infection in the dangerous area of the face)
- Obstruction of venous drainage of the brain leads to Cerebral edema and raised ICP



## Thank You & Good Luck