


SENSORY
(ASCENDING)
SPINAL
TRACTS


*Dr. Jamila
El-Medany*

*Dr. Essam Eldin
Salama*



OBJECTIVES


- *By the end of the lecture, the student will be able to:*
- *Define the meaning of a tract.*
- *Distinguish between the different types of tracts.*
- *Locate the position of each tract.*
- *Describe the sensory pathway.*
- *Identify the different sensory spinal tracts and their functions.*
- *Identify the course of each of these tracts.*
- *Know some associated lesions regarding the main tracts.*



- ❑ The **grey matter** of the spinal cord is completely surrounded by the white matter
- ❑ The **White matter** of the spinal cord consists of **Ascending** and **Descending Nerve Fibers**.
- ❑ It is divided into **Dorsal, Lateral & Ventral Columns** or **Funiculi**.

WHITE MATTER TRACTS

- *Bundles or fasciculi of fibers that occupy more or less definite positions in the white matter.*
- *They have the same **Origin,** **Termination** and carry the same **Function.***




CLASSIFICATION OF WHITE MATTER TRACTS

- They are classified into:

1- Short Tracts;
intersegmental or propriospinal).

*Fibers occupy narrow band immediately peripheral to the grey matter (**fasciculus proprius**).*

*They interconnect adjacent or distant spinal segments
And Permit **intersegmental coordination***






• 2-Long Tracts:

(a) Ascending (sensory or afferent).

(b) Descending (motor or efferent).


They serve to join the brain to the spinal cord.





□ Ascending Tracts;


- Carry impulses from pain, thermal, tactile, muscle and joint receptors to the brain.
- *Some of this information eventually reaches a conscious level (at the cerebral cortex),*
- *while some is destined for subconscious centers (e.g at the cerebellum).*




- Pathways that carry information to a **conscious level** share certain common characteristics:
- There is a *sequence of **Three Neurons*** between the peripheral receptors and the cerebral cortex.

The axons of the **first-order neurone** (or primary afferent neurone) enters the spinal cord through the dorsal root of a spinal nerve and its cell body lies in the **dorsal root ganglion**.


The main fiber remains on the ipsilateral side of the cord and terminates in synaptic contact with the **second neurone which lies** either in the **spinal grey matter** or in the **medulla oblongata of the brain stem**.



- The axon of the **second order neurone crosses over (decussates)** to the opposite side of the CNS and ascends to the **thalamus**, where it terminates.
- **The third-order neurone** has its cell body in the **thalamus**.
- Its axon passes to the **somatosensory cortex** of the parietal lobe of the cerebral hemisphere.




- **Three major pathways carry sensory information:**
 - **Dorsal (Posterior) column (Gracile & Cuneate fasciculi)**
 - **Anterolateral pathway (Spinothalamic)**
 - **Spinocerebellar pathway**




Dorsal Column

- Composed of two tracts; ***Fasciculus Gracilis (FG)*** & ***Fasciculus Cuneatus (FC)***
- **They Carry** the axons of primary afferent neurons that have entered cord through dorsal roots of spinal nerves, from **ipsilateral side** of the body
- **They are concerned with:**
 1. **Discriminative Touch:**
 - Ability to localise accurately the area of the body touched.
 2. **Two Point Discrimination:**
 - To be aware that two points are touched simultaneously even they are close together.
 3. **Proprioception** from muscles and joints for the movement & position of different parts of the body ,




- **Fasciculus Gracilis** contains fibers that are received at sacral, lumbar and lower thoracic levels,
- **Fasciculus Cuneatus** contains fibers that are received at upper thoracic and cervical levels



Dorsal Column

- Fibers ascend without interruption where they terminate upon **2nd order neurons** in **Nucleus Gracilis** and **Nucleus Cuneatus**.
- The axons of the 2nd order neurons decussate in the medulla as **Internal Arcuate Fibers** and ascend through the brain stem as **Medial Lemniscus**.
- The medial lemniscus terminates in the **Ventral Posterior Nucleus of the Thalamus** (3rd order neurons), which project to the somatosensory cortex (**thalamocortical fibers**)




Tabes Dorsalis

A late manifestation of **syphilitic infection** on the CNS.

Affects the **lumbosacral dorsal spinal roots and dorsal columns of the spinal cord.**

Leads to **loss of proprioception** which is manifested by a **high Step Page** and **unsteady gait (Sensory Ataxia)**



Identifying gait abnormalities

SPASTIC GAIT



SCISSORS GAIT



PROPULSIVE GAIT



STEPPAGE GAIT



WADDLING GAIT



Subacute Combined Degeneration of the spinal cord


A systemic disease results from B12 deficiency

It produces Sensory Ataxia

Lateral columns are also affected (combined) causing **weak and spastic limbs**

It is completely recovered by proper treatment with B12

Subacute Combined Degeneration



Multiple Sclerosis

An immune disease affects specifically **fasciculus Cuneatus** of the cervical region. Leads to **loss of proprioception in hands and fingers (Asteriognosis)**

Main symptoms of Multiple sclerosis

Central:

- Fatigue
- Cognitive impairment
- Depression
- Unstable mood

Visual:

- Nystagmus
- Optic neuritis
- Diplopia

Speech:

- Dysarthria

Throat:

- Dysphagia

Musculoskeletal:

- Weakness
- Spasms
- Ataxia

Sensation:


- Pain
- Hypoesthesias
- Paraesthesias

Bowel:

- Incontinence
- Diarrhea or constipation


Urinary:

- Incontinence
- Frequency or retention



Spinothalamic Tracts

- The spinothalamic tracts contain axons of *second-order neurones*, the cell bodies of which lie in the **contralateral** dorsal horn.
- Located lateral and ventral to the ventral horn.
- Carry impulses concerned with; **pain and thermal sensations (Lateral tract)** and **Non-Discriminative touch and pressure (Anterior tract)**, from the contralateral side.
- In brain stem, the two tracts constitute the **Spinal Lemniscus**.
- Information is sent to the primary sensory cortex on the opposite side of the body.




❑ Function:

- Carries **Pain & Temperature** to thalamus and sensory area of the cerebral cortex.

❑ Neurones: 3 Neurones

- **Neurone I:** Small cells in the dorsal root ganglia.
- **Neurone II:** Cells of **Substantia Gelatinosa of Rolandi** in the posterior horn.
- **Neurone III:** Cells of **(VP) nucleus of the thalamus**.

Lateral Spinothalamic Tract



Anterior Spinothalamic Tract

❑ Function:


- Carries **Light**, **Crude Touch** (non discriminative) & **Pressure** to thalamus and sensory cortex.

❑ Neurones: 3 Neurones

- **Neurone I:**
Medium sized cells in the dorsal root ganglia.
- **Neurone II:**
Cells of main sensory nucleus or (**Nucleus Proprius**).

Fibers arising from **Substantia Gelatinosa & Nucleus Proprius** decussate in the **Anterior White Commissar**

- **Neurone III:**
Cells of VP nucleus of thalamus.



Spinothalamic Tracts Lesion


It is selectively damaged in **Syringomyelia**

The central canal becomes enlarged forming a cavity compressing the adjacent nerve fibres

Fibres serving pain and temperature are damaged first as they decussate in the ventral white commissure close to the central canal causing selective loss of pain and temperature in the upper limbs (**dissociate sensory loss**)


Light touch and proprioceptive sensations are retained

Joints of the limbs become disorganized without discomfort (**Charcot's joint**)




Spinocerebellar Tracts

- The Spinocerebellar system consists of a sequence of **only two neurons**;
- **Neurone I**: Large cells of dorsal root ganglia.
- **Neurone II**: cells of the nucleus dorsalis; **Clark's nucleus** (column) .
- **Two tracts: Dorsal & Ventral**
- Located near the dorsolateral and ventrolateral surfaces of the cord
- Contain axons of the second order neurons
- Carry unconscious information derived from **muscle spindles, Golgi tendon, tactile receptors joints, skin and subcutaneous tissue** to the cerebellum for the control of posture and coordination of movements




- Present only above level L3
- The cell bodies of 2nd order neuron lie in **Clark's column**
- Axons of 2nd order neuron terminate **ipsilaterally** (uncrossed) in the cerebellar cortex by entering through the **inferior cerebellar peduncle**.
- **Posterior spinocerebellar tract convey sensory information to the same side of the cerebellum**

Posterior Spinocerebellar Tract



Ventral (Anterior) Spinocerebellar Tract


- The cell bodies of 2nd order neuron lie in base of the dorsal horn of the lumbosacral segments
- Axons of 2nd order neuron **cross to opposite side**, ascend as far as the midbrain, and then make a sharp turn caudally (the fibers cross the midline for the **second time**) and enter the **superior cerebellar peduncle** to terminate in the cerebellar cortex
- So Ventral spinocerebellar tract conveys sensory information to the same side of the cerebellum



Lesion of the Spinocerebellar Tracts


Friedrichs ataxia

- An inherited degenerated disease
- Affecting the spinocerebellar tracts
- Leading to **incoordination of arms, intense tremor, wide base reeling gait ataxia**
- It begins in childhood
- Wheelchair is bound by 20 years of age



Spinotectal Tract

- Ascends in the **anterolateral** part, in close association with spinothalamic system.
- Primary afferents reach dorsal horn through **dorsal roots** and terminate on 2nd order neurons
- The cell bodies of 2nd order neuron lie in **base of the dorsal horn**.
- Axons of 2nd order neuron cross to opposite side, and project to the **periaquiductal gray matter** and **superior colliculus in the midbrain**.
- Involved in reflexive turning of the head and eyes (spino visual reflex) toward a point of cutaneous stimulation.



Spino - olivary Tract


Indirect spinocerebellar pathway (spino-olivo-cerebellar)

Impulses from the spinal cord are relayed to the cerebellum via **Inferior Olivary Nucleus.**

Conveys sensory information to the cerebellum.


Fibers arise at all levels of the spinal cord.

Contribute to movement coordination associated primarily with Balance.



Spinoreticular Tract

- Originates in the dorsal horn, and ascend in the ventrolateral region of the cord
- Contains uncrossed fibers that end in **medullary reticular formation** & both
- crossed & uncrossed fibers that terminate in **pontine reticular formation**, finally to the thalamus; to **Activate the cerebral cortex**
- Forms part of the ascending reticular activating system.
- **Involved in perception of dull aching (slow pain)**



Thank you