



Central Nervous System

Pathology Practicals

Prepared by:

- *Prof. Ammar Al Rikabi*
- *Dr. Sayed Al Esawy*
- *Dr. Abdullah Basabein*

Head of Pathology Department Dr. Hisham Al Khalidi

OBJECTIVES:

At the end of this practical sessions, the students will be able to:

- Recognize, describe and understand the morphological appearance (both macroscopic and microscopic) of some of the common diseases and disorders of the CNS.

• CONTENTS:

Study of the macroscopic and microscopic features through case studies of the following CNS diseases:

- Hydrocephalus.
- Meningitis.
- Brain abscess
- Brain hemorrhage.
- Alzheimer's disease.

REFERENCE: Robbin's Basic Pathology, 9th edition by Vinay Kumar, et al.



Hydrocephalus


CASE 1:

- *A 9 months infant was suffering from enlarged head size and admitted to hospital with convulsions, went into coma and died. Autopsy was done and the brain was large with dilated ventricles .*
- *What is your provisional diagnosis?*

Hydrocephalus




Hydrocephalus – Gross



This is hydrocephalus. Note the marked dilation of the cerebral ventricles. Hydrocephalus can be due to lack of absorption of CSF or due to an obstruction to flow of CSF.

Hydrocephalus vs Normal – MRI view



An MRI scan of a brain with hydrocephalus (left) and a normal MRI scan (right). The large dark area on the left is the ventricles, made bigger by a build-up of CSF

Hydrocephalus – MRI view



Mid Sagittal MRI of a child with communicating hydrocephalus, involving all ventricles.




Pyogenic (Bacterial) Meningitis

CASE 2 :

- *4 years old child who was treated from otitis media and suddenly complained from headache, vomiting, fever and stiff neck. CSF was found to be clouded with abnormal increase of neutrophils, increased protein and absence of sugar. Gram stain of the CSF fluid showed meningococci .*


What is your diagnosis ?

Bacterial Meningitis – Gross




Bacterial meningitis is the infection of the arachnoid membrane, subarachnoid space, and cerebrospinal fluid by bacteria. A creamy purulent exudate covers the cerebral hemispheres

Bacterial Meningitis – Gross




A creamy purulent exudate covers the cerebral hemispheres and settles along the base of the brain, around cranial nerves and the openings of the fourth ventricle

Acute Bacterial Meningitis – Gross




Here is another example of an acute meningitis from bacterial infection. The cerebrospinal fluid (CSF) in such cases typically has a low glucose, high protein, and many PMN's. A gram stain should be done to identify organisms.

Bacterial Meningitis – LPF Microscopy




A neutrophilic exudate is seen involving the meninges at the left, with prominent dilated vessels. There is edema and focal inflammation (extending down via the Virchow-Robin space) in the cortex to the right. This acute meningitis is typical for bacterial infection

Bacterial Meningitis – LPF Microscopy



Neutrophils in the subarachnoid space infiltrate and damage cranial nerves resulting in cranial nerve deficits, and invade leptomeningeal vessels causing phlebitis and arteritis with thrombosis and ischemic infarction

Bacterial Meningitis – CSF Gram stain



Microscopically, a gram stain of CSF sample reveals gram negative diplococci within a neutrophil, typical for Neisseria meningitidis



Cerebral Abscess

CASE 3:



- *A 35 years old lady complains from otitis media . Suddenly she suffers from headache and convulsions. Brain MRI reveals 5 cm. fluid filled cavity in the temporal lobe. Examination of the CSF shows increased pressure with lymphocytes and increased protein but there is no change of sugar content.*
- *What is your diagnosis ?*

Cerebral Abscess - Gross




This is a cerebral abscess. There is a liquefactive center with yellow pus surrounded by a thin wall associated with cavity formation and purulent/pus material. Abscesses usually result from hematogenous spread of bacterial infection, but may also occur from direct penetrating trauma or extension from adjacent infection in sinuses.

Cerebral Abscess - Gross




Cerebral Abscess – CT scan




This CT scan of the head in transverse view demonstrates an abscess in the brain consist of cavity with enhanced periphery (red arrow) in a patient who had septicemia. Predisposing conditions include: bacterial endocarditis, cyanotic congenital heart diseases and chronic pulmonary infections, e.g. bronchiectasis.

Cerebral Abscess – MRI scan




This MRI scan of the head in transverse (axial) view demonstrates a small abscess in the brain (Red arrow) in a patient who had septicemia

Cerebral Abscess – Microscopic view



This trichrome stain demonstrates the light blue connective tissue in the wall of an organizing cerebral abscess. Normal brain is at the right and the center of the abscess at the left.




***Ruptured Berry Aneurysm
causing subarachnoid
hemorrhage***

CASE 4:

- *A previously healthy 31-year-old woman experiences a severe headache and loses consciousness within an hour. An emergent head CT scan reveals extensive subarachnoid hemorrhage at the base of the brain. She is afebrile. A lumbar puncture yields cerebrospinal fluid with many red blood cells, but no white blood cells. The CSF protein is slightly increased, but the glucose is normal.*

What is your provisional diagnosis ?

Common locations of intracranial aneurysms




Saccular aneurysms most frequently form in first- and second-order arteries originating from the cerebral arterial circle (circle of Willis) at the base of the brain

Circle of Willis – Berry aneurysms




The circle of Willis has been dissected, and three berry aneurysms are seen. Multiple aneurysms are seen in about 20-30% of cases of berry aneurysm.

Circle of Willis: Ruptured Berry Aneurysm - Gross




Circle of Willis: Berry Aneurysm Ruptured- Gross natural color close-up view of base of brain showing subarachnoid hemorrhage over anterior surface of pons and a large aneurysm at top of photo which is located in the right internal carotid artery

Circle of Willis: Ruptured Berry Aneurysm - Gross



The subarachnoid hemorrhage from a ruptured aneurysm is more of an irritant producing vasospasm than a mass lesion.

Berry Aneurysm - LPF



Berry Aneurysm: Micro low mag H&E section of basilar artery and adjacent a portion of the aneurysm which was at the posterior inferior cerebellar artery good photo to show lack of medial structures in wall of aneurysm




Alzheimer disease


CASE 5:

- *A 85 years old man complains of progressive loss of memory, disorientation and alterations in mood and behavior since 20 years. He was admitted to hospital because he was disabled and immobile and he died in hospital after one week of admission. Autopsy was done and the brain cortex was found to be atrophied.*
- *What is your diagnosis ?*

Healthy Brain vs Alzheimer's Brain



Healthy Brain vs Alzheimer's Brain - Gross




Alzheimer disease: A. Normal Brain – B. The brain of a patient with Alzheimer shows cortical atrophy with thin gyri and prominent sulci

Alzheimer's Brain - Gross




The cerebral atrophy seen here mainly in the frontal and parietal regions is characterized by narrowed gyri and widened sulci. The atrophy seen here was due to senile dementia of the Alzheimer's type (Alzheimer's disease).

Alzheimer's disease - Illustration




Microscopic lesions of Alzheimer disease

Alzheimer's disease Neuritic plaques - LPF




The characteristic microscopic findings of Alzheimer's disease include "senile plaques" which are collections of degenerative presynaptic endings along with astrocytes and microglia. These plaques are best seen with a silver stain, as seen here in a case with many plaques of varying size.

*Alzheimer's disease
Neuritic plaques - LPF*





The plaques of Alzheimer's disease are seen here with a silver stain. Such neuritic (senile) plaques are most numerous in the cerebral cortex and hippocampus. This dementia is marked mainly by progressive memory loss.

*Alzheimer's disease
Neuritic plaques - LPF*



Alzheimer disease. A neuritic (senile) plaque with a rim of dystrophic neurites surrounding an amyloid core.

Alzheimer's disease Neurofibrillary tangles - HPF



Alzheimer Disease. Neurofibrillary tangles (arrows) are present within the neurons. They are composed of cytoskeletal intermediate filaments.

THE END