Prof. Ahmed Fathalla Ibrahim **Professor of Anatomy College of Medicine King Saud University** E-mail: ahmedfathala@gmail.com

BRAIN STEM

EXTERNAL FEATUR

OBJECTIVES

At the end of the lecture, students should:

- List the components of brain stem.
- Describe the site of brain stem.
- Describe the relations between components of brain stem & their relations to cerebellum.
- Describe the external features of both ventral & dorsal surfaces of brain stem.
- List cranial nerves emerging from brain stem.
- Describe the site of emergence of each cranial nerve.

DEVELOPMENT OF BRAIN




Figure 18 – 20. Diagrammatic sketches of the brain vesicles, indicating the adult derivatives of their walls and cavities. *The rostral part of the third ventricle forms from the cavity of the telencephalon; most of this ventricle is derived from the cavity of the diencephalon.


DEVELOPMENT OF BRAIN

- The brain develops from the cranial part of neural tube.
- The cranial part divides into 3 parts:
 - ***FOREBRAIN:** subdivides into:
 - <u>1-Two cerebral hemispheres (cavities: 2 lateral ventricles).</u>
 - 2-Diencephalon (cavity: 3rd ventricle) :
 - thalamus, hypothalamus, epithalamus & subthalamus
 - *MIDBRAIN (cavity: cerebral aqueduct).
 - *HINDBRAIN (cavity: 4th ventricle): subdivides into
 - 1-Pons.
 - 2-Cerebellum.
 - 3- Medulla oblongata.


BRAIN STEM

SITE:


- It lies on the basilar part of occipital bone (clivus).
- **PARTS:** From above downwards:
- Mid brain, pons & medulla oblongata
- CONNECTIONS WITH CEREBELLUM:
- Each part of brain stem is connected to cerebellum by cerebellar peduncles (superior, middle & inferior).








SAGITTAL SECTION OF BRAIN




IMPORTANCE OF BRAIN STEM

- 1. Pathway of tracts between cerebral cortex & spinal cord.
- 2. Site of origin of nuclei of cranial nerves (from 3rd to 12th).
- 3. Site of emergence of cranial nerves (from 3rd to 12th).
- 4. Contains groups of nuclei & related fibers known as **reticular formation** responsible for: control of level of consciousness, perception of pain, regulation of cardiovascular & respiratory systems.




BRAIN STEM – VENTRAL SURFACE




MEDULLA – VENTRAL SURFACE

- Ventral median fissure:
- It divides the medulla into 2 halves
- 2. Its lower part is masked by decussation of most of **pyramidal (corticospinal)** fibers (75%-90%).
- Pyramid: -
- 1. It lies on either side of ventral median fissure
- 2. It is an elevation produced by corticospinal tract.




MEDULLA – VENTRAL SURFACE

- Olive:
- 1. It lies lateral to the pyramid.
- 2. It is an elevation produced by inferior olivary nucleus (important in control of movement).
- Nerves emerging from Medulla (4 nerves):
- 1. Hypoglossal (12th): between pyramid & olive
- 2. Glossopharyngeal (9th), vagus (10th) & cranial part of accessory (11th): dorsolateral to olive (from above downwards)



PONS – VENTRAL SURFACE


- Basilar sulcus:
- 1. It divides the pons into 2 halves.
- 2. It is occupied by basilar artery.
- Transverse pontine (pontocerebellar) fibers:-
- 1. It originates from pontine nuclei.
- 2. It cross midline & pass through contralateral middle cerebellar peduncle to enter the opposite cerebellar hemisphere.



PONS – VENTRAL SURFACE


- Nerves emerging from Pons (4 nerves):
- 1. Trigeminal (5th): from the middle of ventrolateral aspect of pons, as 2 roots: a small medial motor root & a large lateral sensory root.
- 2. Abducent (6th): at junction between pons & pyramid.

3. Facial (7th) & vestibulocochlear (8th): at cerebellopontine angle (junction between medulla, pons & cerebellum). Both nerves emerge as 2 roots: from medial to lateral: motor root of 7th, sensory root of 7th, vestibular part of 8th & cochlear part of 8th




MID BRAIN – VENTRAL SURFACE

- It is formed of a large column of descending fibers (crus cerebri or basispedunculi), on either side.
- The 2 crura cerebri are separated by a depression (interpeduncular fossa).
- Nerve emerging from Midbrain (one):
- Occulomotor (3rd): from medial aspect of crus cerebri.




SAGITTAL SECTION OF BRAIN



Cavity: central canal.

- **Composed of:**
- 1. Dorsal median sulcus: divdes the closed medulla into 2 halves.
- 2. Fasciculus gracilis: on either side of dorsal median sulcus.
- 3. Gracile tubercle: an elevation produced at the upper part of fasciculus gracilis, marks the site of gracile nucleus.
- 4. Fasciculus cuneatus: on either side of fasciculus gracilis.
- 5. Cuneate tubercle: an elevation produced at the upper part of fasciculus cuneatus, marks the site of cuneate nucleus.


CLOSED MEDULLA



- **Cavity: 4**th ventricle
- On either side, an inverted V-shaped sulcus divides the area into 3 parts (from medial to lateral):
- Hypoglossal triangle:


 overlies hypoglossal
 nucleus.
- 2. Vagal triangle: overlies. dorsal vagal nucleus.
- Vestibular area: overlies ~ vestibular nuclei.

OPEN MEDULLA



PONS – DORSAL SURFACE


- Separated from the medulla by an imaginary line passing between the caudal margins of middle cerebellar peduncle.
- On either side, a sulcus divides the area into 2 parts (from medial to lateral):
- 1. Medial eminence: overlies abducent nucleus.
- 2. Vestibular area: overlies vestibular nuclei.



MID BRAIN – DORSAL SURFACE

- Marked by 4 elevations:
- Two superior colliculi:

 concerned with visual reflexes.
- Two inferior colliculi: forms part of auditory pathway.
- Nerve emerging from Midbrain (one):
- Trochlear (4th): just caudal to inferior colliculus (The only cranial nerve emerging from dorsal surface of brain stem).



SUMMARY

- The brain stem is composed (from above downwards) of: midbrain, pons & medulla oblongata which are continuous with each other, with diencephalon above & with spinal cord below.
- The brain stem is connected with cerebellum through cerebellar peduncles.
- The brain stem is the site of cranial nuclei, the pathway of important ascending & descending tracts & the site of emergence of cranial nerves (from 3rd to 12th).
- Cranial nerves (with the exception of 4th) emerge from ventral surface of brain stem.

QUESTION 1

- □Which one of the following cranial nerves emerges from ventral surface of midbrain?
- 1. Occulomotor (3rd).
- 2. Trochlear (4th).
- 3. Abducent (6th).
- 4. Facial (7th).

QUESTION 2

- Regarding the medulla oblongata, which one of the following is correct?
- **1.** The pyramid is lateral to olive.
- 2. The hypoglossal nerve is the most lateral nerve emerging from it.
- 3. The cuneate tubercle is lateral to gracile tubercle.
- 4. The cerebellum is connected to it by middle cerebellar peduncle.

QUESTION 3

- Which one of the following is the site of the inferior colliculus?
- 1. In the ventral surface of medulla, lateral to the olive.
- 2. In the dorsal surface of medulla, medial to the vagal triangle.
- 3. In the ventral surface of midbrain, lateral to the medial eminence.
- 4. In the dorsal surface of midbrain, above the trochlear nerve.

