

ULTRASOUND OF LIVER AND GALL STONE

(LECTURE 2)

Radiology



Objectives:

- ▶ Introduction to US.
- ▶ Indications of liver and gall bladder US.
- ▶ Normal anatomy and radiological appearance.
- ▶ Pathology of liver and gall bladder.
- ▶ Common pathological cases.

The image features two thick black L-shaped corner brackets. One is positioned in the top-left corner, and the other is in the bottom-right corner. They are oriented towards each other, framing the central text.

INTRODUCTION TO US

Definition:

- ▶ a diagnostic technique in which ULTRA=high-frequency sound waves penetrate the body, bounce around, and produce multiple echoes; these echo patterns can be viewed as an image on a computer screen.
- ▶ Frequency ranges used in medical Ultrasound imaging are 2 - 20 MHz

US machine



ROBES

MACHINE


B- MODE.




M- MODE.



DUPLEX



COLOR DOPPLER



Advantages of US

- ▶ **noninvasive**
- ▶ inexpensive.
- ▶ Easy and available.
- ▶ Safe and **non-ionizing**.

Disadvantages of US

- ▶ Inability to penetrate gas or bone.
- ▶ Operator dependant.
- ▶ Less sensitive in some situations.



Indications of liver and gall bladder US

- ▶ Right upper quadrant pain.
- ▶ Jaundice.
- ▶ High liver function test.
- ▶ Fever work up.
- ▶ Screening for metastasis.

Normal anatomy and radiological appearance



Cont.



Pathology of the liver:

- ▶ Size.
- ▶ Diffuse liver disease.
- ▶ Focal liver disease.
- ▶ Hepatic vascularity.
- ▶ Biliary system obstruction/pathology.


Size abnormality

- ▶ Normal liver size:



- Myeloproliferative disorder eg. Polycythaemia rubra vera.

Cont.




▶ +/- focal lesion.



Diffuse abnormality

- ▶ Diffuse increase parenchymal echogenicity
(whiter than normal)
- ▶ Diffuse fatty infiltration
- ▶ Other infiltrative:
 - Malignant
 - Infectious
 - Glycogen storage disease




Cont.

- ▶ Diffuse decrease in parenchymal echogenicity.

(darker than normal)

- ▶ Acute hepatitis.
- ▶ Other:
- ▶ Malignant infiltration.



Focal liver lesions

- ❑ **Benign tumor:**

- ▶ Hemangioma.

- ❑ **Malignant tumor:**

- ▶ Primary eg. Hepatocellular carcinoma.

- ▶ Secondary metastasis eg. Colon breast.

- ❑ **Infective:**

- ▶ Abscess


- ▶ hydated cyst.

- ❑ **Congenital:**

- ▶ Hepatic cyst.

Cont.


Liver abscess




hemangiomas




metastasis




HCC



Cont.



Hydatid cyst



Vascular abnormality



❑ Portal venous system:

- ▶ thrombosis.
- ▶ Portal hypertension.

❑ Hepatic venous system:

- ▶ Thrombosis
- ▶ (Budd Chiari syndrome).

Cont.



Hepatic vein thrombosis



PV thrombosis

Biliary abnormality

- ▶ Intra-hepatic biliary radicals.

Less than 3mm

- ▶ Extra-hepatic “CBD”

Less than 8mm

- ▶ Causes of dilatation & obstruction:

- o Intra-luminal:



- ✓ Stone & mass.

- o Mural:

- ✓ stricture (benign & malignant)

- o Extrinsic:

- ✓ Compression mass & Lymph node




Pathology of gall bladder

- ▶ Intra-luminal pathology.
- ▶ Mural pathology.

Intra-luminal pathology


▶ Gall stone:

Acoustic shadowing




▶ Polyps

No acoustic shadowing.



Cont.

- ▶ Intraluminal:
Mass lesion
+- invasion
Gall bladder carcinoma.



Mural pathology

❑ Mural thickening:

➤ Primary:

Cholecystitis.

➤ Secondary:


✓ Cardiac failure.


✓ Cirrhosis.

✓ ascites

✓ Hypoalbuminaemia

✓ Renal failure.





COMMON
PATHOLOGICAL
CASES

Case one

- ▶ Middle age women presented to ED with fever, RUQ pain


- ▶ On exam

She looks ill, febrile and on pain

Abdomen: RUQ tenderness

- ▶ Lab high LFTs & WBC.

Cont.




- ▶ Thickening of GB wall $>3\text{mm}$.
- ▶ Distended GB
- ▶ Pericholecystic fluid.
- ▶ Hyperemia.
- ▶ Gall stone
- ▶ Acute calcular cholecystitis.

Case two

- ▶ Middle age women presented to surgical out patient clinic with 2 years history of recurrent RUQ pain mild to moderate in severity radiated to the right shoulder aggravated by fatty meal.
- ▶ On exam:
obese lady well not distressed, febrile or jaundiced.
- ▶ Lab LFTs normal.

Cont.




- ▶ Multiple oval shaped echogenic structures seen within GB causing acoustic shadowing
- ▶ GB stones

Case three

- ▶ Middle age man presented to ER with severe RUQ pain and yellowish discoloration of skin and sclera.
- ▶ On exam:
he looks ill, jaundiced and on pain but not febrile
- ▶ Lab high LFTs.

Cont.



- ▶ Dilated intra-hepatic and extra-hepatic biliary system
- ▶ Echogenic structure seen within CBD
- ▶ CBD stone causing biliary obstruction.

Case four

- ▶ Old man recently discovered to have colonic cancer presented to primary health care clinic with vague upper abdominal pain


- ▶ On exam:

he was thin, ill not febrile or jaundiced.

Mild abdominal tenderness enlarged liver with irregular outline.

- ▶ Lab mildly elevated LFTs.

Cont.



- ▶ Multiple hypoechoic focal hepatic lesions
- ▶ Metastatic liver lesions.

Case five

▶ Middle age man known case of HCV+ for 10 years presented to GI out patient clinic with history of weight loss, indigestion and mild abdominal pain. No fever.



▶ On exam:

he was ill, slim ,mildly jaundice not febrile.

Abdomen: bulging flanks, dilated tortuous vessels around umbilicus. Mild diffuse abdominal tenderness.

▶ Lab high LFTs.

Cont.




- ▶ Shrunken liver with irregular outline.
- ▶ Heterogeneous appearance.
- ▶ Focal hypoechoic lesion.

- ▶ Cirrhotic liver with HCC.

Case six

- ▶ Young man known IV drug addict presented to ER with high fever, chills, upper abdominal pain and vomiting
- ▶ On exam:
- ▶ He looks very ill, febrile and on pain.
- ▶ Abdomen: RUQ tenderness.
- ▶ Lab high LFTs & WBC.

Cont.



- ▶ Focal hypoechoic liver lesion with ill defined outline.
- ▶ Liver abscess.



thank you

Radiology:
The Eye of
Medicine