

Obesity


Dr. Sumbul Fatma

Obesity

- A disorder of body weight regulatory systems
- Causes accumulation of excess body fat
 - >20% of normal body weight

Body Mass Index (BMI)

- BMI is an indirect measure of obesity
- Correlates height, weight and amount of body fat in an individual




Copyright © 2014 Wolters Kluwer Health | Lippincott Williams & Wilkins

	BMI	GRADE
UNDER WEIGHT	≤ 18.5	
NORMAL	18.5 – 24.9	
OVER WEIGHT	25.0 – 29.9	
OBESE	30.0 – 34.9	I
OBESE	35.0 – 39.9	II
HIGHLY OBESE	≥ 40	III

High BMI is associated with increased mortality risk

Obesity is associated with a high risk of:

- Diabetes mellitus
- Hypercholesterolemia
- High plasma triglycerides
- Hypertension
- Heart disease
- Cancer
- Gallstones, arthritis, gout
- Mortality



Copyright © 2008 Wolters Kluwer Health | Lippincott Williams & Wilkins


Anatomic differences in fat deposition

- Health risks depend on the pattern of fat deposition

Android, “apple-shaped,” or upper body obesity

excess body fat deposited in the central abdominal area

- Associated with risk of hypertension, insulin resistance, diabetes, dyslipidemia, and coronary heart disease




Anatomic differences in fat deposition

Gynoid, “pear-shaped,” or lower body obesity

Fat deposited around the hips or gluteal region.

- Associated risks are lower

Small amounts
of visceral fat



Pear-shaped =
lower body
obesity





Different Fat Depots in the Body

■ Subcutaneous Fat

- The fat stored just under the skin in the abdominal and gluteal-femoral region
- Constitutes 80-90% of the total fat in the body

■ Visceral Fat

- Composed of omental and mesenteric fat present in close association with digestive tract



Biochemical differences in fat deposits

Abdominal fat	Gluteal Fat
Smaller cells	Larger cells
More responsive to hormones (both visceral and subcutaneous)	Less responsive (subcutaneous)
Release substances via portal vein to the liver	Release substances to circulation with no effect on the liver

Adipocytes

- Triacylglycerols (fats) are deposited in adipocytes (fat cells) which can increase in size up to a limit.

Prolonged overnutrition stimulates



Pre-adipocytes in adipose tissue



Proliferation / differentiation into mature fat cells




Increases adipocyte number



Adipocytes

- Thus obesity is due to a combination of increased fat cell size (hypertrophy) and number (hyperplasia).
- **Fat cells, once gained, are never lost**
- Reduction in weight causes adipocytes to reduce in size but not in number

Modest weight gain or loss in a non-obese person mainly affects the size, but not the number of adipocytes.



When adipocytes reach their maximum size, further weight gain is achieved by recruitment and proliferation of new pre-adipocytes.



Weight reduction is difficult after cell proliferation has occurred because the fat cells must become smaller than their normal size.

Ectopic Fat

- Excessive calories that cannot be stored in adipose tissue “spill over” into other tissues such as muscle and liver
- It is called “**ectopic fat**” that is strongly associated with insulin resistance

Factors contributing to obesity

- **Genetic:** familial tendency
- **Environmental and behavioral**
 - **Sex:** women more susceptible
 - **Activity:** lack of physical activity
 - **Psychogenic:** emotional deprivation/depression
 - **Alcohol:** problem drinking
 - **Smoking:** cessation of smoking???
- **Drugs:** e.g. tricyclic derivatives

Causes of weight Gain



- Energy imbalance
 - calories consumed not equal to calories used
- Over a long period of time
- Due to a combination of several factors
 - Individual behaviors
 - Social interactions
 - Environmental factors
 - Genetics



-
- More in and less out = weight gain
 - More out and less in = weight loss
 - Hypothalamus
 - control center for hunger and satiety
 - Endocrine disorder
 - Hormonal imbalance

Hormonal control

- Appetite is influenced by
 - afferent neural signals, circulating hormones, and metabolites
- These signals cause the release of hypothalamic peptides and activate efferent neural signals
- Adipocytes also function as endocrine cells
- They release many regulatory molecules:
 - Leptin, adiponectin, resistin



Leptin

- A protein hormone produced by adipocytes that is required to keep the body weight under control
- Signals the brain about fat store level
- Regulates the amount of body fat by:
 - **Controlling appetite and energy expenditure**
- Leptin secretion:
 - Suppressed in starvation (depletion of fat stores)
 - Enhanced in well-fed state (expansion of fat stores)
- Leptin causes overweight mice to lose weight and maintain weight loss

Leptin Resistance

- Leptin increases metabolic rate and decreases appetite in humans
- Plasma leptin level in obese humans is usually normal for their fat mass
- Resistance to leptin has been found in obese humans
- The receptor for leptin in the hypothalamus is produced *db* gene
- Mutation in *db* gene causes leptin resistance in mice
- Leptin resistance may have some role in human obesity
 - Dieting decreases leptin levels
 - Reducing metabolism, stimulating appetite

Adiponectin

- A protein hormone exclusively and abundantly secreted from adipocytes
- Promotes the uptake and oxidation of fatty acids and glucose by muscle and liver
- Blocks the synthesis of fatty acids and gluconeogenesis by hepatocytes
- Net effect is to increase the sensitivity to insulin, and improve glucose tolerance

Adiponectin

- Adiponectin levels are inversely correlated with body fat percentage and parallels with the HDL level
- Low levels are seen in metabolic syndrome and diabetes mellitus

Other Hormones

Ghrelin: A peptide hormone secreted by stomach

- Stimulates appetite
- Secretion increases just before meals and drops after meals
- Increases food intake
- Decreases energy expenditure and fat catabolism
- Levels in dieters are *higher* after weight loss
- The body steps up ghrelin production in response to weight loss
- The higher the weight loss, the higher the ghrelin levels

Other Hormones

Cholecystokinin: Peptides released from the gut after a meal

satiety signals to the brain

Insulin: Promotes metabolism

Metabolic Changes in Obesity

- Adipocytes send signals that cause abnormal metabolic changes such as:
 - Dyslipidemia
 - Glucose intolerance
 - Insulin resistance

Benefits of Weight Loss in obesity


- Weight loss decreases risk factors for obesity
- Leading to:
 - Lower blood pressure
 - Decreased serum triacylglycerols
 - Lower blood glucose levels
 - Increase in HDL levels
 - Decreased mortality
 - Beneficial changes in BMR
 - Decreased energy requirement
- Slow weight loss is more stable

Treatment options

- Physical activity combined with healthy diet decreases level of obesity
- Reduces risk for heart disease and diabetes

Dieting

- Use of low-calorie diet
- Restriction of excessive energy intake



Drugs

Orlistat

A pancreatic and gastric lipase inhibitor
Decreases the breakdown of dietary fat

Lorcaserin

promotes satiety

Surgery

- Surgical procedures are designed to reduce food consumption in patients with BMI >40
- Used when other treatment options fail

Reference

- Lippincott's Illustrated Reviews of Biochemistry 5th Edition

Thank You!

