PANCREAS


Objectives

The student should be able to describe:

- 1.The <u>endocrine part</u> of the pancreas within the <u>exocrine part</u>.
- 2.The <u>histological features</u> of the cells of islet of Langerhans.
- 3.The <u>function</u> of different cells of islets of Langerhans.


PANCREAS


- Stroma: capsule, septa & reticular fibers.
- Parenchyma: Pancreas is a mixed gland:
 - Exocrine part (acini & ducts): produces digestive pancreatic enzymes.
 - Endocrine part (islets of Langerhans): produces hormones.



Pancreatic Acini:


- They are serous acini: secreting a thin fluid rich in digestive pancreatic enzymes.
- <u>Centroacinar cells</u>: Their nuclei appear in the center of the acini. They represent the beginning of the ducts.
 <u>No myoepithelial cells</u> around the acini.






Pancreatic Acinar Cells: Pyramidal in shape. Nuclei are basal, rounded and vesicular. Cytoplasm: Basal part basophilic (due to abundant rER). Apical part acidophilic (due to secretory granules).

Serous acinus ——— (exocrine)








Duct System:

- Centroacinar cells.
- Intercalated ducts (low cuboidal).
- Intralobular ducts (NOT prominent).
- Interlobular ducts.

Main pancreatic duct.









Islets of Langerhans:

- Pale-staining spherical collections of endocrine cells, scattered among the acini.
- Richly vascularized by fenestrated capillaries.
- Each islet is surrounded and supported by reticular fibers.
- 1 million islets in human pancreas.
- Most numerous in the tail of pancreas.



Cells of the Islets:

- 5 types of cells in each islet:
 1, β (B) cells: secrete insulin.
 2, α (A) cells: secrete glucagon.
 3, δ (D) cells: secrete somatostatin.
 - 4. G cells: secrete gastrin.
 - 5. PP cells: secrete pancreatic polypeptide.
- Cannot be differentiated from one another by routine stains.




Cells of the Islets:


β (B) cells:

- Constitute 70% of islet cells.
- Concentrated in islet center.
- Function: secrete insulin which ↓ blood sugar.

α (A) cells:

- Constitute 15-20%.
- Concentrated in islet periphery.
- Granules are much more numerous, more tightly packed, smaller, and denser than those of β cells.
- Function: secrete <u>glucagon</u> which ↑ blood sugar.





Cells of the Islets:

δ (D) cells:

- Constitute 5-10% of islet cells.
- Scattered throughout the islet.
- Granules are less dense than those of β and α cells.
- Function: secrete
 <u>somatostatin</u> which ↓ release
 of hormones from endocrine
 pancreas and enzymes from
 exocrine pancreas.



Cells of the Islets:

G cells:

- Constitute 1% of islet cells.
- Scattered throughout the islet.
- Function: secrete <u>gastrin</u> which ↑ production of HCI by parietal cells of the stomach.

PP cells:

- Constitute 1% of islet cells.
- Scattered throughout the islet.
- Function: secrete pancreatic polypeptide which ↓ exocrine secretions of pancreas.



