

ENDOCRINOLOGY


Posterior pituitary

Dr. Hana Alzamil

PHYSIOLOGY OF POSTERIOR PITUITARY GLAND

- Hypothalamic control
- Posterior pituitary hormones
 - ADH
 - Physiological functions
 - Control of secretion
 - Osmotic stimuli
 - Non-osmotic stimuli
 - Oxytocin
 - Physiological functions
 - Control of secretion

POSTERIOR PITUITARY GLAND




- ◉ Does not synthesize hormones
- ◉ Consists of axon terminals of hypothalamic neurons

HYPOTHALAMIC CONTROL OF PITUITARY SECRETIONS

- Secretions of the posterior pituitary are controlled by
 - Nervous signals from hypothalamus

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.






Herring Body

Pituicytes function

It forms physical and chemical barrier between nerve terminal and blood vessels

Amplify auto receptor negative feedback





ANTIDIURETIC HORMONE

(vasopressin)

SYNTHESIS OF ADH

- ⦿ It is synthesized as pre-prohormone and processed into a nonapeptide (nine amino acids).
- ⦿ ADH synthesized in the cell bodies of hypothalamic neurons (supraoptic nucleus)
- ⦿ ADH is stored in the neurohypophysis (posterior pituitary)


RECEPTORS OF ADH (VASOPRESSIN)

- There are 3 types of receptors for ADH:
 - V1A
 - V1B
 - V2
- V1A receptors mediate vasoconstriction
- V1A receptors also found in the liver (glycogenolysis)
- V1B receptors are unique to anterior pituitary and mediate increased ACTH secretion
- V2 receptors are located in the principle cells in distal convoluted tubule and collecting ducts in the kidneys

MECHANISM OF ACTION OF ADH:

- ◉ ADH binds to V2 receptors on the peritubular (serosal) surface of cells (principle cells) of the distal convoluted tubules and medullary collecting ducts.
- ◉ Via adenylate cyclase/cAMP induces production and insertion of **aquaporin2** into the luminal membrane and enhances permeability of cell to water.
- ◉ Increased membrane permeability to water permits back diffusion of solute-free water, resulting in increased urine osmolality (concentrates urine).

MECHANISM OF ACTION OF ADH



THE SINGLE MOST IMPORTANT FUNCTION
OF ADH IS TO CONSERVE BODY WATER BY
REDUCING URINE OUTPUT



SECRETION OF ADH

OSMOTIC STIMULI


- ◉ If plasma osmolality is directly increased by administration of solutes, such as sodium, cause ADH release.
- ◉ Conversely, substances that enter cells rapidly, such as urea, do not change osmotic equilibrium and thus do not stimulate ADH release.
- ◉ ADH secretion is very sensitive to changes in osmolality.
- ◉ Changes of 1-2% result in increased ADH secretion.

SECRETION OF ADH

NON-OSMOTIC STIMULI

- Hypovolemia is perceived by “pressure receptors” -- carotid and aortic baroreceptors, and stretch receptors in left atrium and pulmonary veins.
- Normally, pressure receptors tonically inhibit ADH release.
- Decrease in blood pressure induces ADH secretion by reducing input from pressure receptors.
- The reduced neural input to baroreceptors relieves the source of tonic inhibition on hypothalamic cells that secrete ADH.
- Sensitivity to baroreceptors is less than osmoreceptors- senses 15 to 25% change in volume

FUNCTION OF ADH (VASOPRESSIN)




- 1 High blood osmotic pressure stimulates hypothalamic osmoreceptors
- 2 Osmoreceptors activate the hypothalamic neurosecretory cells that synthesize and release ADH
- 5 Low blood osmotic pressure inhibits hypothalamic osmoreceptors
- 6 Inhibition of osmoreceptors reduces or stops ADH secretion

Osmoreceptors

- 3 Nerve impulses liberate ADH from axon terminals in the posterior pituitary gland into the bloodstream

ADH



- 4 Kidneys retain more water, which decreases urine output

Sudoriferous (sweat) glands decrease water loss by perspiration from the skin

Arterioles constrict, which increases blood pressure


REGULATION OF ADH

Dehydration

- ADH released

Overhydration


- ADH inhibited



**Plasma Osmolarity stimulates both
ADH release and thirst via
OSMORECEPTORS**

Effects on Blood Vessels

Normal




Angiotensin II

ADH




Increased
Blood
Pressure

Vasoconstriction



Vasodilation



OTHER STIMULI THAT AFFECT ADH SECRETION

- ◉ Stimuli that increase ADH secretion:
 - Pain
 - Nausea
 - Surgical stress
 - Emotional stress
- ◉ Stimuli that decrease ADH secretion:
 - Alcohol intake

TABLE SUMMARIZES THE MAJOR CHARACTERISTICS OF OSMORECEPTORS AND BARORECEPTORS

Receptors	Osmoreceptors	Baroreceptors
Location	Anterolateral hypothalamus	Carotid sinus, aortic arch, pulmonary veins, atria
Value Measured	Plasma osmolality	Circulating volume
ADH Release Stimulated By	Activation of receptor	Suppression of receptor
Change Required for Action	1% above 280 mosm/kg	15-25% decrease
Resulting Amount of ADH	Small (antidiuretic)	Large (vasoconstriction)
Override Other?	no	yes

CONTROL OF ADH RELEASE

⊙ **Osmotic pressure:**

- Osmoreceptor mediated
- \uparrow osmolality \rightarrow \uparrow ADH secretion
- \downarrow osmolality \rightarrow \downarrow ADH secretion

⊙ **Volume effects**

- Baroreceptor mediated (vagus nerve)
- \uparrow blood pressure \rightarrow \downarrow ADH secretion
- \downarrow blood pressure \rightarrow \uparrow ADH secretion

وَبَشِّرِ الصَّابِرِينَ



OXYTOCIN

FUNCTION OF OXYTOCIN

◉ **Breast-feeding**

- Contracts the myoepithelial cells of the alveoli


(classic neuroendocrine reflex)

◉ **Childbirth** (parturition)

- In late pregnancy, uterine smooth muscle (myometrium) becomes sensitive to oxytocin


(positive feedback)

BREAST FEEDING



CHILDBIRTH

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.





OTHER STIMULI THAT CONTROL RELEASE OF OXYTOCIN


- In humans, oxytocin is thought to be released during hugging, touching, and orgasm in both sexes.
- Release increased during stress
- Release inhibited by alcohol
- In males secretion increases at time of ejaculation (contraction of smooth muscle of vas deferens)



OXYTOCIN AND AUTISM

- Autistic group had significantly lower plasma oxytocin levels than in the non-autism group
- Elevated oxytocin was associated with higher scores on social and developmental measures for the non-autistic children

SUMMARY OF POSTERIOR PITUITARY HORMONES ACTIONS



CLINICAL APPLICATION

- What will happen if the pituitary stalk cut above the pituitary gland ?
 1. Secretion of hormones stop totally.
 2. Secretion of hormones will not be affected.
 3. Secretion of hormones decreases then return to normal level after few days.

