
# BREASTFEEDING

BY: RUBANA BAABBAD
CONSULTANT NEONATOLOGIST
INTERNATIONAL BOARD OF LACTATION CONSULTANT
EXAMINERS



## contents

- Introduction
- Breastfeeding definitions
- Breast milk contents
- Advantages of breastfeeding
- Preparation of the prospective mother
- Establishing and maintaining the milk supply
- Technique of breastfeeding
- Expression of breast milk
- Contraindications

قال الحق تبارك وتعالى: (وَالْوَالِدَاتُ يُرْضِفْنَ أَوْلَادَهُ نَّ حُولَيْنِ كَامِلَيْنِ لِمَنْ أَرَادَ أَنْ يُتِمَّ أَوْلَادَهُنَّ حُولَيْنِ كَامِلَيْنِ لِمَنْ أَرَادَ أَنْ يُتِمَّ الرَّضَاعَة) [البقرة: 233].

ان كلمة الرضاعة والفصال ومشتقاتهما قد تكررت في القرآن الكريم أربع عشرة مرة في سبع سور وثماني آيات كريمات

ت	اسم السورة	رقم الآية	الكلمة
.1	سورة البقرة	233	يرضعن، الرضاعة، فصالاً، تسترضعوا
.2	سورة النساء	23	أرضعنكم، الرضاعة
.3	سورة الحج	2	مرضعة، أرضعت
.4	سورة القصص	7.12	ارضعيه، المراضع
.5	سورة لقمان	14	وفصاله
.6	سورة الاحقاف	15	وفصاله
.7	سورة الطلاق	6	ارضعن، فسترضع

# introduction

- Breastfeeding is the normal feeding for infants during the first months of life which can 't be replicated .
- It contains over 200 known component.
- Breast milk bring both nutritive& non nutritive signals to the neonate.

# HUMAN MILK UNIQUE COMPOSITION

Exclusive human milk feeding for the first 6 mon. of life, with continued breastfeeding for 2 years of life is the normative standard for infant feeding because of Its unique nutritive composition and non nutritive bioactive factors that promote survival and healthy development.

### CONT.

- Human milk composition is **dynamic**, and varies within a feeding, diurnally, over lactation, and between mothers and populations.
- Influences on compositional differences include maternal and environmental factors and the expression and management of milk (e.g. Storage and pasteurization)

# Nutritional component of human milk

• Macronutrients: varies within mothers and across lactation but is remarkably conserved across populations despite variations in maternal nutritional status.

differs between preterm and term milk, with preterm milk tending to be higher in protein and fat .


- It contain fat ,carbohydrate, proteins
- menirals,

### Micronutrients


- many micronutrients vary in human milk depending on maternal diet and body stores .
- including vitamins A, B1, B2, B6, B12, D, and iodine.
- Regardless of maternal diet, Vitamin K is extremely low in human milk and thus, the American Academy of Pediatrics recommends an injection of this vitamin to avoid hemorrhagic disease of the newborn
- .Vitamin D also occurs in low quantity in human milk, particularly with low maternal exposure to sunshine.

- Bioactive factor including:
- living cells (macrophages and stem cells)
- anti-infectious
- anti-inflammatory agents
- growth factors
- Prebiotics
- Probiotics
- vitamins, hormons, enzymes complements,
- lysozymes, immunoglobulins
- cytokines
- mucins

# Breast milk under microscopy



# Comparison of microscopic picture



# Colostrum vs. breast milk



# colostrum

 Colostrum, the yellowish, sticky breast milk produced at the end of pregnancy, is recommended by WHO as the perfect food for the newborn, and feeding should be initiated within the first hour after birth.

# colostrum

- Colostrum
  - The often yellow and sometimes clear fluid that is released by a new mother's breasts before her breast milk comes in.
- This fluid has often been referred to as "liquid gold" and it resembles blood more than it does milk as it contains protective white blood cells capable of attacking harmful bacteria. and it also acts to "seal" the inside of the baby's intestines thus preventing the invasion of bacteria. Colostrum is an ideal first food for baby as it is high in protein and low in sugar and fat, thus making it easy to digest.

### colostrum

- Is the first fluid produced after delivery up to 3-5 days and some times longer in premature delivery and with maternal obesity.
   The colostrum
  - ----primary functions is immunological and trophic

## Basic nutritional info on Breast milk

		Colostrum	Breast milk	Cows milk
	Calories	58	70	65
•	Protein	3.7gm	1.3gm	3.4gm
•	Carbohy	5.3gm	7.4gm	4.8gm
•	Fats	2.9gms	4.2gms	3.7gm

- Colostrum-> Thin , yellow , Low on fat & carbs.
- Breast milk -> White, thin, watery & sweet.
- Foremilk-> Watery, low fat & high carbs.
- Hindmilk-> Creamier, thick, high fat

# Colostrum in comparison to transitional and mature milk

### Higher in

- Rich in **immunological** components(s. IgA, lactoferrin,leukocytes, developmental factors such as epidermal growth factors.
- Na
- **Cl**
- Mg


#### Lower in

- Low volume
- lactose
- Potassium
- calcium

# BIOACTIVE COMPONENTS AND THEIR SOURCES

- **defined as** elements that "affect biological processes or substrates and hence have an impact on body function or condition and ultimately health".
- Bioactive components in human milk come from a variety of sources;
- 1. produced and secreted by the mammary epithelium,
- 2. produced by cells carried within the milk,
- 3. drawn from maternal serum and carried across the mammary epithelium by receptor-mediated transport.

# Sources of mother milk immune factors



## **BIOACTIVE FACTORS**

recognition of potent, bioactive human milk factors indicates the importance of preserving their biologic activity, to the extent possible, through the process of milk collection, storage, and pasteurization.

### IMMUNOLOGICAL FACTORS

- Transfer of living protection and programming:
   Cells of human milk
- Human milk contains a variety of cells, including macrophages, T cells, stem cells, and lymphocytes.
- Communication between cells: Cytokines and chemokines
- Cytokines are multi-functional peptides that act in autocrine/paracrine fashion.
- Chemokines are a special class of chemotactic cytokines that induce movement of other cells.

### Cont.

- Finally, recognition of the unique mechanisms by which human milk protects and enhances development provides models for new preventive and therapeutic approaches in medicine.
- Many of these factors act synergistically, such that consumption of human milk is superior to supplementation with individual factors or their combinations.

# **Breastfeeding Definitions**

• Exclusive breastfeeding (Breast milk (including milk expressed or from a wet nurse)

- Predominant breast milk: Breast milk (including milk expressed or from a wet nurse) as the predominant source of nourishment
- Complementary feeding; Breast milk (including milk expressed or from a wet nurse) and solid or semi-solid foods

May include anything else: any food or liquid including nonhuman milk and formula

# Cont.

- Breastfeeding: Breast milk (including milk expressed or from a wet nurse) and solid or semi-solid foods, may include anything else: any food or liquid including non-human milk and formula
- Bottle-feeding: Any liquid (including breast milk) or semi-solid food from a bottle with nipple/teat may include anything else: any food or liquid including non-human milk and formula

# WHO recommendation

- Breastfeeding is the normal way of providing young infants with the nutrients they need for healthy growth and development. Virtually all mothers can breastfeed, provided they have accurate information, and the support of their family, the health care system and society at large.
- Exclusive breastfeeding is recommended up to 6 months of age, with continued breastfeeding along with appropriate complementary foods up to two years of age or beyond.

# The Breastfed Baby

#### Immune system.

Responds better to vaccinations. Human milk helps to mature immune system. Decreased risk of childhood cancer.

Throat.

Children who are

likely to require

tonsillectomies.

breastfed are less

#### Skin.

Less allergic eczema in breastfed infants.

#### Joints and muscles.

Juvenile
rheumatoid
arthritis is
less common
in children
who were
breastfed.

#### Bowels.

Less constipation.

#### Urinary tract.

Fewer infections in breastfed infants

#### Appendix.

Children with acute appendicitis are less likely to have been breastfed.

Eyes.

Ears.

Visual acuity is

human milk.

Breastfed babies

get fewer ear

infections.

higher in babies fed

Kidneys.
With less salt
and less protein,
human milk is
easier on a baby's
kidneys.

#### Digestive system.

Less diarrhea, fewer gastrointestinal infections in babies who are breastfeeding. Six months or more of exclusive breastfeeding reduces risk of food allergies. Also, less risk of Crohn's disease and ulcerative colitis in adulthood.

#### Higher IQ.

Cholesterol and other types of fat in human milk support the growth of nerve tissue.

### Endocrine system.

Reduced risk of getting diabetes.

#### Mouth.

Less need for orthodontics in children breastfed more than a year. Improved muscle development of face from suckling at the breast. Subtle changes in the taste of human milk prepare babies to accept a variety of solid foods.

#### Respiratory system.

Breastfed babies have fewer and less severe upper respiratory infections, less wheezing, less pneumonia and less influenza.

#### Heart and circulatory system.

Breastfed children have lower cholesterol as adults. Heart rates are lower in breastfed infants.

#### thealphaparent.com

# Benefits of breastfeeding for mothers

- Helps the uterus to regress to its size before pregnancy.
- Losing accumulated fat during pregnancy.
- Empowerment
- Decrease risk of osteoporosis
- Improve blood sugar control for diabetics
- Decrease the incidence of high cholesterol, diabetes, breast, cervical, ovarian, uterine cancers, less chance for gallstone formation and rheumatoid arthritis
- Bonding ,decrease post partum depression

# Advantages of breastfeeding

### Superior health outcomes in breastfed infant

Protection during breastfeeding	Protection after weaning in early childhood	Protection later in childhood
Gastrointestinal and respiratory infections	Gastrointestinal and respiratory infections	Obesity
Urinary infections	Wheezing	Types I and II diabetes
Sepsis and meningitis	Celiac disease	Leukemia/lyphomas
Atopic dermatitis	Growth faltering	Crohn disease
Food allergies	Cognition	Cognition
Wheezing	Visual acuity	Strong, secured personality
Necrotizing enterocolitis		
Celiac diseas		
Growth faltering		
Visual acuity		

## Preparation of the prospective mother

- Most women are physically capable of breastfeeding, provided the receive sufficient encouragement and are protected from discouraging experiences and comments while the secretion of breast milk is becoming established.
- Physical Factors: leading to a good breastfeeding include: good health, having enough rest, freedom of worry, treatment of any disease, and adequate nutrition.
- Retracted & inverted nipples.

A retracted nipple is a nipple that turns inward instead of outward, except when stimulated. This type of nipple is sometimes referred to as an inverted nipple. ... Some experts make a distinction between retracted and inverted nipples, referring to a retracted nipple as one that lies flat against the breast, rather than indenting in.

#### Hoffman technique

The Hoffman technique for drawing out inverted nipples has been in use since the 1950s. To try it:

- •Place your thumbs on either side of your nipple. Be sure to place them at the base of the nipple, not the outside of the areola.
- •Press firmly into your breast tissue.
- •While still pressing down, gently pull your thumbs away from each other.
- •Move your thumbs all around the nipple and repeat.


# Establishing and maintaining the milk supply

- The most satisfactory stimulus to the secretion of human milk is regular and complete emptying of the breast; milk production is reduced when the secreted milk is not drained.
- The breastfeeding should begin as soon after delivery as the condition of the mother and the baby permits, preferably within the first hour.

# Breastfeeding nosition



# Manual expression of breast milk







Expression of breastmilk

## Contraindications

- It is important to look at the entities that put the mother or infant at significant risk and are not remedial.
- Infectious Diseases:
- Life threatening illnesses in the mother:
- Medications:

# **Correct latching**



