

# Hormones affecting female breast



DR. MOHAMMED ALOTAIBI


# Objectives



**By the end of this lecture, you should be able to:**

- Know the physiologic anatomy of the breast.
- Describe the physiological changes that occur in the breast during mammogenesis, lactogenesis, and galactopoiesis and the hormones involved.
- Recognize the phases of lactogenesis.
- Describe the endocrine and autocrine control of lactation.
- Explain the physiological basis of suckling reflex and its role in lactation.


# The structure of the breast and mammary glands



## Each breast consists of 15-20


### lobes of secretory tissue:

- a. Each lobe has one lactiferous duct
- b. Lobes (and ducts) are arranged radially
- c. Lobes are composed of lobules
- d. Lobules are composed of alveoli




# Where does milk come from?

The fundamental secretory unit of the breast is the **alveolus**



# Lobule-Alveolar System



The function of the alveolar epithelial cells is to remove nutrients from the blood and transform these nutrients into the components of milk.

# Stages of Mammary Gland Development



- 1) **Mammogenesis** (*growth and development of mammary gland to a functional state*)
- 2) **Lactogenesis** (*initiation of milk secretion*)  
  
**Lactogenesis 1**  
**Lactogenesis 2**
- 3) **Galactopoiesis** (*maintenance of milk secretion in the postpartum period*)
- 4) **Involution** (*cessation of milk production*)

# Hormones affecting female breast



**Mammogenic Hormones** (promoting the proliferation of ductal and alveolar cells) [ductal & alveolar morphogenesis]

**Estrogens**

**Progesterone**



**During puberty** (ovarian hormones stimulate mammary growth)

**Prolactin (PRL)**

**Growth hormone**

**Glucocorticoid (Cortisol)**




# Ovarian Hormones

## ○ Estrogen

- ✦ Growth & branching of **ductal system** (with GH)
- ✦ Fat deposition in the stroma.

## ○ Progesterone

- ✦ Growth of **lobule-alveolar system** (budding of alveoli and secretory changes in epithelial cells).



Although progesterone and estrogen are essential for physical development of the breasts, they inhibit the actual secretion of milk during pregnancy by interacting with prolactin receptors.

# Lactogenesis


- **Lactogenesis:** Cellular changes by which **alveolar epithelial cells** switch from a **non-secretory** to a **secretory tissues** (initiation of milk secretion).

- **Involves 2 Phases:**

- **Lactogenesis 1**

- **Lactogenesis 2**

❖ **Lactogenic hormones** (promoting initiation /onset of milk production by alveolar cells)



# Lactogenesis 1



(Histological and enzymatic differentiations of alveolar epithelial cells).

- Starts in **mid-pregnancy** and characterized by expression of many (but not all) of genes involved in the synthesis of milk components (increases in uptake transport systems for amino acids, glucose, and calcium required for milk synthesis).
- Hormones involved:
  - Progesterone
  - Prolactin (PRL)
  - hPL (hCS)
  - Growth hormone
  - Glucocorticoid (Cortisol)

} **Alveologenesi**s

Production and secretion of milk components in this stage are restricted to a limited number of alveolar epithelial cells with incompletely developed secretory mechanisms.

# Lactogenesis 2



(Copious secretion of all milk components), **starts 2-3 days postpartum**

- At parturition, withdrawal of **progesterone** + high level of **prolactin** leads to:
  - ✦ Further increase in expression of milk protein genes
  - ✦ Glands absorb large quantities of metabolic substrates from the blood
  - ✦ Movement of cytoplasmic lipid droplets and casein into alveolar lumen
  - ✦ Transfer of immunoglobulins (IgA)
  - ✦ Secretion of colostrum followed by milk
  - ✦ Suckling stimulates further increase in expression of most of genes involved in milk secretion with expansion of alveolar epithelium
  - ✦ Switch from endocrine to autocrine control of milk production.

# Hormones affecting female breast



## Lactogenesis 2 hormones

**Prolactin (PRL)**

**Growth hormone**

**Glucocorticoid (Cortisol)**

**Thyroxine**


**Insulin**

**All are required to facilitate the mobilization of nutrients and minerals**

# Prolactin (PRL)



- Secreted from the anterior pituitary gland (Lactotrophs).
- Its level rises **steadily** from the 5th week of pregnancy until birth (10-20 times the nonpregnant level) (enhanced by Estrogen)



# Galactopoeisis



- Galactopoeisis is defined as the maintenance of lactation once lactation has been established. **starts 9-15 days postpartum**

**Galactopoietic Hormones:** maintaining milk production after it has been established

**PRL** (primary)

**Cortisol, Insulin, Thyroid and Growth hormones** (permissive)

**Oxytocin**

**Estrogen** (very low dose is galactopoietic)

**Progesterone** (alone has no effect on galactopoeisis because it has


no receptors in mammary glands at this stage)

# The alveolar cell secretes the components of milk through five pathways

D SECRETORY EPITHELIAL CELL

PRL stimulates transcription of the genes that encode several milk proteins.

Alveolar lumen




Transcellular salt and water transport through channels and transporters.

Paracellular pathway for ions and water.



# Oxytocin and psychogenic stimuli initiate milk ejection (“let-down”)




**Galactokinetic Hormones:** (promoting contraction of myoepithelial cells, and thus milk ejection).

- Oxytocin (OT)

- **Milk Ejection Reflex:**

*Oxytocin* contracts the myoepithelial cells, forcing milk from the alveoli into the ducts and sinuses where it is removed by the infant (**galactokinetic effect**).

# After delivery of placenta (withdrawal of high P conc.)



**Alveolus of Mammary Gland**

# Autocrine Control of Lactation




## Influence of Local Factors Acting on the Breasts

- It is not just the level of maternal hormones, but the efficiency of *milk removal* that governs the volume product in each breast
- A protein factor called *feedback inhibitor of lactation (FIL)* is secreted with other milk components into the alveolar lumen
- *FIL*, insensitive to prolactin → **⬇️ milk production**

# Autocrine Control of Lactation



## INHIBITOR IN BREASTMILK




- If breast remains full of milk, secretion stops

Control of breastmilk production within the breast.

# Suckling and Prolactin Secretion

Suckling is the most powerful physiological stimulus for PRL release




## Psychogenic stimuli

# Suckling reflex



**Prolactin**

suckling stimulates the release of **TRH** from the hypothalamus



# Milk production



- Milk production is a "use it or lose it" process. The more often and effectively the baby nurses, more milk will be produced.
- Lactation is maintained by continuous removal of milk.
- Milk production <100 ml/day in day 1 postpartum.
- Milk production by day 3 reaches 500 ml/day.
- **Involution:** when the breasts stop producing milk completely after weaning.

# AAP Recommendations

- Exclusive breastfeeding for the first **six** months of life
- Continued breastfeeding for at least one year, 'As long as is desired by mother and child'



يقول تعالى (والوالدات يرضعن أولادهن حولين كاملين لمن أراد أن يتم الرضاعة) [البقرة: 233]



**The End**

**Thank You**