



Editing File

Mnemonic File

Reproduction Block

Pharmacology team 438

Drugs Used in Treatment of Gonorrhoea and Syphilis

Objectives:

By the end of the lecture , you should know:

- ◆ List the drugs used in the treatment of syphilis & gonorrhoea
- ◆ Describe the mechanism of action and adverse effects of each drug
- ◆ Describe the contraindications of drugs used
- ◆ Describe the recommended regimens used for the treatment of syphilis & gonorrhoea
- ◆ Know the alternative treatments in allergic patients

Color index:

Black : Main content

Red : Important


Blue: Males' slides only

Purple: Females' slides only

Grey: Extra info or explanation

Green : Dr. notes

Syphilis

- Is a sexually transmitted disease caused by bacterium *Treponema Pallidum* (a spiral -shaped, Gram-negative highly mobile bacterium).
- *T. pallidum* enters the body via skin and mucous membranes through abrasions during sexual contact or blood transfusion or placental transfer from a pregnant woman to her fetus.
 - **Congenital Syphilis:** If a woman is pregnant and has symptomatic or asymptomatic early syphilis, disseminating organisms may pass through the placenta to infect the fetus.
 - Manifestation: Perforation of Palate. → 
- May become chronic without treatment.

Stages of Syphilis

Disease progresses in stages (primary, secondary, latent and tertiary), signs and symptoms vary depending upon stage of disease.

01



Primary stage :

- Painless skin ulceration (a chancre).



02



Secondary stage :

- Diffuse skin rash & mucous membranes lesions.



03



Latent stage :

- In latent syphilis there are little to no symptoms which can last for years.
- 70% may have no symptoms


04



Tertiary stage :

- Approximately 30% of untreated patients progress to the tertiary stage within 1 to 20 years.
- Rare because of the widespread use of antibiotics.
- Manifestations as cardiovascular syphilis (syphilitic aortitis and an aortic aneurysm).

Drugs used in the treatment of Syphilis ¹



β-Lactam Antibiotics:

1) Natural Penicillins

Drug	Benzylpenicillin (penicillin G)	Procaine penicillin G	Benzathine penicillin G
MOA	<ul style="list-style-type: none"> Inhibits bacterial cell wall synthesis through inhibition of transpeptidase enzyme required for crosslinks. Bactericidal. 		
P.K	<ul style="list-style-type: none"> Given I.V Short duration of action 	<ul style="list-style-type: none"> Given I.M. delayed absorption. Long acting (24 hrs) 	<ul style="list-style-type: none"> Given I.M. Delayed absorption. Long acting (days), 2.4 million units is given once.
ADR	<p>All these penicillin preparations are:</p> <ul style="list-style-type: none"> Acid unstable.² Penicillinase (β-lactamase) sensitive Not metabolized. Excreted unchanged in urine through acid tubular secretion. Renal failure prolong duration of action. 		
ADR	<ul style="list-style-type: none"> Hypersensitivity. Convulsions with high doses or in renal failure. Super infections. 		

2) 3rd generation cephalosporins

Drug	Ceftriaxone
MOA	<ul style="list-style-type: none"> Inhibits bacterial cell wall synthesis. Bactericidal.
P.K	<ul style="list-style-type: none"> Given parenterally (i.v.) Eliminated via biliary excretion Long Half-life
ADR	<ul style="list-style-type: none"> Hypersensitivity reactions Thrombophlebitis Superinfection Diarrhea

1) Not allergic to penicillins? Prescribe Penicillin. Allergic and not pregnant? Tetracycline. Allergic and pregnant? Cephalosporins or Macrolides.

2) Thus never given orally.

Tetracyclines

Drug	Doxycycline
MOA	<ul style="list-style-type: none"> Inhibit bacterial protein synthesis by reversibly binding to 30S bacterial ribosomal subunits. Bacteriostatic.
P.K	<ul style="list-style-type: none"> Given orally. Well absorbed orally. Long acting. 100 mg twice daily for 14 days.
ADR	<ul style="list-style-type: none"> Nausea, vomiting, diarrhea & epigastric pain (given with food). ★ Brown discoloration of teeth in children. ★ Deformity or growth inhibition of bones in children. Avoid dairy products. Hepatic toxicity in prolonged therapy with high dose. Vertigo. Superinfections. Phototoxicity.
C.I	<ul style="list-style-type: none"> Pregnancy. Breast feeding. Children (below 10 yrs).

Macrolides

Drug	Azithromycin
M.O.A	<ul style="list-style-type: none"> Inhibits bacterial protein synthesis by binding to bacterial 50S ribosomal subunits. Bacteriostatic
P.K	<ul style="list-style-type: none"> Acid stable → orally once daily Penetrates into most tissues except CSF. T_{1/2} 2-4 days . Should be given 1 hour before or 2 hours after meals. ★ No effect on cytochrome P450.
ADR	<ul style="list-style-type: none"> GIT upset: Nausea, vomiting, abdominal pain and diarrhea. Allergic reactions: urticaria and mild skin rashes.

WHO Guideline for Treatment of Syphilis

Prof Hanan: Doses aren't important, just know the drugs and in late stage the duration will increase

Early syphilis

Adults

(primary, secondary and early latent syphilis of not more than two years duration)

- **benzathine penicillin G** 2.4 million units once I.M.
- **procaine penicillin G** 1.2 million units I.M. for 10–14 days .

If penicillin is not allowed due to allergy, use :

- **Doxycycline** 100 mg twice daily orally for 14 days
- **Ceftriaxone** 1 g IM once daily for 10–14 days
- **Azithromycin** 2 g once orally.

Pregnant woman

- **benzathine penicillin G** . 2.4 million units once I.M.
- **procaine penicillin G** . 1.2 million units I.M. for 10–14 days

If penicillin is not allowed due to allergy, use :

- **Erythromycin** 500 mg orally 4 times daily for 14 days
- **Ceftriaxone** 1 g I.M once daily for 10–14 days
- **Azithromycin** 2 g once orally.

Late syphilis

Adults

(infection of more than two years duration without evidence of treponemal infection)

- **benzathine penicillin G** . 2.4 million units nits I.M. once weekly for three consecutive weeks.
- **procaine penicillin G** . 1.2 million units I.M. for 20 days .

If penicillin is not allowed due to allergy, use :

- **Doxycycline** 100 mg twice daily orally for 30 days

Pregnant woman

- **benzathine penicillin G** . 2.4 million units nits I.M. once weekly for three consecutive weeks.
- **procaine penicillin G** . 1.2 million units I.M. for 20 days

If penicillin is not allowed due to allergy, use :

- **Erythromycin** 500 mg orally 4 times daily for 30 days
- **Ceftriaxone** 1 g IM once daily for 10–14 days
- **Azithromycin** 2 g once orally.

Congenital Syphilis

In infants with confirmed congenital syphilis or infants who are or clinically normal, but whose mothers had untreated syphilis

- **Aqueous benzylpenicillin** (I.V.) 100 000–150 000 U/kg/day for 10–15 days
- **Procaine penicillin** (I.M.) 50 000 U/kg/day single dose for 10–15 days

Gonorrhoea

- Caused by **Neisseria gonorrhoea** [a gram -ve cocci], a pus producing bacteria .
- Transmitted during sexual contact with affected person.
- Many people have no symptoms
- Men may have burning with urination, discharge from the penis or testicular pain.
- Women may have burning with urination, vaginal discharge, vaginal bleeding between periods or pelvic pain.

Treatment of uncomplicated Gonorrhoea

Drugs	3rd generation cephalosporins E.g Ceftriaxone Cefixime	Fluoroquinolones E.g Ciprofloxacin
MOA	-	<ul style="list-style-type: none"> • Inhibit DNA synthesis by inhibiting DNA gyrase enzyme (required for DNA supercoiling) • Bactericidal
P.K	<ul style="list-style-type: none"> • Ceftriaxone (500 mg I.M) • Cefixime (400 mg orally) • Typically given in combination with a single dose of: <ul style="list-style-type: none"> ○ azithromycin(1gm orally) or doxycycline(100 mg orally twice daily) for 7 days 	<ul style="list-style-type: none"> • Single oral dose of : <ul style="list-style-type: none"> ○ Ciprofloxacin (500 mg) ○ Ofloxacin (400 mg)
Uses	★ 1st line treatment for uncomplicated gonorrhoeal infections	
ADR	-	<ul style="list-style-type: none"> • Nausea ,vomiting & diarrhoea . • Headache & dizziness. ★ May damage growing cartilage & cause arthropathy. ★ Phototoxicity, avoid excessive sunlight
C.I	-	<ul style="list-style-type: none"> • Pregnancy • Nursing mothers • Children under 18 years.

Drug	Spectinomycin
MOA	<ul style="list-style-type: none"> • Inhibits protein synthesis by binding to 30S ribosomal subunits
P.K	<ul style="list-style-type: none"> • Given 2g I.M, once
Uses	★ Alternative treatment in patient cannot tolerate or be treated with cephalosporins or quinolones
ADR	<ul style="list-style-type: none"> • Pain at the site of injection . • Fever and Nausea . ★ Nephrotoxicity (not common)

According to WHO recommendations:

- First choice is 3rd generation cephalosporins.
- Second choice is Fluoroquinolones.
- Third choice is Spectinomycin.

Treatment of complicated Gonorrhoea

- **Complicated gonorrhoeal infections** : Spread through bloodstream into:
 - Eye
 - Joints
 - Heart valves
 - Brain
- It can also spread from a mother to a fetus during birth
 - Newborn eye infection (conjunctivitis) may lead to blindness
- WHO guidelines suggest one of the following options for topical application to **both eyes immediately after birth**:
 - Silver nitrate 1% solution
 - Erythromycin 0.5% eye ointment
 - Tetracycline hydrochloride 1% eye ointment
 - Povidone iodine 2.5% solution (water-based)
 - Chloramphenicol 1% eye ointment

Drugs	Silver nitrate 1% solution	Erythromycin
Info	<ul style="list-style-type: none"> ● It has germicidal effects due to precipitation of bacterial proteins by liberated silver ions. 	<ul style="list-style-type: none"> ● 0.5% ointment for treatment & prevention of corneal & conjunctival infections.
Use	<ul style="list-style-type: none"> ● Put into conjunctival sac immediately after birth (no later than 1 hr after delivery) 	

Quiz

MCQ

Q1- A 22-year-old sexually active man presents to the Ambulatory care clinic with dysuria, penile discharge, and a swollen right knee. A joint aspirate of his right knee reveals many neutrophils as well as some gram negative diplococci. Which is the best choice to treat his condition?

A- Ceftriaxone B- Cephalexin C- Dexamethasone D - Meropenem

Q2-A 26-year-old sexually active HIV-negative man presents to his primary care physician with a nonpruritic maculopapular rash on his palms. He reports that about 6 weeks ago, he developed a non painful ulcer on his penis that healed spontaneously. He is injected with a single dose of benzathine penicillin G intramuscularly and sent home. What, if anything, should have been done differently for this patient's care?

A - A different antibiotic class should have been used
B - A different preparation of penicillin G (not benzathine) should have been used
C- Nothing—the course of action taken is entirely appropriate
D - Penicillin V should have been given instead of penicillin G

Q3-Which one of the following is the drug of choice for treatment of syphilis?

A.- ceftriaxone B - Benzylpenicillin penicillin G C. silver nitrate D - Ciprofloxacin

Q4- Which of the following is mechanism of action of Azithromycin?

A- Inhibit bacterial cell wall synthesis
B- Inhibit DNA synthesis by inhibiting DNA gyrase enzyme
C- Inhibits bacterial protein synthesis by binding to bacterial 50S ribosomal subunits.
D-Inhibit bacterial protein synthesis by reversibly binding to 30 S bacterial ribosomal subunit

Q5-Which of the following is a contraindication for Fluoroquinolones ?

A - Pregnancy B - Renal failure C- Hepatic failure D- Bleeding disorders

SAQ

A 24- year- old sexually active young man presents to the emergency department with dysuria, penile discharge, and a swollen right knee .A joint aspirate of his right knee reveals many neutrophils as well as some gram negative diplococci. Which is the best choice to treat his condition?

Q1-Which is the best choice to treat his condition?

Q2- Mention its MOA.

Q3- Enumerate 3 ADRS.

Answers:

MCQ

Q1	A
Q2	C
Q3	B
Q4	C
Q5	A

SAQ

Q1	Ceftriaxone
Q2	Inhibits bacterial cell wall synthesis.(Bactericidal)
Q3	Hypersensitivity reactions,Thrombophlebitis,Super infection Diarrhea

Thank you for all the love and support you gave the team in those two years!

Hope we made the context much easier to study.

God bless you, Future doctors.



Team Leaders:

May Babaeer

Zyad Aldosari

This Amazing Work was Done By:

Mohsen Almutairi

Abdullah Alassaf

Bader Aldhafeeri

Note writers

Nouf AlShammari

Quiz writers

Abdullah Alassaf