Vitamins B₆ and B₁₂

- General biochemistry
- Functions
- Deficiency diseases

Classification of Vitamins



Water-Soluble Vitamins

B vitamins

 Thiamin (B₁), riboflavin (B₂), niacin (B₃), pantothenic acid (B₅), pyridoxine (B₆), biotin (B₇), cobalamin (B₁₂), folate

- Not significantly stored in the body
- Must be supplied regularly in the diet
- Excess excreted

Vitamin B Complex

- Present in small quantities in different types of food
- Important for growth and good health
- Help in various biochemical processes in cell
- Function as coenzymes



Three forms

- Pyridoxine
- Pyridoxal
- Pyridoxamine
- Active form

 All 3 are converted to pyridoxal phosphate (PLP)



Figure 28.11. Structures of vitamin B₆.

Textbook of Biochemistry With Clinical Correlations, Sixth Edition, Edited by Thomas M. Devlin. Copyright © 2006 John Wiley & Sons, Inc.

Functions of Vitamin B_6

As coenzyme for

- Transamination
- Deamination
- Decarboxylation
- Condensation reactions



Figure 28.12. Some important metabolic roles of pyridoxal phosphate.

Textbook of Biochemistry With Clinical Correlations, Sixth Edition, Edited by Thomas M. Devlin. Copyright © 2006 John Wiley & Sons, Inc.

Condensation Reaction

Formation of ALA by ALA synthase, The regulatory step in hemoglobin synthesis



Decarboxylation Reaction:

Formation of Chatecholamines: Dopamine, norepinephrine and epinephrine





Decarboxylation Reaction:

Formation of Serotonin



Transamination Reaction



Disorders of Vitamin B₆ Deficiency

- Dietary deficiency is rare, but it was observed in:
 - Newborn infants fed on formulas low in B6
 - Women on oral contraceptives
 - Alcoholics

 Isoniazid treatment for tuberculosis can lead to vitamin B₆ deficiency by forming inactive derivative with PLP

Disorders of Vitamin B₆ Deficiency

- Deficiency leads to poor activity of PLP-dependent enzymes Causing:
 - Deficient amino acid metabolism
 - Deficient lipid metabolism
 - Deficient neurotransmitter synthesis
 [serotonin, epinephrine, norepinephrine and gamma amino butyric acid (GABA)]

Disorders of Vitamin B₆ Deficiency Cont'd

- PLP is involved in the synthesis of sphingolipids
- Its deficiency leads to demyelination of nerves and consequent peripheral neuritis

Disorders of Vitamin B₆ Deficiency

- Mild deficiency leads to:
 - Irritability
 - Nervousness
 - Depression

Severe deficiency leads to:
 Peripheral neuropathy
 Convulsions

Forms of Vitamin B₁₂

- Cyanocobalamin
- Hydroxycobalamin
- Adenosylcobalamin (major storage form in the liver)
- Methylcobalamin (mostly found in blood circulation)

Coenzyme forms of B_{12}

- Adenosylcobalamin and Methylcobalamin
 Coenzymes for metabolic reactions
- Body can convert other cobalamins into active coenzymes



Copyright © 2008 Wolters Kluwer Health | Lippincott Williams & Wilkins



(Cobalamin)

 Mainly found in animal liver bound to protein as
 Methylcobalamin or
 5'-deoxyadenosylcobalamin

Vitamin B₁₂

- Essential for normal nervous system function and red blood cell maturation
- Not synthesized in the body and must be supplied in the diet
- Binds to intrinsic factor and absorbed by the ileum
- Intrinsic factor is a protein secreted by cells in the stomach





Vitamin B₁₂ Storage

Liver stores vitamin B₁₂ (4-5 mg)

- Other B vitamins are not stored in the body
- Vitamin B₁₂ deficiency is observed in patients with IF deficiency due to autoimmunity or by partial or total gastrectomy
 - Clinical deficiency symptoms develop in several years

Functions of Vitamin B₁₂

- Two reactions require B₁₂ (1) Conversion of propionyl-CoA to succinyl-CoA
- The enzyme in this pathway, methylmalonyl-CoA mutase, requires B₁₂



Copyright © 2008 Wolters Kluwer Health | Lippincott Williams & Wilkins

Functions of Vitamin B₁₂

- (2) Conversion of homocysteine to methionine
- Methionine synthase requires B₁₂ in converting homocysteine to methionine



B₁₂ Deficiency and Folate Trap

- Homocysteine re-methylation reaction is the only pathway where N⁵-methyl TH4 can be returned back to tetrahydrofolate pool
- Hence folate is trapped as

N⁵-methyltetrahydrofolate (folate trap)

 This leads to folate deficiency and deficiency of other TH4 derivatives (N⁵-N¹⁰ methylene TH4 and N¹⁰ formyl TH4) required for purine and pyrimidine syntheses

TH4: Tetrahydrofolate

Interconversion between TH4 carrier of "one-carbon units"



Disorders of Vitamin B₁₂ Deficiency

Pernicious anemia

- Megaloblastic anemia
- Vitamin B₁₂ deficiency is mainly due to the deficiency of intrinsic factor

Disorders of Vitamin B₁₂ Deficiency

Demyelination

- Myelin sheath of neurons is chemically unstable and damaged
- Neuropathy
- Peripheral nerve damage

Disorders of Vitamin B₁₂ Deficiency

Causes of neuropathy

- Deficiency of vitamin B₁₂ leads to accumulation of methylmalonyl CoA
- High levels of methylomalonyl CoA is used instead of malonyl CoA for fatty acid synthesis
- Myelin synthesized with these abnormal fatty acids is unstable and degraded causing neuropathy

Neuropsychiatric symptoms of Vitamin B₁₂ Deficiency

Neurological symptoms

- Paraesthesia (abnormal sensation) of hands and feet
- Reduced perception of vibration and position
- Absence of reflexes
- Unsteady gait and balance (ataxia)

Neuropsychiatric symptoms of Vitamin B_{12} Deficiency

- Psychiatric symptoms
- Confusion and memory loss
- Depression
- Unstable mood

References

- Lippincott's Illustrated Reviews in Medical Biochemistry
- Textbook of Medical Biochemistry with Clinical Correlations by Thomas M Devlin
- Harper's Illustrated Biochemistry