

# Pathophysiology of Epilepsy



- 
- \* Seizures
  - \* Epilepsy

# Definition

## \* Seizure (Convulsion)

- Clinical manifestation of synchronised electrical discharges of neurons
- \* Seizures are symptoms of a disturbance in brain function , which can be due to epilepsy or other causes
- \* A seizure is a sudden surge in electrical activity in the brain that causes an alteration in sensation, behavior, or consciousness

# Epilepsy

Present when 2 or more unprovoked seizures occur at an interval greater than 24 hours apart

- \* Sudden recurrent episodes of sensory disturbance
- \* +/- Loss of consciousness, or convulsions
- \* Associated with abnormal electrical activity in the brain
- \* Abnormal , excessive electrical discharge of a group of neurons within the brain.
- \* When a person has recurrent ( 2 or more) , unprovoked seizures → " epileptic " .
- \* Seizures is symptom of epilepsy

## \* Provoked seizures

\* Seizures induced by somatic disorders originating outside the brain

- \* Fever
- \* Infection
- \* Syncope
- \* Head trauma
- \* Hypoxia
- \* Toxins
- \* Cardiac arrhythmias



# Classification of Seizures

- \* Seizures
- \* Partial
- \* or
- \* Generalized

## Partial (or Focal) Seizures

- **Simple Partial**
  - Awareness not impaired
- **Complex Partial**
  - Awareness impaired/lost
- **Partial Seizures secondarily generalizing**

## Generalized Seizures

- **Absence**
  - Typical
  - Atypical
- **Myoclonic**
- **Clonic**
- **Tonic**
- **Tonic-Clonic**
- **Atonic**

\* Focal / Partial seizures → their onset ( start) is limited to part of the cerebral hemisphere

\* Generalized seizures → those that involve the cerebral cortex diffusely ( whole of it ) from the beginning  
(*generalized seizures*)

## \* partial seizures

### a. Simple partial seizures

manifest

motor, somatosensory, and psychomotor symptoms  
without impairment of consciousness

### \* b. Complex partial seizures

manifest

impairment of consciousness with or without simple partial  
symptoms




**Impairment of consciousness:**  
cognitive, affective symptoms



Dreamy state; blank, vacant expression; déjà vu; jamais vu; or fear

## Complex Partial Seizures



**Formed auditory hallucinations.** Hears music etc



**Formed visual hallucinations.** Sees house, trees that are not there



Bad or unusual smell


**Olfactory hallucinations**

**Psychomotor phenomena.** Chewing movements, wetting lips, automatisms (picking at clothing)



**Dysphasia**

### EEG: left temporal lobe seizure




Repetitive sharp waves over left temporal region

## Generalized seizures

- \* Manifest a loss of consciousness  
Convulsive or non-convulsive
- \* (1) Generalized tonic-clonic seizures  
(Grand Mal epileptic seizure )
- \* (2) Absence seizures  
(Petit mal epileptic seizures)


# Primary generalized seizure



## A Partial seizure


### 1 Spread

Seizure focus



### 2 Secondary generalization

Seizure focus





\* The onset of a seizure:


Small group of abnormal neurons undergo

- Prolonged depolarizations
- Rapid firing of repeated action potentials

\* Spread to adjacent neurons or neurons with which they are connected into the process.



# Generalized

- \* 1- Generalized tonic-clonic (grand mal) seizure
- \* +/- aura
- \* (peculiar sensation or dizziness; then sudden onset of seizure with loss of consciousness)
- \* Tonic phase : Rigid muscle contraction in which clenched jaw and hands; eyes open with pupils dilated; lasts 30 to 60 seconds
- \* Clonic phase: Rhythmic, jerky contraction and relaxation of all muscles in with incontinence and frothing at the lips; may bite tongue or cheek, lasts several minutes.
- \* Postictal state: Sleeping or dazed for up to several hours.




A clinical seizure occurs when the electrical discharges of a large number of cells become abnormally linked together, creating a storm of electrical activity in the brain.

- \* Seizures may then spread to involve adjacent areas of the brain or through established anatomic pathways to other distant areas.




## \* 2. Absence (petit mal) seizure

- \* a. Loss of contact with environment for 5 to 30 seconds.
- \* b. Appears to be day dreaming or may roll eyes, nod head, move hands, or smack lips.
- \* c. Resumes activity and is not aware of seizure.





- 
- \* The clinical manifestations of a seizure reflect the area of the brain from which the seizure begins (i.e., seizure focus) and the spread of the electrical discharge.

\* Clinical manifestations accompanying a seizure are numerous and varied, including →

- \* (1) indescribable bodily sensations,
- \* (2) "pins and needles" sensations,
- \* (3) smells or sounds,
- \* (4) fear or depression,
- \* (5) hallucinations,
- \* (6) momentary jerks or head nods,
- \* (7) Déjà vu (over familiarity)
- \* (8) staring with loss of awareness, and
- \* (9) Convulsions → i.e., involuntary muscle contractions) lasting seconds to minutes.



# Aetiology of seizures


- **Epileptic**
  - Idiopathic (70-80%)
  - Cerebral tumor
  - Neurodegenerative disorders
  - Secondary to
    - Cerebral damage: e.g. congenital infections, intraventricular hemorrhage
    - Cerebral dysgenesis/malformation: e.g. hydrocephalus

# Aetiology of seizures

- **Non-epileptic**
  - Febrile convulsions
  - Metabolic
    - Hypoglycemia
    - HypoCa, HypoMg, HyperNa, HypoNa
  - Head trauma
  - Meningitis
  - Encephalitis
  - Poisons/toxins

# Pathophysiology of Epilepsy ( at molecular level)

- Cortical cell membrane level
  - Instability of the nerve cell membrane → Polarization abnormalities (excessive polarization ,hypopolarization , or lapses in repolarization),
  - The cell to be more susceptible to activation → Hypersensitive neurons with lowered thresholds for firing and firing excessively , related to →

- 
- (1) Excess of Excitatory ( acetylcholine- or Glutamate - related activity )
  - (2) Decreased inhibitory ( GABA -related activity)
  - Together and/or (2) above → leading to instability of cell-membrane & lowered threshold for excitation → excessive polarization, hypopolarization allowing the cell to be more susceptible to activation spontaneously or by any ionic imbalances in the immediate chemical environment of neurons







## \* Electroencephalogram ( EEG )

- \* EEG >>>> diagnosis, classifying seizures >>>> therapeutic decisions
- \* **spikes or sharp waves** (Epileptiform EEG patterns)
  - Focal epileptiform discharges indicate focal epilepsy
  - Generalized epileptiform activity indicates a generalized form of epilepsy.

# Normal EEG Awake




# Generalized Tonic Clonic



# EEG: Partial Seizures



- 
- \* Some types linked to genes
  - \* (run in families)
  - \* Genetic abnormalities >>>> increasing a person's susceptibility to seizures that are triggered by an environmental factor
  - \* Several types of epilepsy have now been linked to defective genes for ion channels

# Pathophysiology

- \* Genetic factors
  - \* At least 20 %
  - \* Some examples
    - \* Benign neonatal convulsions--20q and 8q
    - \* Juvenile myoclonic epilepsy--6p
    - \* Progressive myoclonic epilepsy--21q22.3

