

# Physiology of Motor Tracts

*Dr. Salah Elmalik*

# Neuropsychiatry Block

## Chapter 56

### Cortical & Brain Stem Control of Motor Function

(**Guyton & Hall**)

Reference book/ Ganong review of  
medical physiology

# Learning objectives:

At the end of this lecture the student should :

- Appreciate what is upper motor neuron and lower motor neuron .
- Explain the origin , course and functions of the following motor tracts:
  
- Pyramidal tracts:
  - (corticospinal & corticobulbar tracts)
- Extrapyramidal tracts as:
  - ❖ Rubrospinal .
  - ❖ Vestibulospinal .
  - ❖ Reticulospinal .
  - ❖ Olivospinal .
  - ❖ Tectospinal

# Components of Motor Neurons

For performance of motor acts we need:

## Upper motor neurons (UMN)

- These are the motor neurons whose cell bodies lie in the motor cortex, or brainstem and they activate the lower motor neuron (LMN)
- There are two UMN Systems :-
- 1- Pyramidal system (corticospinal tracts ).
- 2- Extrapyramidal system

## Lower motor neurons: (LMN)

- These are the motor neurons of the spinal cord (AHCs) and brain stem motor nuclei of the cranial nerves that innervates skeletal muscle directly.

# Classification of descending motor systems


- The descending motor pathways have commonly been divided into "*pyramidal*" and "*extrapyramidal*" tracts.
- This classification is based on the finding that the motor tract which originates from the cerebral cortex and descends to the spinal cord (the corticospinal tract) passes through the pyramids of the medulla, and therefore has been called the "*the pyramidal tract*".
- The rest of the descending motor pathways do not travel through the medullary pyramids, and are therefore collectively gathered under the heading: "*the extrapyramidal tracts*".

The following are the important sets of descending motor tracts, named according to the origin of their cell bodies and their final destination:

- 1) Corticospinal tract,
- 2) Corticobulbar tract.

(= **Pyramidal tract**)

- 1) Rubrospinal tract,
- 2) Reticulospinal tracts,
- 3) Vestibulospinal tracts,
- 4) Tectospinal tract, and
- 5) Olivospinal tract .





**Extrapyramidal  
tracts**

# Motor Areas

## (1) The primary motor area ( M1 . Motor area 4)

- occupies the precentral gyrus & contains large , highly excitable Betz cells.
- MI of one side controls skeletal muscles of the opposite side of the body
- Feet are at the top of the gyrus and face at the bottom , arms and the hand area in the mid portion
- Facial area is represented bilaterally, but rest of the representation is generally unilateral
- Area of representation is proportional with the complexity of function done by the muscle. So, muscles of hands and speech (lips, tongue , and vocal cord) occupies 50% of this area
- The neurons of this area arranged in vertical Columns.



MOTOR HOMUNCULUS



## 2. The Supplementary Motor Area (M2)


- Lies in front of area 4 and above the premotor area
- Stimulation of this area leads to bilateral grasping movements of both hands simultaneously.
- This area involved in organizing or planning motor sequence

### 3. Premotor Area ( M3)

- lies in front of the primary motor area & below supplementary motor area.
- Stimulation of the premotor area produces complex coordinated movements, such as setting the body in a certain posture to perform a specific task.
- It works in association with the supplemental motor area, establishing the motor programs necessary for execution of complex movements
- It contains mirror neurons which are important for understanding the action of other people and for learning new skills by imitation.

# Premotor Area cont....

- ▶ A few highly specialized motor centers have been found in the premotor areas of the human cerebral cortex:
- ▶ Broca's Area for Speech
- ▶ The Frontal Eye Movements Area:
  - ✓ located above Broca's area in the frontal lobe
  - ✓ controls voluntary movements of the eyes toward different objects in the visual field.
- ▶ Head Rotation Area:
  - ✓ located just above the eye movement area in the motor cortex .
  - ✓ directing the head toward different visual objects .
- ▶ Area for Hand Skills
  - *Damage to this area causes motor apraxia*




# Pyramidal Tracts


# Corticospinal ( Pyramidal) Tracts

## Cells of origin :


- 30% originate from the primary motor area,
- 30% from the premotor areas , and supplementary motor areas
- 40% from the somatic sensory areas posterior to the central sulcus.
  - ✓ 3% of the fibres are large myelinated fibres, derived from the large , highly excitable pyramidal Betz cells of MI . These fibers form monosynaptic connections with motor neurons of the spinal cord.
  - ✓ But most of pyramidal fibers are unmyelinated
  - ✓ Fibers from the cerebral cortex descend in corona radiata to reach the posterior limb of internal capsule (between caudate and putamen nuclei of the basal ganglia)

- Then descend through the cerebral peduncle of midbrain and then through pons.
- ✓ Some fibers cross in brainstem to supply contralateral cranial nerve nuclei constitute the Corticobulbar tract .
- In the lower medulla
- around 80% of the fibres cross to the opposite side and descend in the lateral column of spinal cord as the Lateral Corticospinal Tract .
- Some of these fibres end monosynaptically on contralateral anterior horn cells but the majority end on interneurons in the intermediate region of the spinal cord grey matter
- *These fibers controls and initiates fine discrete skilled movement of distal limb muscles (i.e. Fingers and hands)*






Descending lateral corticospinal pathway





- The remaining 20 % of corticospinal fibers do not decussate in the medulla .
- They descend ipsilaterally in the ventral column of the spinal cord white matter , Constituting the Ventral (Anterior) Corticospinal Tract .
- Finally they decussate (cross to the opposite side ) & synapse on the contralateral spinal motor neurons
- *These fibers control the axial and proximal limbs muscles so it concern with control of posture.*



## Course and Termination

*fibers of the CBS tract descend from the cerebral cortex*

*collect together and descend through the posterior limb of the internal capsule*

*through the middle portion of the cerebral peduncles of the midbrain*

*The fibers are separated by transverse pontine fibers in the pons*

*In the upper medulla oblongata where they form the pyramids of the medulla*

*In the lower region of the medulla, most of the fibers cross to the opposite site forming the "medullary decussation"*

*About 80% to 90% cross to the opposite side of the spinal cord (contralaterally) and continue as the "lateral corticospinal tract".*

*fibers which do not decussate in medulla (about 10-20%) descend on the same side of the spinal cord (ipsilaterally) as the "ventral corticospinal tract".*

# Functions of Pyramidal System

- 1) Initiation of voluntary movements
- 2) The lateral corticospinal tract fibers that descend in the spinal cord for control of muscles of the distal parts of the limbs, especially the hand and digits muscles, which sub serve fine skilled movements used in manipulation by hand and fingers, and other accurate motor actions done by the limbs .
- 3) The ventral corticospinal tracts control posture of axial and proximal muscles for balance , climbing and walking.
- 4) Facilitation of muscle tone and deep reflexes through gamma motor neurons
- 5) Those fibers originate from parietal lobe are for sensory-motor coordination
- 6) Corticobulbar tracts /control face & neck muscles & facilitate their tone, and are involved in facial expression, mastication, swallowing

# Extrapyramidal Tracts

# Extrapyramidal Tracts

- The extrapyramidal system is made up of all those part in the central nervous system that are concerned with motor control , other than pyramidal system. They consist of ;
- 1.Cortical motor areas, especially the premotor area and parietal cortex
- 2. The basal ganglia
- 3. The reticular formation , the red nuclei , the tectum of brain and vestibular nuclei
- They include the following tracts:

# Rubrospinal Tracts


- After emerging from Red Nucleus in midbrain, fibers decussate at same level of red nucleus
- It receives afferent connections from:
  - Ipsilateral cortical motor area (corticorubral tract)
  - Contralateral side of cerebellum
  - Basal ganglia




- Descend with the lateral corticospinal tract



- In spinal cord tract occupies the lat. white column, & fibers synapse on the contralateral AHCs





**Figure 55-5**


Corticorubrospinal pathway for motor control, showing also the relation of this pathway to the cerebellum.

# Functions of Rubrospinal Tract


- 1) An additional pathway for transmission of cerebral cortical motor commands to the lower motor neurons similar to those of the corticospinal tract. When the corticospinal fibers are destroyed, discrete movement can still occur but fine control of the fingers and hand is impaired..
  - 2) This tract is excitatory for flexors & inhibitory for extensors (anti-gravity muscles).
- ✓ Rubrospinal tract lies in the lateral columns of the spinal cord, along with the corticospinal tract, and terminates on the interneurons and motor neurons that control the more distal muscles of the limbs.
  - ✓ Therefore, corticospinal & rubrospinal tracts together are called the lateral system of the cord, in contradistinction to a vestibulo-reticulospinal system which lies mainly medially in the cord and is called **the medial motor system of the cord**



**A** Medial brain stem pathways




**B** Lateral brain stem pathways




# Vestibulospinal Tracts

# Vestibulospinal Tracts

- 2 components of the tract:
- Major component → Lat. V.S Tract
- Minor component → Med. V.S Tract.
- Lat. V.S Tract ← lat. Vest. Nuclei.
- Med. V.S Tract ← med Vest. Nuclei.





**Figure 55-13**

Connections of vestibular nerves through the vestibular nuclei (the large oval white area) with other areas of the central nervous system.

# Functions of Vestibulospinal Tracts

- Vestibulospinal tracts control reflexes e.g. Postural & righting ( which correct body position ) + control eye movements.

## The lateral vestibulospinal


- Cells of origin : Lateral Vestibular Nucleus
- Axons descend in the ipsilateral ventral white column of spinal cord .
- *This tract mediates excitatory influences upon extensor motor neurons to maintain posture (when the otolith organs signal deviations from stable balance and upright posture)*

## The medial vestibulospinal tract :

- Cells of origin : Medial Vestibular Nucleus
- As its axons descend ipsilaterally in the ventral white column of spinal cord , they form part of the Medial Longitudinal Fasciculus
- *The medial longitudinal fasciculus consists of both ascending & descending fibers that link vestibular nuclei to nuclei supplying the extraocular muscles for coordination of head and eye movements*


# Reticulospinal tracts

- Tract arises from reticular formation (groups of scattered neurons along with nerve fibers present in midbrain, pons & medulla).
- It has two components:
  - Pontine reticulospinal tract
  - Medullary reticulospinal tract



Vestibulospinal and reticulospinal tracts descending in the spinal cord to excite (*solid lines*) or inhibit (*dashed lines*) the anterior motor neurons that control the body's axial musculature.

(B) RETICULOSPINAL TRACT



# The Reticulospinal Tracts

## Divisions:

1) *Medial (pontine) reticulospinal tract.* 2) *Lateral (medullary) reticulospinal tract.*

### *Medial reticulospinal tract*

- Arises from neurons of the "pontine reticular formation"
- descends to all levels of the spinal cord
- Terminate mainly on interneurons in the spinal gray matter which excite the medially situated  $\alpha$ - and  $\gamma$ -MNs innervating the antigravity muscles, that is, the muscles of the vertebral column and the extensor muscles of the lower limbs.

### *Lateral reticulospinal tract*

- arises from neurons in the "medullary reticular formation"
- its fibers descend to all levels of the spinal cord
- synapse with interneurons that inhibit the  $\alpha$ - and  $\gamma$ -MNs of antigravity and extensor muscles, but they facilitate the  $\alpha$ - and  $\gamma$ -MNs of flexor muscles.



## ***Medial reticulospinal tract***

- The pontine reticular formation has a high degree of ***natural excitability***.
- In addition, it receives strong excitatory signals from the vestibular nuclei and the neocerebellum

### **Functions:**

- ***the medial (or pontine) reticulospinal tract is excitatory for extensors & inhibitory for flexors (unlike rubro-spinal).***

## ***Lateral reticulospinal tract***

- The medullary reticular formation receives afferent signals from: (i) the premotor area of cerebral cortex, (ii) the paleocerebellum, and (iii) red nucleus.

### **Functions:**

- ***The medullary reticulospinal tract is inhibitory for extensors (like rubro-spinal).***

# Tectospinal Tract

- Originates in Superior Colliculus in the tectum of midbrain → then decussate in the dorsal tegmentum



- Axons descend in ventral white column of spinal cord




- And terminate on Contralateral cervical AHCs

## Function :

- *This tract produces reflex turning of the head and neck ( in response to visual and auditory stimuli)*


(C) COLLICULOSPINAL TRACT



# Olivospinal Tract

- Originates in Inferior olivary nucleus of the medulla is found only in the cervical region of the spinal cord.
- Function is uncertain, but thought to **facilitate muscle tone**

# Summary of motor Tracts



# General functions of Extrapyramidal tract

1. Help pyramidal tract in initiation of voluntary movement
2. Share in planning and programming of voluntary movement
3. Responsible for subconscious gross movements (swinging of arms during walking)
4. Keep equilibrium and adjust body posture
5. Regulation of muscle tone

# Removal of (Area Pyramidalis) of the Primary Motor Cortex

- Removal of the area that contains the giant Betz pyramidal cells (Area Pyramidalis) causes loss of voluntary control of discrete movements of the distal segments of the limbs, especially of the hands and fingers (This does not mean that the hand and finger muscles themselves cannot contract (paralysis), *but the ability to control the fine movements is gone*)
- That is because area pyramidalis is essential for voluntary initiation of finely controlled movements, especially of the hands and fingers

## Effects of Lesions in the Motor Cortex or in the Corticospinal Pathway(The stroke)

- ▶ The motor control system can be damaged by the "stroke -the result is loss of blood supply to the cortex or to the corticospinal tract where it passes through the internal capsule.
- ▶ Muscle Spasticity Caused by Lesions That Damage Large Areas Adjacent . to the Motor Cortex
- ▶ The primary motor cortex normally exerts a continual tonic stimulatory effect on the motor neurons of the spinal cord; when this stimulatory effect is removed, hypotonia results.
- ▶ Most lesions of the motor cortex, especially those caused by a stroke, involve the primary motor cortex & adjacent parts of the brain such as the basal ganglia. In these instances, muscle spasm occurs in the muscles on the opposite side of the body.

**THANK YOU**