
Biochemistry of Vitamin K

GI and Nutrition Block

Dr. Sumbul Fatma

Objectives

Upon completion of this lecture, the Second Year students will be able to:

- Identify the types and sources of vitamin K
- Understand the role of vitamin K in blood coagulation
- Recognize the importance of γ -carboxylation of glutamic acid in coagulation proteins
- Understand the role of anticoagulant drugs in affecting vitamin K function
- Discuss the causes and disorders of vitamin K deficiency

Overview

- Types, chemistry and sources of vitamin K
- Sources and daily requirements
- Functions:
- Synthesis of γ -carboxyglutamate in:
 - Prothrombin and blood clotting factors
 - Interaction of prothrombin with platelets
 - Osteocalcin
 - Protein C and S (anticogaulant proteins)
- Deficiency and disorders
- Clinical manifestations

Types and Sources



Occurs in several forms:

- Vitamin K₁ (Phylloquinone)
- Vitamin K₂ (Menaquinone)
- Vitamin K₃ (Menadione) – synthetic form

Dietary sources:

- Cabbage, kale, spinach, egg yolk, liver



Cabbage




Kale




Spinach


Chemistry of Vitamin *V*



Vitamin K₁ (Phylloquinone)



Vitamin K₂ (Menaquinone)



Vitamin K₃ (Menadione)

Sources of Vitamin K




- **Phylloquinone:** Green leafy vegetables
- **Menaquinone:** Intestinal bacteria
 - Intestinal bacterial synthesis meets the daily requirement of vitamin K even without dietary supplement
- **Menadione:** synthetic form
 - A precursor of menaquinone

RDA for Vitamin K

($\mu\text{g}/\text{day}$)



- Infant (0-1 year): 2-2.5
 - Children (1-8): 30-55
 - Men (19+): 120
 - Women (19+): 90
 - Pregnancy / lactation: 90 / 90
 - UL: Not established
-



Functions of Vitamin K



- Coenzyme for the synthesis of prothrombin and blood clotting factors in the liver
 - Prothrombin and clotting factors are protein in nature
 - Synthesis of prothrombin, clotting factors II, VII, IX, X require carboxylation of their glutamic acid (Glu) residue





Figure 28.7. Function of Vitamin K.



Functions of Vitamin K



- ❑ Mature prothrombin and clotting factors contain γ -carboxyglutamate (Gla) after carboxylation reaction
- ❑ Vitamin K is essential for the carboxylase enzyme involved
- ❑ Dihydroquinone form of vitamin K is essential for this reaction





Figure 28.7. Function of Vitamin K.




Analogues of Vitamin K



- Anticoagulant drugs: warfarin and dicoumarol
 - Structural analogs of vitamin K
- They inhibit the activation of vitamin K to hydroquinone form (inhibiting the reductase enzyme)
- Prothrombin and clotting factors are not carboxylated
- Hence blood coagulation time increases upon injury

- Carboxylation of glutamate requires vitamin K
- The process is inhibited by warfarin





Functions of Vitamin K



- **Prothrombin – platelet interaction**
 - ❑ Carboxylated prothrombin contains two carboxylate groups (COO^-)
 - ❑ These groups bind to Ca^{2+} forming prothrombin-calcium complex
 - ❑ The complex then binds to phospholipids on the surface of platelets (important for blood clotting)
 - ❑ Converting prothrombin to thrombin and initiating clot formation
-

Prothrombin – platelet interaction





Functions of Vitamin K



- **Synthesis of γ -carboxyglutamate in osteocalcin**
 - ❑ Osteocalcin is a bone turnover protein
 - ❑ Also called Bone Gla Protein (BGP)
 - ❑ Involved in bone formation, mineralization and resorption
 - ❑ γ -Carboxyglutamate is required for osteocalcin binding to hydroxyapatite (a calcium mineral) in the bone
 - ❑ The binding mechanism is similar to that of prothrombin-platelet binding
-

Deficiency of Vitamin K



- Deficiencies are rare: it is synthesized by intestinal bacteria
- Hypoprothrombinemia: increased blood coagulation time
- Some second-generation cephalosporin drugs cause this condition due to warfarin-like effects (antibiotics given with vit. K)
- May affect bone growth and mineralization
- Lipid malabsorption can lead to vitamin K deficiency

Deficiency of Vitamin K



- Prolonged antibiotic therapy
 - Especially in marginally malnourished individuals (eg debilitated geriatric patients)

 - Gastrointestinal infections with diarrhea

 - Both of the above destroy the bacterial flora leading to vitamin K deficiency
-

Deficiency of Vitamin K



- Deficiency most common in newborn infants
 - Newborns lack intestinal flora
 - Human milk can provide only 1/5th vitamin K
 - Supplements are given intramuscularly at birth

Clinical Manifestations of the Deficiency



- Hemorrhagic disease of the newborn
 - Bruising tendency, ecchymotic patches (bleeding underneath the skin)
 - Mucus membrane hemorrhage
 - Post-traumatic bleeding / internal bleeding
 - Prolonged prothrombin time
-

Toxicity of Vitamin K



- Prolonged supplementation of large doses of menadione can cause:
 - Hemolytic anemia
 - Jaundice
 - Due to toxic effects on RBC membrane
-

Take home message



- Vitamin K is essential for blood coagulation process
 - It mediates the process by γ -carboxylation of glutamic acid residues of prothrombin and coagulation factors
-

References



- Lippincott's Biochemistry 6th Edition
Chapter 28, pp. 389-391
Richard Harvey and Denise Ferrier
Lippincott Williams & Wilkins, USA
 - Textbook of Biochemistry with Clinical
Correlations by Thomas M Devlin. 6th Edition
Chapter 28, pp. 1099-1101
-