Myeloproliferative Neoplasms

- 1. Myeloproliferative neoplasms (MPN)
 - 1.1. Chronic myelogenous leukemia, BCR-ABL1-positive (CML)
 - 1.2. Polycythemia vera (PV)
 - 1.3 Essential thrombocythemia (ET)
 - 1.4 Primary myelofibrosis (PME)
 - 1.5. Chronic noutrophilic loukomia (CNL)
 - 1.6. Chronic eosinophilic leukemia, not otherwise specified (CEL-NOS)
 - 1.7. Mast cell disease (MCD)
 - 1.8. MPN, unclassifiable

MPN features

- Cytosis
- Organomegaly (mainly splenomegaly)
- High uric acid
- Hypercellular bone marrow
- Progression to acute leukaemia (mainly AML)

Table 1. Classification of Myeloid Neoplasms According to

 the 2008 World Health Organization Classification

 Scheme

1. Myeloproliferative neoplasms (MPN)

- 1.1. Chronic myelogenous leukemia, BCR-ABL1-positive (CML)
- 1.2. Polycythemia vera (PV)
- 1.3. Essential thrombocythemia
- 1.4. Primary myelofibrosis (PMF)
- 1.5. Chronic neutrophilic leukemia
- 1.6. Chronic eosinophilic leukemia

BCR-ABL <u>must be</u> negative

- 1.7. Mast cell disease (MCD)
- 1.8. MPN, unclassifiable
- 2. Myeloid and lymphoid neoplasms with eosinophilia and abnormalities of PDGFRA, PDGFRB, and FGFR1

3. MDS/MPN


- 3.1. Chronic myelomonocytic leukemia (CMML)
- 3.2. Juvenile myelomonocytic leukemia (JMML)
- 3.3. Atypical chronic myeloid leukemia, BCR-ABL-negative (aCML)
- 3.4. MDS/MPN, unclassifiable
- 4. Myelodysplastic syndromes (MDS)
- 5. Acute myeloid leukemia (AML)

1/18/21


Polycythemia


- In Greek "too many cells in the blood.".
- Absolute increase in total body red cell volume (or mass)
- Manifests itself as a raised Hb or packed cell volume (PCV)
- Hb is >16.5or 18.5 g/dl in women and men, respectively

Regulation of Erythropoiesis



Classification of Polycythemia





Polycythemia Vera

•MPN characterized by increased red blood cell production independent of the mechanisms that normally regulate erythropoiesis.

Diagnostic Features:

- •HB >18.5g/dl in men ,16.5g/dl in women
- •Hypercellular bone marrow
- •JAK2 mutation in >95% of cases
- Low Serum erythropoietin level

Clinical features of PV

1-Increased blood viscosity

- Hypertension
- Headache, dizziness, visual disturbances & paresthesia

2- Thrombosis

- Deep vein thrombosis
- Myocardial infarction
- Mesenteric, portal or splenic vein thrombosis

3-Splenomegaly in 70% 4-Hepatomegaly in 40%

Investigations

CBC:

*RBC: increased *Hb: increased *WBC & PLT :mildly increased (usually) Blood smear:

- Excess of normocytic normochromic RBC
- ±Leukocytosis &thrombocytosis


Bone marrow


Hypercellular

Blasts -


- Predominant erythroid precursors
- ± Increased megakaryocytes & Myeloid precursors.

AL transformation





Complication & treatment



Primary Myelofibrosis

1. Myeloproliferative neoplasms (MPN)

- 1.1. Chronic myelogenous leukemia, BCR-ABL1-positive (CML)
- 1.2. Polycythemia vera (PV)
- 1.3. Essential thrombocythemia (ET)
- 1,4. Primary myelofibrosis (PMF)
- Chronic neutrophilic leukemia (CNL)
- 1.6. Chronic eosinophilic leukemia, not otherwise specified (CEL-NOS)
- 1.7. Mast cell disease (MCD)
- 1.8. MPN, unclassifiable


Primary Myelofibrosis

Clonal MPN characterized by a proliferation of megakaryocytes & granulocytes in the bone marrow that associated with deposition of fibrous connective tissue and extramedullary haematopoiesis


Clinical features

•Anemia


- Leukoerythroblastic blood picture.
- Massive splenomegaly
- Fibrotic bone marrow
- •JAK2 mutation (50%)
- Risk of AML transformation (20%)



Bone marrow in Myelofibrosis



Stages of PMF



Essential Thrombocythemia

1. Myeloproliferative neoplasms (MPN)

- 1.1. Chronic myelogenous leukemia, BCR-ABL1-positive (CML)
- 1.2. Polycythemia vera (PV)
- 1.3. Essential thrombocythemia (ET)
 - .4 Primary myelofibrosis (PMF)
- 1.5. Chronic neutrophilic leukemia (CNL)
- 1.6. Chronic eosinophilic leukemia, not otherwise specified (CEL-NOS)
- 1.7. Mast cell disease (MCD)
- 1.8. MPN, unclassifiable


Essential Thrombocythemia


ET is MPN that involves primarily the megakaryocytic lineage. & characterized by sustained thrombocytosis .

Diagnostic Features

- Sustained thrombocytosis ≥450×10⁹.
- Hypercellular BM with megakaryocytic proliferation
- Exclusion of: CML, MDS, PV & Primary Myelofibrosis
- JAK2 mutation (60%), if negative; no evidence of reactive thrombocytosis:

Iron def. ,splenectomy, surgery, infection, autoimmune disease....





Essential Thrombocythemia

Clinical Presentation

- Asymptomatic (50%)
- Thrombosis
- Bleeding
- Mild splenomegaly (50%)
- Mild hepatomegaly (20%)


Very indolent (5% risk of AML transformation)

Treatment

Aspirin ±Hydroxyuria

JAK2 Mutation

JAK2: Non receptor protein tyrosine kinase involved in signal transduction pathway



JAK2 mutation :

Point mutation (at codon 617 in JH2) leads to loss of auto inhibitory control over JAK2.

The mutated JAK2 is in a constitutively active state,

JAK2 Mutation

