



Intestinal Helminths

Dr. Ibrahim Alkhalife

PROTOZOA

Unicellular
Single cell for all functions

- 1: Amoebae: move by pseudopodia
- 2: Flagellates: move by flagella
- 3: Ciliates: move by cilia
- 4: Apicomplexa (Sporozoa)
tissue parasites

HELMINTHS

Multicellular
Specialized cells

- Round worms (Nematodes):
- elongated, cylindrical, unsegmented.
- Flat worms:
- Trematodes: leaf-like, unsegmented.
 - Cestodes: tape-like, segmented.

Nematodes: General features

1. **Elongated worm, cylindrical, unsegmented and tapering at both ends.**
2. **Variable in size, measure <1 cm to about 100cm.**
3. **Sex separate and male is smaller than female**



Nematodes: Location in the human body

- **Intestinal nematodes**
- **Tissue nematodes**

Nematodes: common intestinal infections

1. *Enterobius (Oxyuris) vermicularis* (Pinworm, seatworm, threadworm)
2. *Trichuris trichiura* (whipworm)
3. *Ascaris lumbricoides* (roundworm)
4. *Ancylostoma duodenale* & *Necator americanus* (hookworms)
5. *Strongyloides stercoralis*

1- Enterobius vermicularis (THREAD WORM)

(Common names: Pin worm, seat worm)

- Found all over the world but more common in temperate regions.
- Children are more often involved than adults, it tends to occur in groups living together such as families, army camps or nursery.
- Adult worms are mainly located in lumen of cecum and the female migrate to rectum to deposit her eggs on perianal skin.





Direct human - human infection occurs mainly by swallowing the eggs.

Autoinfection occurs by contamination of the fingers.

It can be seen by naked eye as white thread \pm 1cm.

Male is smaller than female \pm 0.5cm, with coiled end.

Enterobius vermicularis



Enterobius vermicularis (*Oxyuris*)

Pathology

- Most infections are asymptomatic
- Main clinical presentation **pruritus ani** which can be very troublesome and occurs more often during the night, persistent itching may lead to inflammation and secondary bacterial infection of the peri-anal region

***Enterobius vermicularis* (Oxyuris)**

Infected children may suffer from emotional disturbance, insomnia, anorexia, loss of weight and loss of concentration and enuresis.

Ectopic enterobiasis occurs in infected adult female when invade vulva and vagina result in valvo-vaginitis, salpingitis, also adult worm can lodge in the lumen of appendix cause appendicitis.

Enterobius vermicularis (Oxyuris)

DIAGNOSIS :

Unlike other intestinal Nematodes, the eggs are not usually found in feces. The best method is to look for them around the anus by taking an anal swab or by using


CELLULOSE ADHESIVE TAPE

the examination should be done before defecation or bathing.


Treatment

Albandazole or Mebendazole for whole family

Enterobius vermicularis (Oxyuris)



Ascaris lumbricoides (roundworm)



Ascaris lumbricoides (roundworm)

The commonest human helminths infection all over the world.

The large round worm which is normally located in **the small intestine**


- Found in **jejunum** and upper part of **ileum**
- Female \pm **20** cm longer than male \pm **10** cm
- Feed on semi digested food.




Ascaris lumbricoides (roundworm)

Infective stage:
embryonated egg


Diagnostic stage:
unembryonated egg



Ascaris lumbricoides life cycle



Ascaris eggs



Ascaris egg (embryonated)

Ascaris larva emerging from egg



Ascaris lumbricoides (roundworm)

Pathology:

- **1-Adult worm:** (small intestine)

Light infection : asymptomatic


Heavy infection : intestinal obstruction

Migrating adult : to bile duct-jaundice

- **2-Larvae:** Loeffler`s syndrome


Pneumonitis and broncho-spasm,
cough with bloody sputum, Eosinophilia, urticaria

Ascaris lumbricoides (roundworm)



Loeffler`s syndrome: Larvae in lung
pneumonia, cough, bloody sputum

Ascaris lumbricoides (roundworm)



Ascaris larva in lung

Ascaris lumbricoides (roundworm)

Diagnosis:

- eggs in stool.
- larvae in sputum.
- adult may pass with stool.




Treatment:
Albendazole or Mebendazole


2-Trichuris trichiura (Whipworm)



Life Cycle of
Trichuris trichiura



Trichuris trichiura



Infective stage:
embryonated egg

Diagnostic stage:
egg in stool

Trichuris trichiura (whipworm)

World wide, common in poor sanitation

- It coexists with *Ascaris* because of similar requirements (the eggs needs 3 weeks in the soil to be embryonated which is the infective stage).
- Adult live in **large intestine** especially **caecum** and **appendix**
- in heavy infection the whole length of large intestine affected.
- Male and female worm have **narrow anterior** portion penetrate the intestinal mucosa

Trichuris trichiura (Whipworm)

Pathology

- **light infection:** asymptomatic
- **heavy infection:** abdominal pain, bloody diarrhea.

Rectal prolapse in children is a common complication.



Trichuris trichiura (Whipworm)



- **Diagnosis:** egg in stool characterized by its barrel shape with mucoid plugs at each pole.



- **Treatment:** Albendazole.

Hook worms

Ancylostoma duodenale & *Necator americanus*




Its buccal capsule (mouth) lined with hard hooks, triangular **cutting plates** and **anticoagulant glands**.

There are no specific symptoms or signs of hookworm infection, but they give rise to a combination of intestinal inflammation and progressive iron-deficiency anemia and protein deficiency

Filariform Larval (infective stage) invasion of the skin can produce a skin disease called cutaneous larva migrans also known as *creeping eruption*, this is commonly caused by walking barefoot through areas contaminated with fecal matter. Larva migrate through the vascular system to the lungs, and from there up the trachea, and are swallowed. They then pass down the esophagus and enter the digestive system, finishing their journey in the small intestine where the larvae mature into adult worms. They mate inside the host, females laying up to 30,000 eggs per day, which pass out in feces (diagnostic stage). The eggs need to be in soil for about one week to become ***FILARIFORM LARVA***





Life cycle of Hook worms

Pathology & clinical picture

- larvae:

- **At the site** of entry of larvae intense itching (ground itch) and dermatitis.
- **Migration phase**
 - cough with bloody sputum
 - pneumonitis and bronchitis but less severe than *Ascaris*,
 - eosinophilia urticaria.

- Adult worm:

- low worm burden (INFECTION): **no** symptoms.
- Moderate to heavy burden:
 - Epigastric pain, vomiting, hemorrhagic enteritis.
 - Protein loss: hypo-proteinaemia edema.
 - **Anemia**: due to withdrawal of blood by parasites and hemorrhage from punctured sites lead to **severe anemia** = **microcytic hypochromic anemia**.

Hook worms

Diagnosis and treatment

■ Diagnosis:

- Eggs in stools;
- occult blood (+)



Treatment: Albendazol, Mebendazole


Strongyloides stercoralis


- Widely distributed in tropical area at Asia, Africa & South America
- Fatal dissemination in **immuno-compromised host.**
- It is **smallest** pathogenic nematodes \pm **2.5mm.**
- adult live membrane of duodenum, jejunum rarely mucous membrane of bronchus
- **Autoinfection** is very important criteria

Strongyloides stercoralis life cycle

- The parasite shows 3 different modes of development:
- **1-Direct development:** The rhabditiform larva pass from stool and become directly a Filariform larva if the environment of the soil is suitable.
- **2-Indirect development:** In external environment Rh. larva becomes free living adults, produce eggs, rhabditiform larva and Filariform larva (Infective stage).
- **3-AUTOINFECTION:**
 - Internal: when the rhabditiform larva become a filariform larva in the intestine and penetrate the intestine
 - External: fecal contamination of skin –Rh larva > filariform penetrates the skin

Strongyloides stercoralis





Strongyloides stercoralis

Pathology and clinical picture

- **Cutaneous** little reaction on penetration.
sever dermatitis at perianal region in case of external autoinfection
- **Migration:** pneumonitis during larval migration.
- **Intestinal:** inflammation of upper intestinal mucosa, diarrhea, upper abdominal pain in the epigastria colicky in nature.
- **Disseminated strongyloidiasis:** in patient with **immunodeficiency**, uncontrolled diarrhea –granulomatous changes –necrosis--perforation--peritonitis—death.

Strongyloides stercoralis

Diagnosis

rhabditiform larvae
diagnostic stage in:

- Stool examination
- Duodenal aspirate




Treatment

Albendazole, Mebendazole

Common Tapeworm Infections

<u>TAPEWORM</u>	DISEASE	TRANSMISSION OF INFECTION	LOCATION OF ADULT IN HUMANS	LOCATION OF LARVA IN HUMANS	CLINICAL PICTURE	LAB.
						DIAGNOSIS
<i>Taenia saginata</i>	taeniasis	ingestion of larva in undercooked beef	Small Intestine	not present	vague digestive disturbances	eggs or proglottids in stools
<i>Taenia solium</i> - <u>ADULT</u>	taeniasis	ingestion of larva in undercooked pork	Small Intestine	not present	vague digestive disturbances	eggs or proglottids in stools
<i>Taenia solium</i> - <u>LARVA</u> (cysticercus cellulosae)	Cysticercosis	ingestion of egg	not present (except in Autoinfection, small intestine)	sub-cutaneous muscles brain, eyes	depending on locality: from none to epilepsy	X-ray, CT, MRI Serology
<i>Hymenolepis nana</i>	hymenolepiasis	ingestion of egg	Small Intestine	Intestinal Villi	Enteritis diarrhoea	eggs in stools
<i>Echinococcus granulosus</i>	hydatid disease	ingestion of egg	not present	Liver, lungs, Bones etc	depending on locality	X-ray, CT, US Serology Hydatid sand




Taenia saginata


Taenia saginata

- Is an obligatory parasite of man, the adult worm live in the **SMALL INTESTINE**
- **CATTLE** become infected by ingesting grass contaminated with **eggs or gravid segments** which passed from human faeces. In the cattle the onchosphere hatches out go to circulation and transformed to cysticercus stage in the muscle known as **CYSTICERCUS BOVIS**
- Man become infected by eating **undercooked** or improperly cooked beef, the adult worm lives in **small intestine** of man passing eggs and gravid proglottids to the environment.
- The majority of cases are Asymptomatic, some patients have vague intestinal discomfort, vomiting and diarrhoea

Life cycle of *Taenia saginata*




Taenia solium



Dwarf tape worms

Hymenolepis nana

Life cycle



Pathology and Clinical features

Often there are none, but with heavy infection there may be abdominal pain and diarrhoea. Anaemia and nervous symptoms, including dizziness and irritability, can occur in children.


Laboratory diagnosis

Eosinophilia may be present. Ova found in faeces.

Distribution

36 million people are infected worldwide.


Hymenolepis nana






Hymenolepis nana

Echinococcus granulosus






Contamination by
food and fingers



Human
intermediate
host

Location of hydatid cyst *Echinococcus granulosus*




Intrasosseous cyst
Spreads along
medulla by budding
outside cyst.
Semisolid; no fibrosis.

Hydatid cyst




Cerebral hydatidosis



protoscoleces (hydatid sand)

(by P.W. Pappas and S.M. Wardrop)



Hydatid cyst

Diagnosis of Hydatid cyst

- Imaging: computed tomography (CT), magnetic resonance imaging (MRI) revealed a cystic swelling with smooth outline.
- Microscopy: hydatid sand
- Serologic tests; to detect specific antibodies

Treatment of Tapeworms

- Intestinal stages: Praziquantel
- Tissue stages (Hydatid, cysticercosis):
 - Depends on clinical condition: Surgical and/or Albendazole