

King Saud University

College of medicine

US of liver and gall stone

DR. MOHAMMED AYESH
CONSULTANT RADIOLOGIST

outline:

- ▶ Introduction to US.
- ▶ Indications of liver and gall bladder US.
- ▶ Normal anatomy and radiological appearance.
- ▶ Pathology of liver and gall bladder.
- ▶ Common pathological cases.

Introduction to US



History OF US

- ▶ Piezoelectricity discovered by the Curies in 1880 using natural quartz.
- ▶ Piezoelectric Effect is the ability of certain materials to generate an electric charge in response to applied mechanical stress.
- ▶ (US) SONAR was first used in 1940's war-time
- ▶ Diagnostic Medical applications in use since late 1950's

Definition:

- ▶ a diagnostic technique in which high-frequency sound waves penetrate the body, bounce around, and produce multiple echoes; these echo patterns can be viewed as an image on a computer screen.
- ▶ Frequency ranges used in medical Ultrasound imaging are 2 - 15 MHz

US machine



MACHINE




PROBES


B- MODE.




M- MODE.



DUPLEX



COLOR DOPPLER



US uses:

- ▶ Cardiology

Echocardiography is an essential tool in cardiology, valvular heart disease.

- ▶ Emergency Medicine:

for Trauma patient and acute abdomen.

- ▶ Gastroenterology:

In abdominal sonography, the solid organs of the abdomen such as the pancreas, aorta, inferior vena cava, liver, gall bladder, bile ducts, kidneys, spleen and appendix.

- ▶ Gynecology:

to assess female pelvic organs, uterus ovaries.

- ▶ Obstetrics:

- ▶ sonography is commonly used during pregnancy to check on the development of the fetus.

Cont.

- ▶ Neurology
- ▶ for assessing blood flow and stenoses in the carotid arteries (Carotid ultrasonography)

- ▶ Neonatology:
for basic assessment of intracerebral structural abnormalities, bleeds, ventriculomegaly or hydrocephalus.

- ▶ Urology:
to study a patient's bladder, prostate or testes.

- ▶ Musculoskeletal
- ▶ For assessing tendons, muscles, nerves, ligaments, soft tissue masses, and bone surfaces

- ▶ vascular system:
- ▶ To assess patency and possible obstruction of arteries Arterial doppler, diagnose DVT venous doppler and determine extent and severity of venous insufficiency

Advantages of US

- ▶ noninvasive
- ▶ inexpensive.
- ▶ Easy and available.
- ▶ Safe and non ionizing.

Disadvantages of US

- ▶ Inability to penetrate gas or bone.
- ▶ Operator dependant.
- ▶ Less sensitive in some situations.



Indications of liver and gall bladder US

- ▶ Right upper quadrant pain.
- ▶ Jaundice.
- ▶ High liver function test.
- ▶ Fever work up.
- ▶ Screening for metastasis.

Normal anatomy and radiological appearance



Cont.



Pathology of the liver:

- ▶ Size.
- ▶ Diffuse liver disease.
- ▶ Focal liver disease.
- ▶ Hepatic vascularity.
- ▶ Biliary system obstruction/pathology.

Size abnormality

- ▶ Normal liver size:
15 cm at MCL.
- ▶ Hepatomegaly:
 - Infective eg viral hepatitis.
 - Neoplastic eg. Metastasis.
 - Degenerative eg. early cirrhosis.
 - Raised venous pressure eg. Congestive cardiac failure.
 - Storage disorder eg. Amyloidosis.
 - Myeloproliferative disorder eg. Polycythaemia rubra vera.




Cont.

- ▶ Small shrunken liver (Late cirrhosis):
- ▶ Shrunken liver with irregular outline
- ▶ Ascitis
- ▶ Portal hypertension.
- ▶ +/- focal lesion.



Diffuse abnormality

- ▶ Diffuse increase parenchymal echogenicity (whiter than normal)
- ▶ Diffuse fatty infiltration.
- ▶ Other infiltrative:
 - Malignant
 - Infectious
 - Glycogen storage disease



Cont.

- ▶ Diffuse decrease in parenchymal echogenicity.

(darker than normal)

- ▶ Acute hepatitis.

- ▶ Other:

Malignant infiltration.




Focal liver lesions

- ❑ Benign tumor:
 - ▶ Hemangioma.
- ❑ Malignant tumor:
 - ▶ Primary eg. Hepatocellular carcinoma.
 - ▶ Secondary metastasis eg. Colon breast.
- ❑ Infective:
 - ▶ Abscess
 - ▶ hydated cyst.
- ❑ Congenital:
 - ▶ Hepatic cyst.

Cont.


Liver abscess




hemangiomas



metastasis



HCC



Cont.



Vascular abnormality



□ Portal venous system:

- ▶ thrombosis.
- ▶ Portal hypertension.

□ Hepatic venous system:

- ▶ Thrombosis
- ▶ (budd chiari syndrome).

Cont.




Hepatic vein thrombosis



PV thrombosis

Biliary abnormality

- ▶ Intra-hepatic biliary radicals.
Less than 3mm
- ▶ Extra-hepatic “CBD”
Less than 8mm
- ▶ Causes of dilatation & obstruction:
 - Intra-luminal:
 - ✓ Stone & mass.
 - Mural:
 - ✓ stricture (benign & malignant)
 - Extrinsic:
 - ✓ Compression mass & Lymph node




Pathology of gall bladder


- ▶ Intra-luminal pathology.
- ▶ Mural pathology.

Intra-luminal pathology

▶ Gall stone:
Acoustic shadowing



▶ Polyps
No acoustic shadowing.




Cont.

- ▶ Intraluminal:

Mass lesion

+ - invasion

Gall bladder carcinoma.



Mural pathology

❑ Mural thickening:

➤ Primary:

Cholecystitis.

➤ Secondary:


✓ Cardiac failure.

✓ Cirrhosis.

✓ ascitis

✓ Hypoalbuminaemia

✓ Renal failure.



Common pathological cases

Case one

- ▶ Middle age women presented to ED with fever, RUQ pain



- ▶ On exam

She looks ill, febrile and on pain

Abdomen: RUQ tenderness

- ▶ Lab high LFTs & WBC.

Cont.




- ▶ Thickening of GB wall $>3\text{mm}$.
- ▶ Distended GB
- ▶ Pericholecystic fluid.
- ▶ Hyperemia.
- ▶ Gall stone

Acute calcular cholecystitis.

Case two

- ▶ Middle age women presented to surgical out patient clinic with 2 years history of recurrent RUQ pain mild to moderate in severity radiated to the right shoulder aggravated by fatty meal.
- ▶ On exam:
obese lady well not distressed, febrile or jaundiced.
- ▶ Lab LFTs normal.

Cont.




- ▶ Multiple oval shaped echogenic structures seen within GB causing acoustic shadowing

GB stones

Case three

- ▶ Middle age man presented to ER with severe RUQ pain and yellowish discoloration of skin and sclera.
- ▶ On exam:
he looks ill, jaundiced and on pain but not febrile
- ▶ Lab high LFTs.

Cont.




- ▶ Dilated intra-hepatic and extra-hepatic biliary system
- ▶ Echogenic structure seen within CBD

CBD stone causing biliary obstruction.

Case Four

- ▶ Old man recently discovered to have colonic cancer presented to primary health care clinic with vague upper abdominal pain
- ▶ On exam:
He was thin & ill but not febrile or jaundiced.
Mild abdominal tenderness enlarged liver with irregular outline.
- ▶ Lab mildly elevated LFTs.

Cont.



- ▶ Multiple hypoechoic focal hepatic lesions

Metastatic liver lesions.

Case Five

▶ Middle age man known case of HCV+ for 10 years presented to GI out patient clinic with history of weight loss, indigestion and mild abdominal pain. No fever.



▶ On exam:

He was ill, slim, mildly jaundice but not febrile.

Abdomen: bulging flanks, dilated tortuous vessels around umbilicus. Mild diffuse abdominal tenderness.

▶ Lab high LFTs.

Cont.




- ▶ Shrunken liver with irregular outline.
- ▶ Heterogeneous appearance.
- ▶ Focal hypoechoic lesion.
- ▶ Cirrhotic liver with HCC.

Case Six

- ▶ Young man known IV drug addict presented to ER with high fever, chills, upper abdominal pain and vomiting
- ▶ On exam:
He looks very ill, febrile and on pain.
Abdomen: RUQ tenderness.
- ▶ Lab high LFTs & WBC.

Cont.



- ▶ Focal hypoechoic liver lesion with ill defined outline.

Liver abscess.



thank you

Radiology:
The Eye of
Medicine