

MJerobjology Praetjeal

GI BLOCK; 2021

Objectives:

- 1. Understand the use of viral serological studies for the diagnosis of hepatitis A, B & C infections.
- 2. To know measures to prevent hepatitis A & B infections.
- 3. To know the viral serological tests used to screen blood donors.
- 4. Risk of transmission of HBV

Case 1

Mohammed Khan is a 20 year-old male who has recently arrived from India to work as a food handler in a restaurant in Riyadh. Three weeks after his arrival, he was seen in A&E Dept. of KKUH because of repeated vomiting, abdominal pain and fever. On examination, his temperature was 38°C, his pulse rate 110/min and BP 120/80mmHg, he was jaundiced and had tenderness in the right upper quadrant of his abdomen.

Case 1

Mohammed Khan is a 20 year-old male who has recently arrived from India to work as a food handler in a restaurant in Riyadh. Three weeks after his arrival he was seen in A&E Dept. of KKUH because of repeated vomiting, abdominal pain and fever. On examination, his temperature was 38°C, his pulse rate 110/min and BP 120/80mmHg, he was jaundiced and had tenderness in the right upper quadrant of his abdomen.

QUESTJONS

1. What are the possible causes for his presentation?

2. What investigations would you like to order for him? Explain how these investigations would help you.

Test	How this investigation will help you?
1. CBC & ESR	
2. Blood Film	
3. Liver function test	
4. Viral Hepatitis screening	
5. Blood Culture	

Test	How this investigation will help you?		
1. CBC & ESR	Shows non-specific signs of infections or inflammation		
2. Blood Film	To exclude malaria		
3. Liver function test	To asses liver function		
4. Viral Hepatitis screening	To exclude viral hepatitis		
5. Blood Culture	To exclude typhoid fever		

Investigation

	CBC		LFTs	
Hb	14.2 g/L	AST	1557 IU/L	(12~37)
WBCs	6100 mm^3	ALT	1879 IU/L	(20~65)
Platelet	271 g/L	ALP	441 IU/L	(175~476)
ESR	4mm/h	Albn	42.3 g/L	(30-50)
Malaria E	Blood film -ve.	Bilirubin	n 86 µmol/	'L (3~17)
Blood cu	lture is negative.			

3. Based on these findings what is the most likely diagnosis?

4. What further investigations would you like to order?

5. The serologic results were as follows:

TEST	RESULT
Anti~HAV~IgM	Positive
HBsAg	Negative
Anti~HCV	Negative
Anti~HEV IgM	Negative

6. Based on the serologic results, what is the diagnosis?

7. Briefly outline the management of this patient.

Case 2

Mohammed Abdullah is a 34 year old married Saudi male who has donated two units of blood at KKUH for a relative undergoing an operation. Two days later, the Blood Bank called him because of abnormal blood test results and advised him to see his physician.

On arrival to the blood bank, the doctor informed him that his blood is not suitable for transfusion because of the presence of infection.

QUESTJONS

1. What type of infectious agents can be transmitted through blood transfusion? (List 4 infections).

The next day Mohammed came to see his general practitioner with a letter from the Blood Bank. The letter revealed the result shown below.

What is your interpretation?

Test	Result
HBsAg	Negative
Anti~HBc	Negative
Anti~HCV	Positive
HIV-Ag/Ab	Negative
Anti~HTLV	Negative

What do you do next?


How would you interpret the results ordered by the GP?

Lab. Test	Patient Result	Normal Range
ALT	49	20~65 IU
AST	29	12~37 IU
Bilirubin	4	3~17 mol/L
HIV-Ag/Ab	Negative	~
Anti~HCV	Positive	~
HBsAg	Negative	~
Anti~HBc	Negative	~
Anti~HBs	Negative	~


4. How do you diagnose HCV infection?

Serological assay

- → Screening for (Anti~HCV) by ELISA
- Confirmatory test by
 recombinant immunoblot assay (RIBA)
 Or Line Immunoassay INNO-LIA HCV



Molecular assay



The General practitioner arrange for him to see hepatologist who examine him and review his results. He further added PCR with genotype for Hepatitis C.

What is the significance of these tests & how they can help in the management:

Test	Significance	How it can help?
1. PCR	1~Qualitative: ~ or + (HCV~RNA) 2~Quantitative: viral load	 Confirm the Dx Monitor response to Rx
2. Genotype	Identify the genotype of HCV	Guide the choice & duration of therapy.

Case 3


A 15-weeks pregnant Saudi woman was seen for the first time at the antenatal clinic at KKUH. As part of the antenatal screening, the doctor arranged for blood screening for viral serology.

The results were as follows:

Test	Result
HBsAg	positive
HBeAg	negative
Anti~HBe	positive
Anti~HBc IgM	negative
Total Anti~HBc	positive
HIV Ag/Ab	negative
Anti~HCV	negative


Acute HBV Infection with Recovery

Typical Serologic Course



Progression to Chronic HBV Infection

Typical Serologic Course



1. How would you interpret these results?

2. On the lights of these Laboratory results how would you manage the newborn?

3. Is there a risk of transmission of HBV to the newborn?

3. Is there a risk of transmission of HBV to the newborn?

4. What further management would you offer to the mother?

Pregnant Hepatitis B carriers should be advised to

- ~ Be seen at least annually by their regular medical doctor.
- Obtain vaccination against hepatitis viruses A as indicated.
- Discuss the risk for transmission with their partner and need for testing.
- > Not donate blood, body organs, other tissue.
- Not share any personal items that may have blood on them (e.g., toothbrushes).

Today the mother is admitted in labour and you were among the staff involved in the delivery.

During a repair of the episiotomy, you accidentally prick your finger with a needle stained by the patient blood.

What should you do?

1. What should you do?

- Report occupational exposures immediately.
- The hepatitis B vaccination status and the vaccine-response status (if known) should be reviewed.
 - > HBsAg
 - > Anti~HBc
 - > Anti~HBs

2. What is the risk of infection to you?

the risk of developing serologic evidence of HBV inf if the blood (+) HBsAg

the risk of developing serologic evidence of HBV inf if the blood (+) HBsAg

(~) HBeAg



23~37%

Tests	RESULTS	INTERPRETATION
HBSAG	NEGATIVE	
ANTI-HBC	NEGATIVE	
ANTI-HBS	NEGATIVE	
HBSAG	NEGATIVE	
ANTI-HBC	POSITIVE	
ANTI-HBS	POSITIVE	
HBSAG	NEGATIVE	
ANTI-HBC	NEGATIVE	
ANTI-HBS	POSITIVE	
HBSAG	POSITIVE	
ANTI-HBC	POSITIVE	
IGM ANTI-HBC	POSITIVE	
ANTI-HBS	NEGATIVE	
HBSAG	POSITIVE	
ANTI-HBC	POSITIVE	
IGM ANTI-HBC	NEGATIVE	
ANTI-HBS	NEGATIVE	
HBSAG	NEGATIVE	
ANTI-HBC	POSITIVE	
ANTI-HBS	NEGATIVE	

Tests	RESULTS	INTERPRETATION
HBSAG	NEGATIVE	
ANTI-HBC	NEGATIVE	SUSCEPTIBLE
ANTI-HBS	NEGATIVE	
HBSAG	NEGATIVE	
ANTI-HBC	POSITIVE	IMMUNE DUE TO
ANTI-HBS	POSITIVE	NATURAL INFECTION
HBSAG	NEGATIVE	IMMUNE DUE TO HEPATITIS
ANTI-HBC	NEGATIVE	B VACCINATION
ANTI-HBS	POSITIVE	
HBSAG	POSITIVE	
ANTI-HBC	POSITIVE	ACUTELY
IGM ANTI-HBC	POSITIVE	INFECTED
ANTI-HBS	NEGATIVE	
HBSAG	POSITIVE	
ANTI-HBC	POSITIVE	CHRONICALLY
IGM ANTI-HBC	NEGATIVE	INFECTED
ANTI-HBS	NEGATIVE	
HBSAG	NEGATIVE	FOUR
ANTI-HBC	POSITIVE	INTERPRETATIONS
ANILURS	NEGATIVE	POSSIBLE *



- 1. May be recovering from acute HBV infection.
- 2. May be distantly immune and test not sensitive enough to detect very low level of anti-HBs in serum.
- 3. May be susceptible with a false positive anti-HBc.
- 4. May be undetectable level of HBsAg present in the serum and the person is actually a carrier.

- عن أبي هريره رضي الله عنه
- أن رسول الله صلي الله عليه وسلم قال:

{ كلمتان خفيفتان على اللسان , ثقيلتان في الميزان حبيبتان إلى الرحمن :

سبحال (اللم وبحسره)

سبحاق (لللم (لعظيم)

