

# Vitamin D, Rickets and Osteoporosis

Prof. Rana Hasanato

# Objectives

By the end of this lecture, the students should be able to:

- Understand the functions, metabolism and regulation of vitamin D
- **Discuss the role of vitamin D in calcium homeostasis**
- Identify the types and causes of rickets
- Identify biomarkers used for the diagnosis and follow up of osteoporosis

# Overview

- Vitamin D distribution, metabolism, regulation and functions
- Vitamin D in calcium homeostasis
- Vitamin D deficiency
- Nutritional and inherited rickets
  - Types, diagnosis and treatment
- Osteoporosis
  - Diagnosis, biomarkers, treatment and prevention

# Vitamin D

- Vitamin D is considered a steroid hormone
- Cholecalciferol (vitamin D<sub>3</sub>) is synthesized in the skin by the sunlight (UV)
- The biologically active form is 1,25-dihydroxycholecalciferol (calcitriol)
- Ergocalciferol (vitamin D<sub>2</sub>) is derived from ergosterol in plants
- D<sub>3</sub>, D<sub>2</sub> are also available as supplement


# Vitamin D distribution

- Dietary sources:
  - Ergocalciferol (vitamin D<sub>2</sub>) found in plants
  - Cholecalciferol (vitamin D<sub>3</sub>) found in animal tissue
- Endogenous vitamin precursor:
  - 7-Dehydrocholesterol is converted to vitamin D<sub>3</sub> in the dermis and epidermis exposed to UV in sunlight


*Daily requirement (IU/day):*

*Adults: 600 Children: 400 Elderly: 800*

# Sources of Vitamin D



# Metabolism and actions of vitamin D



# Vitamin D metabolism

## In skin:

- Cholecalciferol (Vitamin D<sub>3</sub>) is derived from 7-dehydrocholesterol by the sunlight

## In liver:

- Cholecalciferol is converted to 25-hydroxycholecalciferol (calcidiol) by the enzyme 25-hydroxylase



# Vitamin D metabolism


In kidneys:

- The 1- $\alpha$ -hydroxylase enzyme converts 25-hydroxycholecalciferol to 1,25-dihydroxycholecalciferol (biologically active)
- Active vitamin D is transported in blood by gc-globulin protein

# Vitamin D regulation and calcium homeostasis

- Vitamin D synthesis is tightly regulated by plasma levels of phosphate and calcium
- Activity of 1- $\alpha$ -hydroxylase in kidneys is:
  - Directly increased due to low plasma phosphate
  - Indirectly increased via parathyroid hormone (PTH) due to low plasma calcium
  - PTH increases vitamin D synthesis in kidneys
- Vitamin D has essential role in calcium homeostasis
- Calcium homeostasis is maintained by parathyroid hormone (PTH) and calcitonin

# Metabolism and actions of vitamin D




# Vitamin D action


- Vitamin D action is typical of steroid hormones
- It binds to intracellular receptor proteins
- The receptor complex interacts with target DNA in cell nucleus
- This stimulates or represses gene expression

# Vitamin D functions

- Regulates plasma levels of calcium and phosphate
- Promotes intestinal absorption of calcium and phosphate
- Stimulates synthesis of calcium-binding protein for intestinal calcium uptake
- Minimizes loss of calcium by the kidneys
- Mobilizes calcium and phosphate from bone to maintain plasma levels

# Vitamin D response to low plasma calcium





Copyright © Pearson Education, Inc., publishing as Benjamin Cummings.

## Calcium homeostasis

# Vitamin D deficiency


- Deficiency most common worldwide
- High prevalence in Saudi Arabia due to:
  - Low dietary intake
  - Insufficient exposure to Sun
- Circulating level of  $>75$  nmol/L is required for beneficial health effects



# Nutritional rickets

- A disease in children causing net demineralization of bone
  - With continued formation of collagen matrix of bone
  - Incomplete bone mineralization
  - Bones become soft and pliable
  - Causes skeletal deformities including bowed legs
  - Patients have low serum levels of vitamin D
- (Osteomalacia: demineralization of bones in adults)*

# Rickets



# rickets

## Causes

- Vitamin D deficiency because of:
  - Poor nutrition
  - Insufficient exposure to sunlight
  - Renal osteodystrophy (causes decreased synthesis of active vitamin D in kidneys)
  - Hypoparathyroidism (hypocalcemia)

# Inherited rickets

## Vitamin D-dependent rickets (types 1 and 2)

- Rare types of rickets due to genetic disorders
- Causing vitamin D deficiency mainly because of genetic defects in:
  - Vitamin D synthesis
  - Vitamin D receptor (no hormone action)

# Diagnosis and treatment of rickets


- **Measuring serum levels of:**
  - 25-hydroxycholecalciferol
  - PTH
  - Calcium
  - Phosphate
  - Alkaline phosphatase
- **Treatment:**
  - Vitamin D and calcium supplementation

# Osteoporosis

- Reduction in bone mass per unit volume
- Bone matrix composition is normal but it is reduced
- Post-menopausal women lose more bone mass than men (primary osteoporosis)
- Increases fragility of bones
- Increases susceptibility to fractures



(a)



(b)

Fig. 1 **Bone showing (a) normal trabeculae and (b) bone loss in osteoporosis.**



Fig. 2 **Crush fractures of vertebral bodies in a patient with osteoporosis.**




Fig. 3 **Elderly woman with so-called 'Dowager's hump' from collapsed vertebrae due to osteoporosis.**



# Osteoporosis

- Secondary osteoporosis may be caused by:
  - Drugs
  - Immobilization
  - Smoking
  - Alcohol
  - Cushing's syndrome
  - Gonadal failure
  - Hyperthyroidism
  - GI disease

# Diagnosis of osteoporosis

- WHO standard: Serial measurement of bone mineral density
- Biochemical tests (calcium, phosphate, vitamin D) alone cannot diagnose or monitor primary osteoporosis
- The test results overlap in healthy subjects and patients with osteoporosis
- Secondary osteoporosis (due to other causes) can be diagnosed by biochemical tests

# Biomarkers of osteoporosis

## Bone formation markers

- **Osteocalcin**

- Produced by osteoblasts during bone formation
- Involved in bone remodeling process
- Released during bone formation and resorption (bone turnover)
- Short half-life of few minutes

# Biomarkers of osteoporosis

- **Bone-specific alkaline phosphatase**
  - Present in osteoblast plasma membranes
  - Helps osteoblasts in bone formation
  - A Non-specific marker
  - Its isoenzymes are widely distributed in other tissues

# Biomarkers of osteoporosis

- P1NP (Procollagen type-1 amino-terminal propeptide)
  - Produced by osteoblasts
  - Involved in the process of type 1 collagen formation
  - Blood levels are highly responsive to osteoporosis progression and treatment

# Biomarkers of osteoporosis

## Bone resorption markers

- CTX-1 (Carboxy-terminal cross-linked telopeptides of type 1 collagen)
  - A component of type-1 collagen
  - Released from type-1 collagen during bone resorption
  - Blood and urine levels are highly responsive to post-resorptive treatment
  - Levels vary largely by circadian variation

# Treatment and prevention of osteoporosis

## Treatment

- In confirmed cases of osteoporosis
  - Treatment options are unsatisfactory
- Oral calcium, estrogens, fluoride therapy may be beneficial
- Bisphosphonates inhibit bone resorption that slows down bone loss

# Treatment and prevention of osteoporosis

## Prevention

- Prevention from childhood is important
- Good diet and exercise prevent osteoporosis later
- Hormone replacement therapy in menopause may prevent osteoporosis



# References

- Lippincott's Biochemistry 6<sup>th</sup> Edition
- Clinical Biochemistry: An illustrated colour text 4<sup>th</sup> Edition by Allan Gaw (Churchill Livingstone)
- Wheater, G. et al. The clinical utility of bone marker measurements in osteoporosis. *J. Trans. Med.* 2013, 11: 201-214.