

Endocrine Block

Glucose Homeostasis

Dr. Usman Ghani

Objectives

- Define glucose homeostasis and the metabolic processes involved
- Differentiate between different phases of glucose homeostasis
- Discuss the primary sources of energy and major organs utilizing glucose during the five phases of homeostasis
- Understand the role of hormones in maintaining glucose homeostasis

Overview

- Introduction
- Sources of glucose
- Phases of glucose homeostasis
- Hormones in glucose homeostasis (actions, role in CHO metabolism)
 - Insulin
 - Glucagon
 - Cortisol
 - Growth hormone
 - Epinephrine

Glucose homeostasis

- A process that
 - Controls glucose metabolism and
 - Maintains normal blood glucose level in the body
- Glucose is a major source of body's energy
- The liver plays a key role in maintaining blood glucose level
- It is tightly controlled as the brain constantly needs glucose
- Severe hypoglycemia can cause coma and death
- Chronic hyperglycemia results in glycation of proteins, endothelial dysfunction and diabetes mellitus

Sources of glucose

Dietary sources:

- Dietary CHO's are digested to monosaccharides
- Starch provides glucose directly
- Fructose and galactose are converted to glucose in the liver

Metabolic sources (via gluconeogenesis):

- Glycerol, lactate, pyruvate, glucogenic amino acids





Fig. 1 **Glucose homeostasis.**

Phases of glucose homeostasis

- Five phases:
 - Phase I (Well-fed state)
 - Phase II (Glycogenolysis)
 - Phase III (Gluconeogenesis)
 - Phase IV (Glucose, ketone bodies (KB) oxidation)
 - Phase V (Fatty acid (FA), KB oxidation)



<u>Phase</u>	<u>ORIGIN OF BLOOD GLUCOSE</u>	<u>TISSUES USING GLUCOSE</u>	<u>MAJOR FUEL OF BRAIN</u>
I	Exogenous	All	Glucose
.....			
II	Glycogen Hepatic gluconeogenesis	All except liver. Muscle and adipose tissue at diminished rates	Glucose
.....			
III	Hepatic gluconeogenesis Glycogen	All except liver. Muscle and adipose tissue at rates intermediate between II and IV	Glucose
.....			
IV	Gluconeogenesis, hepatic and renal	Brain, RBCs, renal medulla. Small amount by muscle	Glucose, ketone bodies
.....			
V	Gluconeogenesis, hepatic and renal	Brain at a diminished rate, RBCs, renal medulla	Ketone bodies, glucose

Phase I (Well-fed state)

- Glucose is mainly supplied by dietary CHO
- Liver removes about 70% of glucose load after a CHO meal
- All body tissues use dietary glucose for energy in this phase
- Some glucose is converted to glycogen for storage in the liver (glycogenesis)

Phase I (Well-fed state)

- Excess glucose is converted to fatty acids and triglycerides in the liver
- These are transported via VLDL (very low density lipoproteins) to adipose tissue for storage
- Gluconeogenesis is inhibited in this phase
 - Cori and glucose-alanine cycles are inhibited

Phase II (Glycogenolysis)

- Phase II starts during early fasting when dietary glucose supply is exhausted
- Hepatic glycogenolysis and gluconeogenesis maintain blood glucose level in this phase
- Major sources of blood glucose in this phase:
 - Glycogenolysis and gluconeogenesis




Figure 22.3. Metabolic interrelationships of major tissues in early fasting state.

Phase III (Gluconeogenesis)

- Phase III starts when glycogen stores in liver are exhausted (< 20 hours)
- Duration of phase III depends on
 - Feeding status
 - Hepatic glycogen stores
 - Physical activity
- Hepatic gluconeogenesis from lactate, pyruvate, glycerol and alanine maintains blood glucose level
- Major source of blood glucose in this phase:
 - Gluconeogenesis

Phase IV (Glucose and KB oxidation)

- Several days of fasting leads to phase IV
- Gluconeogenesis starts to decrease
- FA oxidation increases KB accumulation
- KBs enter the brain and muscle for energy production
- Brain uses both glucose and KB for energy

Phase V (FA and KB oxidation)

- Prolonged fasting leads to phase V
- Less dependence on gluconeogenesis
- All body tissues mainly use FA and KB oxidation for energy production
- Gluconeogenesis somewhat maintains blood glucose level in this phase




Figure 22.4. Metabolic interrelationships of major tissues in fasting state.

Phase V (FA and KB oxidation)

- High KB conc. and glucose levels inhibit proteolysis in muscle (conservation of muscle)
- When all fat and KBs are used up
 - Body uses muscle protein to maintain blood glucose level

Hormones and glucose homeostasis

Hormones that regulate glucose metabolism:

- Insulin (lowers blood glucose level)

- Glucagon

- Cortisol

- Growth hormone

- Epinephrine



Antagonize insulin action

Insulin

- Plays a major role in glucose homeostasis
- Synthesized by the β -cells of islets of Langerhans of pancreas
- A small protein composed of two chains
- Rise in blood glucose level stimulates insulin secretion
- Promotes entry of glucose into cells

Insulin actions






Fig. 2 The actions of insulin.

Mechanism of action

- The insulin receptor is present on the plasma membrane of cell
- Composed of
 - α -subunit (extracellular)
 - β -subunit (cytoplasmic)
- Binding of insulin to α -subunit causes phosphorylation of β -subunit
- This activates the receptor
- The activated receptor then phosphorylates intracellular proteins generating a biological response





Insulin and CHO metabolism

Promotes glucose uptake into cell:


- Glucose is diffused into cells through hexose transporters such as GLUT4
- GLUT4 is present in cytoplasmic vesicles
- Insulin binding to its receptor causes vesicles to diffuse into plasma membrane
- GLUT4 is inserted into the membrane
- Allowing glucose transport into the cell
- Brain and liver have non-insulin dependent glucose transporter

Insulin and CHO metabolism

- Stimulates glycogen synthesis
- Decreases blood glucose levels
- Increases glycolysis
- Stimulates protein synthesis
- Insulin deficiency causes diabetes mellitus
- Hyperinsulinemia is due to insulin resistance in:
 - Diabetes mellitus or
 - Metabolic syndrome

Glucagon

- A peptide hormone secreted by α -cells of pancreatic islets
- Secreted in response to hypoglycemia
- Increases glucose levels
- Stimulates glycogenolysis
- Activates hepatic gluconeogenesis



Glucocorticoids (Cortisol)

- Cortisol is a steroid hormone secreted by adrenal gland
- Contributes to glucose homeostasis
- Maintains normal glucose levels in fasting
- Stimulates gluconeogenesis in the liver
- Mobilizes amino acids for gluconeogenesis
- Stimulates fat breakdown in adipose tissue

Growth hormone

- A protein hormone secreted by anterior pituitary gland
- Maintains blood glucose levels by:
 - Inhibiting insulin action
 - Stimulating gluconeogenesis in the liver

Epinephrine

- A catecholamine hormone secreted by adrenal gland
- Stimulates lipolysis in adipose tissue when blood glucose levels fall
- Promotes glycogenolysis in skeletal muscle

Take home message

- Glucose homeostasis is a process that controls glucose metabolism and maintains blood glucose level in the body
- There are five phases of glucose homeostasis- Phase I (well-fed state), Phase II (glycogenolysis),
- Phase III (gluconeogenesis), Phase IV (glucose, ketone bodies (KB) oxidation), Phase V (fatty acid (FA), KB oxidation)
- Hormones that regulate glucose metabolism include insulin (lowers glucose level) and glucagon (increases glucose level)
- Other hormone such as cortisol, growth hormone and adrenaline are known to antagonize the actions of insulin thus increases the blood glucose level

References

- Textbook of Biochemistry with Clinical Correlations by Devlin 7th Edition.
- http://www.vivo.colostate.edu/hbooks/pathphys/endocrine/pancreas/insulin_phys.html