

1. Sagittal sutures closes at :
 - a. Twenty years
 - b. Thirty years
 - c. Forty years
 - d. Fifty years .

2. Presence of canine in a child indicates that the child is :
 - a. Below 6 months
 - b. Above 6 months and below 12 months
 - c. Above 12 months and below 18 months
 - d. Above 18 months

3. First permanent tooth appears at age of :
 - a. 6 months
 - b. 6 years

4. Age of 16 years old female is estimated by X-ray on:
 - a. Ankle joint
 - b. Humerus
 - c. Wrist joint
 - d. Elbow joint

5. Fusion of all parts of femur in males complete at age of :
 - a. 18 years
 - b. 21 years

6. Red color hypostasis is seen in death from :
 - a. CO poisoning

7. The age of discrimination is :
 - a. Six years
 - b. Seven years

8. Rigor mortis first appears in the muscles of :
 - a. Face
 - b. Trunk
 - c. Upper limb
 - d. Lower limb

9. In death from hanging , the hypostasis will appear in :
 - a. Face and abdomen
 - b. Back
 - c. Lower limb

10. Mummification occurs when the climate is :
 - a. Dry and hot
 - b. Dry and cold
 - c. Moist and cold
 - d. Moist and hot

11. Hypostasis is :
 - a. Postmortem hemolysis
 - b. Extravasation of blood
 - c. PM Gravitation of blood
 - d. Subcutaneous hemorrhage
12. Sure signs of death start to appear :
 - a. At the moment of death
 - b. After 1 hour
 - c. After 2 hours
13. Eruption of permanent teeth occurs between :
 - a. 6 -15 years.
14. X-ray for 21 female will show fusion of
 - a. Iliac crest
 - b. Iliac tuberosity
15. Rigor mortis is due to :
 - a. Depletion of ATP
 - b. Loss of muscle tone
 - c. Loss of reflexes
16. Primary flaccidity is due to :
 - a. Loss of muscle tone
 - b. Depletion of ATP
17. A cold stiff body:
 - a. 8 -12 Hours
18. Healing by color change occurs in :
 - a. Abrasion
 - b. Contusion
 - c. Contused wound
19. A cut wound may look like contused wound in :
 - a. Forehead
 - b. Over the chin of the tibia
 - c. Axilla
 - d. Forehead
 - e. Chest wall
20. Cadaveric spasm is due to :
 - a. Ante-mortem tension
21. The weakest point in iris print is :
 - a. Fear
 - b. Cataract
 - c. Availability of the device
 - d. No large databases

- 22.** The shape of electrical burns depends on :
- Type of current
 - Surface area
 - Pathway of the current
- 23.** Foot prints are most commonly used for:
- Trailing the enemies in desert.
 - None mixing of infants in Hospitals.
 - Solving of crimes.
- 24.** PM Identification is difficult in cases of:
- severe burns.
 - severe putrefaction.
 - multiple stab wounds.
 - complete charring.
 - absence of the head (decapitation)
- 25.** Which instrument :
- Knife
 - Hammer
- 26.** Permanent infirmity means :
- Loss of more than organ
 - Loss of organ or its function
- 27.** Treatment of heat cramps :
- IV fluid
 - Increase salt in diet
- 28.** Heat collapse affected by :
- Age
 - Sex
 - Health
- 29.** Bridging of the tissue :
- Contused wound
 - Cut wound
- 30.** A person has heat cramps he will have :
- Subnormal temperature
 - Normal temperature
 - Fever
 - Hyperpyrexia
- 31.** Which of the following is not a characteristic of abrasions :
- Hemorrhage
 - Pressure
 - Embolism

- 32.** Reliable evidence of sex :
- Ovaries and testes
 - External genitalia
 - Sex hormones
 - Hair distribution
- 33.** Trans-sexuality means :
- Conflict between gender and physical sex
 - Male hermaphrodite
 - Female hermaphrodite
 - None of the above
- 34.** Blue lines on the gum indicates poisoning of :
- Lead
- 35.** Complete identification means identification of :
- Sex and age
 - Individuality
- 36.** ML classification of wound is :
- Anatomical classification
 - Depends on the shape of the wound
 - Depends on the type of instrument
- 37.** Adepocere occurs after submersion of body in water :
- One week
 - Two weeks
 - Three weeks
- 38.** If the knife penetrate the heart :
- Stab punctured wound
 - Stab penetrating wound
- 39.** Bone fracture :
- Fat embolism
- 40.** Heat collapse :
- Subnormal temperature
 - Slow pulse
- 41.** The mechanism in death from cold is :
- Inhibition of respiratory enzyme
- 42.** Injury from ionizing radiation causes:
- Neutropenia
 - Thrombocytosis
 - Reticulocytosis

43. Red scar in burns may stay up to :
- 2 months
 - 6 months
44. Yellow color in abrasions is due to :
- Biliverdin in the 5th day
 - Bilirubin in the 5th day

باقي كم سؤال ماقدرنا نتذكره اعذرونا



موفقين إن شاء الله ,

Vol & MerMaiD