


Thyroid and Parathyroid Scans

Basic Principles and Clinical Applications



Saleh Othman , MD

A. Prof.& Consultant Nuclear Medicine

King Khalid University Hospital & School Of Medicine

King Saud University

Date: To be announced

Thyroid Scan

LEARNING OBJECTIVES...



At the end of the lecture you will be able to answer the following questions:

- How is the thyroid scan performed?
- When is thyroid scanning helpful?
- What is significant about whether a nodule is "hot" or "cold"?
- What is the role of nuclear medicine in the treatment of thyroid disorders?

How is the thyroid scan performed?



Thyroid Scan : Procedure



Tc-99m Pertechnetate

I-123

Dose	0.5-4.0 mCi given IV	0.5 mCi orally
Half Life	6 Hours	13 Hours
Cost	Not Expensive (Generator)	Expensive (Cyclotron)
Time of imaging	20 min post injection	6 and 24 hours post ingestion
Remarks	Trapped not organified	Trapped and organified

Thyroid Scan : Procedure



Patient Preparation:

The patient must be off thyroid hormones :

1. Thyroxine (T-4) for at least 3-4 weeks.
2. Triiodothyronine (T-3) for at least 10 days.

The patient must not be taking antithyroid medications:

1. Propylthiouracil (PTU) and tapazole for at least 3-5 days.

The patient must not i.v iodinated contrast agents

(IVP, CT with contrast, myelogram, angiogram) for at least 3 weeks.

Radiopharmaceutical and dose:

Tc-99m as sodium pertechnetate 0.5 -4.0 mCi given Intravenously. OR

I-123 Sodium Iodide 0.5 mCi orally

Gamma camera: Small or large field of view.

Patient position: Supine with chin tilted up.


Imaging:

20 min. post injection of Tc99 m : ANT, LAO and RAO images obtained.

6 and 24 hours post oral dose for I123 : ANT, LAO and RAO images .



Normal Thyroid Scan



THYROID UPTAKE MEASUREMENT (I-123 Sodium Iodide)



- The Thyroid Uptake Measurement measures the metabolic activity of the thyroid gland as reflected by its extraction of iodine from the blood.

Indications

- Diagnosis of Grave's disease .
- Evaluation of subacute and chronic thyroiditis .
- Thyroid Cancer

Patient Preparation :

- **Must be off thyroid hormones :**

1. Thyroxine (T-4) for at least 3-4 weeks.
2. Triiodothyronine (T-3) for at least 10 days.

- **Must not be taking antithyroid medications :**

1. Propylthiouracil (PTU) and tapazole for at least 3-5 days.

- **Must not have had intravenous or intrathecal iodinated contrast material (IVP, CT with contrast, myelogram, angiogram) for at least 3 weeks .**

- **Other agents may interfere, but usually only to a small extent .**

- **NPO 2-4 hours before and for at least 1 hour after ingesting the radiopharmaceutical**

THYROID UPTAKE MEASUREMENT WITH /WITHOUT IMAGING (I-123 Sodium Iodide)



Equipment :

Uptake only : Uptake probe (single crystal probe with flat field collimator).

Imaging plus uptake studies: Gamma camera

Radiopharmaceutical : dose given orally

Imaging plus uptake studies: I-123: 500 μCi .

Uptake study only:
I-123: 100 μCi

Thyroid uptake measurements may be determined using Tc-99m-pertechnetate.

Patient position: Sitting.

Detector field of view: Neck.



THYROID UPTAKE MEASUREMENT WITHOUT IMAGING (I-123 Sodium Iodide)



Acquisition Protocol:

- Place I123 capsule(s) in neck phantom.
- Acquire counts for 1 minute , record the counts, time of acquisition, and time of day on the thyroid Uptake Worksheet.
- Immediately administer the capsule(s) to the patient.
- At 6 hours position the probe in front of the patient's neck .
- Acquire counts for 1 minute for I-123 and record the counts, time of acquisition, and time of day on the Worksheet.
- Position the probe over the thigh for 6 hour “background” measurement.
- Acquire counts for 1 minute for I-123 and for 2 minutes for I-131; record the counts, time of acquisition, and time of day on the Worksheet.
- Using the Thyroid Uptake Worksheet, calculate the 6 hour thyroid uptakes. Remember to correct the standard counts for decay.

Twenty four hour uptake measurement in the same way as the 6 hours.

Normal Values Of Thyroid Uptake



- I131 OR I-123 RAIU (4 & 24 hours) :

Normal 4 hour RAIU : 5 - 15%

Normal 24 hour RAIU : 8 - 35%

- Tc- 99m Uptake (20 min Uptake) : N (0.5 -4 .0%)

Causes of High Thyroid Uptake



- **Hyperthyroidism : Grave's Disease or TSH-secreting pituitary adenoma**
- **Autonomous toxic nodule**
- **Multinodular toxic goiter (Plumer's Disease)**
- **Enzyme defects : Dyshormonogenesis.**
- **Iodine starvation (Iodine deficiency)**
- **Lithium Therapy**
- **Recovery phase of thyroiditis.**
- **Rebound following abrupt withdrawal of antithyroid meds**



Causes of Low Thyroid Uptake

- **Parenchymal Destruction:**
 - Acute, Subacute and Chronic Lymphocytic Thyroiditis
- **Hypothyroidism:**
 - Primary or Secondary (insufficient pituitary TSH secretion)
 - Surgical/Radioiodine Ablation of Thyroid
- **Blocked Trapping:**
 - Iodine load (most common): Iodinated contrast material, Food rich in iodide: fish , cabbage ,...etc
 - Exogenous thyroid hormone replacement depressing TSH levels (thyrotoxicosis factitia)
 - Ectopic thyroid: Struma Ovarii
- **Blocked Organification:**
 - Antithyroid medication (PTU): Note- Tc-99m uptake should not be affected

Tc-99m Thyroid scan and uptake


Imaging plus uptake studies



ALYAMI EBTISAM HUSS

843223

Study Date
Study Time



THYROID UPTAKE : 2.96 %

Area	29.8	(sqcm)
Mass	53.0	g



Patient Name : ALYAMI EBTISAM HUSS

Patient ID : 843223

Exam Date : 03Jun2007

THYROID UPTAKE

Adac Laboratories BV
Maarsse
The Netherlands

ANTERIOR

THYROID METASTASES STUDY (I-123 or I-131 as Sodium Iodide)



Indications

- Detection and localization of persistent or recurrent functioning thyroid cancer

Patient Preparation

- Stimulation of potentially functioning thyroid tissue:

A. Inject recombinant human thyrotropin on 2 consecutive days and administer the radiopharmaceutical on the third day .

B. Withdraw thyroid replacement hormones :

1. Thyroxine (T-4) for at least 4 weeks.
2. Triiodothyronine (T-3) for at least 10 days.


- The patient must not have had i.v iodinated contrast material (IVP, CT with contrast, myelogram, angiogram) for at least 3 weeks .

- The patient should be NPO for at least 4 hours prior to radiopharmaceutical administration and for at least 1 hour afterwards .

Radiopharmaceutical, Dose, & Technique of Administration

- Radiopharmaceutical: Oral administration
 - a. I-123 as sodium iodide : 2 mCi
 - b. I-131 as sodium iodide : 2-10 mCi






Imaging using Gamma camera : Whole body scan



Negative WBS

THYROID METASTASES STUDY

(I-123 or I-131 as Sodium Iodide)



Local Recurrence

Bone Metastases

Lung Metastases

When is thyroid scanning helpful?



Indications for Thyroid Scan



- Evaluation of thyroid nodules : No. & type
- Evaluation of congenital hypothyroidism : Agenesis Vs. Dyshormonogenesis.
- Evaluation of neck masses : ectopic thyroid, thyroglobulin cyst.
- Evaluation of thyrotoxicosis.

Evaluation of thyroid nodules

Single vs MNG





Solitary cold nodule

Multinodular goiter

The chance of malignancy is more in Solitary cold nodule than in MNG

Evaluation of thyroid nodules

Hot vs Cold vs warm



Hot

< 5% Malignant

Cold


15-20% Malignant

warm


Suspicious

Evaluation of congenital hypothyroidism


Agnesis vs Dyshormonogenesis



Agnesis



Dyshormonogenesis





Perchlorate Discharge Test


- 50 - 80 uCi I^{123} orally.
- 2 hrs RAIU
- 400 mg Kclo4
- RAIU/ 15 min for 2 hrs.
- Positive test : ≥ 15 fall of RAIU below 2 hrs. uptake.

Evaluation of neck masses

ectopic thyroid vs thyroglossal cyst



Lingual thyroid




Thyroglossal cyst

Evaluation of thyrotoxicosis


Thyrotoxicosis with hyperthyroidism



- Graves' Disease .
- Neonatal hyperthyroidism.
- Toxic nodular goiter :
 - MNTG or Plummer's disease
 - ATN or toxic adenoma
- Iodine induced (Jod-Basedow disease)
- Rare causes :
 - Excessive HCG by trophoblastic tumor
 - Hypothalamic pituitary neoplasms (TSH induced)



Graves disease



MNTG

Evaluation of thyrotoxicosis


Thyrotoxicosis **with** hyperthyroidism



- **Graves' Disease .**
- **Neonatal hyperthyroidism.**
- **Toxic nodular goiter :**
 - MNTG or Plummer's disease
 - ATN or toxic adenoma
- **Iodine induced (Jod-Basedow disease)**
- **Rare causes :**

Excessive HCG by trophoblastic tumor

Hypothalamic pituitary neoplasms (TSH induced)




ATN

Evaluation of thyrotoxicosis

Thyrotoxicosis **without** hyperthyroidism



- **Subacute thyroiditis.**
- **Chronic thyroiditis with transient thyrotoxicosis**
- **Thyrotoxicosis factitia** (exogenous hormone).
- **Thyroid extract** (e.g.Hamburger thyrotoxicosis)
- **Ectopic thyroid :**
Metastatic thyroid carcinoma
Struma ovari



SAT

Radioactive Iodine Therapy for Hyperthyroidism



- **Isotope used : I131**
- **Physical Properties:** Solution or capsule
- **Main side effect : Hypothyroidism**
- **Dose :**
 - a. **Calculated : Considering weight and uptake of the gland**
 - b. **Empirical :**
 - Graves: 5-15 mCi**
 - ATN : 15-20 mCi**

Radioactive Iodine Therapy for Thyroid Cancer



Isotope used : I131

Physical Properties: Solution or capsule

- **Thyroid remnant : 80-100 mCi**
- **Lymph Node Mets : 100 mCi**
- **Local Recurrence : 100 mCi**
- **Lung Mets : 150 mCi**
- **Bone Mets : 200 mCi**

Parathyroid Scan





Figure 13.10. Normal and aberrant distribution of the parathyroid glands.

Parathyroid Scan

LEARNING OBJECTIVES...



At the end of the lecture you will be able to answer the following questions:

- Discuss the principles of dual-phase and dual-isotope parathyroid imaging.
- Identify the common imaging features of pathologic parathyroid glands.
- Discuss causes of false negative scans.
- Identify causes of false negative and false positive scans.

Parathyroid Scan Techniques



- TL-201 _ Tc-99m subtraction
- Tc-99m Sestamibi (Dual Phase)
- Tc-99m Tetrofosmin (Dual Phase)

Parathyroid imaging



Radiopharmaceutical

99mTc / 201Tl Subtraction

99mTc sestamibi

Activity administered

80 MBq (2 mCi) 201Tl;
370 MBq (10 mCi) 99mTc

925 MBq (25 mCi)

Images acquired

Inject Tl .rst and acquire 15-min
100 000 count view of neck and
mediastinum.

Then acquire similar Tc images
without moving patient.

Subtract Tc data from Tl after
normalization to equal count densities

Anterior (and oblique)
views at 15 min and
at 2–3 h; SPECT as needed

PARATHYROID IMAGING



Tc-99m-Sestamibi

- The Parathyroid Study depicts hypertrophied parathyroid tissue, probably because of uptake of Tc-99m-sestamibi in the mitochondria of hyperactive cells.

Indications : Detect and localize parathyroid adenomas .

Patient Preparation :None.

Radiopharmaceutical, Dose, & Technique of Administration

- **Radiopharmaceutical:** 25 mCi Tc-99m-sestamibi i.v.
- **Patient position:** Supine with head and neck extended and immobilized.
- **Gamma camera Imaging field:**
 1. Neck.
 2. Upper two thirds of the mediastinum.

Acquire images at 15 minutes and 2-3 hours post injection.

SPECT images improves localization.

- **TI – Tc99m subtraction :** Several protocols have been developed for routine subtraction of thyroid tissue from parathyroid tissue

Parathyroid Scan


Dual phase MIBI Scan (Or Tetrofosmin)



MOHAMMED NAJI


538527

22Apr2002



Parathyroid Scan

Tc-Tl Subtraction Scan



Parathyroid Scan


Sestamibi dual phase

5MIN EARLY F_SS

Se:1
Im:2 (F1/1)

ALSOESSE FATMA AHMED MATAR
Study Date:15/04/1428
Study Time:11:03:33
MFI:740079
02May2007

ALSOESSE FATMA AHME 740079



5MIN EARLY FULL



EARLY ZOOM

PARATHYROID SCAN

TC-MIBI

KKUH MA

Parathyroid Scan

Ectopic Parathyroid



IMG: (F171)



5MIN A FULL



5MIN A FULL


PARATHYROID MIBI

27Jun2007

Study Date:12/06/1426
Study Time:12:49:29
MRN:831769



EARLY ZOOM



DELAY ZOOM

KKUH


ROOM4

RSM

C80
W160

PLANAR vs SPECT/CT




KK16289



5MIN A FULL



2HRS DBL ZOOM





SPECT-CT images accurately localize the adenoma and guide the surgeon to the best surgical approach

Sestamibi Parathyroid Scan Result

High PTH / High Ca

High PTH / High Ca



TP


FN

**Q:What is the cause of the
FN result...?**



A: Mechanism of sestamibi uptake

Parathyroid Cells



Normal parathyroid glands comprise 2 cell types:

- i. Chief cells:** responsible for PTH production
- ii. Oxyphil cells:** eosinophilic cells whose cytoplasm is composed almost entirely of mitochondria.

While the normal oxyphil cell does not synthesize and secrete PTH, the oxyphil cells of pathologic parathyroid glands do secrete the hormone.



Mechanism of Sestamibi uptake

SESTAMIBI : METHOXYISOBUTYLISONITRILE

- Its parathyroid uptake was first reported by Coakley et al. in 1989
- Mechanism of MIBI uptake and retention is still unclear. Multifactors have been proposed:

a. Biochemical properties of the tracer :


Lipophilicity : The lipophilic sestamibi molecule is concentrated by mitochondria. This explains why adenomas with an abundance of mitochondrial-rich oxyphil cells retain the sestamibi

Cationic charge


b. Cell Type : A predominance of oxyphil cells within an adenoma is more likely to lead to a positive scan.

c. Local factors: blood flow, trans-capillary exchange, interstitial transport and negative intracellular charge of both mitochondria and membranes.

Cell Type and Scan Result



Parathyroid adenoma composed entirely of glycogen-rich chief cells.



Parathyroid adenoma composed mainly of mitochondrial-rich oxyphil cells.

Q



**What is the cause of the
FN result...?**



“FN” Sestamibi Scan...?

- **Histologic type** : False-negative scans can occur with parathyroid glands containing predominantly clear cells.
- **Size and Location**: Smaller-volume parathyroid adenomas and those in the upper position are less likely to be localized with sestamibi scans.
- **Number of adenomas**: FN rate is increased with MGD compared with patients with a single adenoma .
- **Decreased tracer concentration** : Possible association.
 - a. P-glycoprotein expression
 - b. Multidrug resistance–related protein expression
- **Variability of radiotracer uptake in parathyroid adenomas**: Related to differences in perfusion and metabolic activity

Even with refinements in sestamibi scanning, the fact that all parathyroid adenomas are not created equal on a cellular level may inevitably lead to FN scans in a certain number of cases.



False-Positive Scintigraphic Findings

- Solitary thyroid adenoma or a multinodular goiter
 - Benign or malignant tumors : breast, lung, and head and neck carcinomas and their lymph node and osseous metastases, as well as bronchial carcinoids.
 - Primary thyroid lymphomas.
 - Cervical L.N. metastasis from PTC carcinoma
 - Reactive lymph nodes
 - Remnant thymus
 - PTH-secreting paraganglioma
 - Enlarged submandibular salivary gland .
- *In the clinical setting of hyperparathyroidism, false-positive findings are uncommon.*

Points To Remember Before Proceeding For Parathyroid Imaging



- **Imaging is not for diagnosis:** High Ca and PTH establish the diagnosis
- **Imaging does not identify normal parathyroids:** These are too small to be seen (20-30mg)
- **Imaging should detect abnormal parathyroid(s) and indicate the approximate size and the precise relationship to the thyroid gland:** lateral , SPECT and SPECT /CT
- **Imaging should identify ectopic glands :** SPECT and SPECT/CT
- **Optimal imaging should be able to differentiate patients with single adenoma from those with MGD**
- **Imaging should identify thyroid nodules which may require concurrent surgical resection.**

Reference book and the relevant page numbers..



- **Handout Nuclear Medicine**

Dr. Saleh Othman -kkuh

- **Nuclear Medicine: The Requisites, Third Edition (Requisites in Radiology) [Hardcover]**



Harvey A. Ziessman MD, Janis P. O'Malley MD, James H. Thrall MD

Relevant Pages :

I- Thyroid and Parathyroid : 71-105

**II- Oncology : 264-274 , 279 -283 ,302 -345 ,119-133
109 -112 ,296 -299**

Nuclear Medicine
Basic Principles And Clinical Applications
Dr. Saleh Othman



Thank You 😊



Radiology (Nuclear Medicine)

Dr. Saleh Othman

XX-XX-XXXX