



# RADIOLOGY

TEAM 435

## Radiology of the cardiac disease

[ Color index: **Important** | **Notes** | Extra ]

[ [Editing file](#) | [Feedback](#) | [Share your notes](#) | [Shared notes](#) ]

### ● Resources:

- 435 Slides
- 434 Team
- 435 Notes

### ● Done by:


- Naif Alsolami
- Mai Alageel

### ● Team Leaders:

- Mohammed Alsahil
- Amjad Alduhaish


### ● revised by:

- Ahmad alyahya



# Cardiac CT Angiogram

- When do we need to do Chest CT? when we have query of cardiac diseases.
- **Examination for cardiac diseases is always done with I.V contrast.**
- Advantages: Safe, fast (takes 5-10 minutes), and more accurate diagnoses.
- Cardiac CT used commonly to diagnose **acute pulmonary embolism**, In the past they used angiography.
- In the past they used CXR to diagnose APE and in case of strong clinical suspicion they would do catheter angiography: incise catheter through lower limb of patient and inject contrast in the pulmonary vessels, patient can develop cardiac arrest "20-30%" during examination and mortality rate in angiography 5-7%
- **Now they use indirect CT** "injecting IV contrast through antecubital vein then they do CT" very safe and contrast injection is very fast + diagnosis is more accurate.





Quick recall! **important!!**

- There are two settings:

mediastinal window "structures"

lung window "with it you'd able to see lungs clearly and the cardiac diseases' effect on the lungs"

# Radiological Anatomy of the Chest




- The most important thing when we see a patient suspected to have pulmonary embolism is “to determine the level”
- Scanogram: is like CXR, this is examination at level of the hilar of the lungs ” it’s the most important level”

- **Mercedes area represents:**




“it is one of important things you need to observe in the CT”

- 1- pulmonary trunk “ main pulmonary artery”
- 2- Left pulmonary artery
- 3- Right pulmonary artery.


- The presence of embolus in these vessels is known as massive embolus because this is major place, If it occurs peripherally it will produce less effect and less severe presentation.
- I.V contrast in optimal CT will show the vessels **homogenous white color** but in case of **embolus** or pulmonary thrombus it will be fully defected & **appears grey or black interrupting this homogenous contrast**
- shows the corresponding anatomy of CT “image in blue”
- this picture is not important



# Pulmonary Artery ( Just for looking )




# Pulmonary Embolism



The most common finding on x-ray in PE is a normal CXR.

**Pic1**

**Pic2**



**Pic3**

★**The gold standard to diagnose acute pulmonary embolism is CT angiography of chest.**

“used to be X-Ray angiography” .

**But** in Emergency the first thing done for patients with chest problems is CXR. It’s **not diagnostic for Acute PE** “non specific signs”; in fact, it is often **NORMAL**.

**Since we know that it will be normal on chest x-ray so why do we do it ?!**

- If the CXR is **normal** and there’s high clinical suspicion and the history is supporting the theory we proceed to CT angiography of chest. **”to exclude PE”**
- If the patient is having PE and we rely only on normal chest X ray (**pic1**) and discharge the patient, he may come after about 24 hours with “massive pulmonary infarction” (**Pic2**).
- So CXR is done to rule out other causes of symptoms: “Pneumothorax, CHF, rib fracture”.


We mentioned before normally blood vessels with contrast would appear homogeneously enhanced “white”. in this image (**Pic3**) we can see that

there is part not well enhanced and that would be considered blood clot or thrombus or **“acute pulmonary embolism”**.


We can diagnose PE by taking a good history and see the patient symptoms.

**Hx:** DVT, Bed ridden, long standing, pregnancy, OCP “oral contraceptive”


**Symptoms:** Dyspnea, orthopnea, sudden pleuritic chest pain.



fetal = fatal \*

<p>These images are taken from base of lung up to the neck.</p>	
<p style="text-align: center;">↓</p>	
<p>pulmonary vessels clearly enhanced “clear enhancing pattern”, so if there is thrombus it will be very clear.</p>	
<p style="text-align: center;">↓</p>	
<p>In the great vessels there is no thrombus but the thrombus is in the small peripheral arteries</p>	
<p>This illustrate what we can see in CT, after axial cut we can change setting to see vessels in details also we can change it into coronal image “as X-ray” also we can see sagittal images .This is axial section</p>	

## CTA Pulmonary Vasculature




- Massive PE “in the major vessels” involves more than one side and even peripheral vessels”.
- Coronal CTA filling defect = defect in contrast enhancement.

- coronal section in CT angiography. (both above images).

# CT Angiogram

- By using CT angiogram we able to "zoom" and see peripheral small vessels.
- In this image we have **filling defect** of the contrast which **means PE**.
- **White = contrast, grey = embolus or pulmonary thrombus** which can be peripheral, central or all




## Aortic Arch Anatomy ( For your knowledge )

We can use **MRI (MR angiography)** for examining great vessels, we can do it even without giving IV contrast, but if we want clear images we will use IV contrast.


But there's **disadvantages**:

- 1- if we didn't give contrast we are not going to see small vessels peripherally
- 2- Takes **45 mins**, cannot be done for a patient with acute PE (orthopnea & dyspnea)
- 3- Patients with cardiac devices such as pacemaker can't do it.



## Aortic Aneurysm ( For your knowledge )


- Right cardiac border is formed by right atrium and left border by Left ventricle and the apex.
- **Aortic knop/knuckle** is part of aortic arch so if there is pouch it's called aortic aneurysm  
It is enlarged if there's:
  1. Increased pressure
  2. Increased flow
  3. Change in aortic wall





# Heart & Vessels ( Just for your knowledge )

**Note:** the doctor didn't explain this, just go through it.

- Any and or all heart chambers may enlarge when the heart becomes diseased. Cardiomegaly = a big heart. The heart may enlarge due to a number of diseases e.g. valve disease, high blood pressure, congestive heart failure.
- If the heart fails, the lung often becomes congested. Early on the pulmonary vessels appear more prominent as in this case. More advanced failure can result in a condition of pulmonary edema which is fluid flooding into the alveoli of the lungs causing the patient marked shortness of breath.



## Anatomy on Normal Chest X-ray ( Just for your knowledge )



- There are 7 contours to the heart in the frontal projection in this system. But only the top five are really important in making a diagnosis.




# Cardiothoracic Ratio

In the past, the only way to know if the heart is enlarged or not is calculating the **cardiothoracic ratio**. **How to do it?**

We take a **transverse line at the widest part of the rib cage** ( just above costophrenic angle), and an **other transverse line at the widest part of the heart** and we divided these two lines. it should be less than the half of chest diameter (**<50%**).

Now, we do echocardiography to know if the heart is enlarged or not.



sometimes Cardiothoracic Ratio >50% but heart is **Normal** "false Positive"

e.g. patients with depressed sternum which makes their hearts' displayed "spread" so Cardiothoracic Ratio will be abnormal while in reality it's normal


sometimes Cardiothoracic Ratio <50% but heart is **Abnormal** " false Negative"

e.g. patient has cardiac problem such as Mitral Stenosis left atrium will be enlarged "has nothing to do with "borders" so in CXR not enlarged but in reality it's enlarged

<ul style="list-style-type: none"><li>- <b>Extracardiac causes of cardiac enlargement:</b> Portable AP films, Obesity, Pregnancy, Ascites, Straight back syndrome, <b>Pectus excavatum</b></li><li>- CTR &gt;50% but heart is normal, this is because there is an extra-cardiac cause for the apparent cardiomegaly (<b>Pic1</b>)</li><li>- On the lateral film(<b>Pic2</b>), the arrows point to the inward displacement of the lower sternum in a pectus excavatum deformity.</li></ul>	Two chest X-rays labeled 'pic1' and 'pic2'. 'pic1' is a frontal view showing a cardiothoracic ratio >50% with a red double-headed arrow and the text '>50%'. 'pic2' is a lateral view showing blue arrows pointing to the inward displacement of the lower sternum, characteristic of pectus excavatum.
<p>-Obstruction to outflow of the ventricles e.g.: ventricular hypertrophy <b>We must look at cardiac contours</b></p> <ul style="list-style-type: none"><li>- Here is an example of a heart which is less than 50% of the CTR in which the heart is still abnormal. This is recognizable because there is an <u>abnormal contour</u> to the heart (arrows)</li></ul>	A frontal chest X-ray showing a cardiothoracic ratio <50% with a red double-headed arrow and the text '<50%'. Three white arrows point to the abnormal contour of the heart, indicating a condition like ventricular hypertrophy.

# Ascending Aorta ( Just for your knowledge )

**Note:** the doctor didn't explain this, just go throw it.





Know the site and if it'd dilated or not



Enlarged with:


- Increased pressure
- Increased flow
- Changes in aortic wall

# Finding the Main Pulmonary Artery ( Just for your knowledge )



- the 2 pictures below are important.

**DEXTROCARDIA (قلب يميني)**



**cardiac displacement with scoliosis**



# The Pulmonary Vasculature

Pulmonary vasculature is an indirect sign of the status of the heart.

This means if the patient is having cardiac disease like mitral stenosis it will reflect on the pulmonary vasculature.


## Five States of the Pulmonary Vasculature:

- Normal
- Pulmonary venous hypertension
- Pulmonary arterial hypertension
- Increased flow
- Decreased flow - mostly unrecognizable even when it is present

### • What to evaluate ?


#### 1- Normal distribution of flow (upper vs. lower lobe):

- In erect position, blood flow to bases (blue circle) is bigger than flow to apices (green circle) because the size of vessels at bases is normally bigger than apical vessels size.
- You can't measure size of vessels at the left base because the heart obscures them




#### 2- Normal distribution of flow (Central Vs peripheral):

- Central vessels give rise to progressively smaller peripheral branches
- Normal tapering of vessels from central to peripheral



## Venous Hypertension

RDPA usually > 17 mm




- Patient present with dyspnea in ER, and did CXR. What is abnormal in this CXR?

1-Consolidation “air space disease” Is it taking the shape of the lobe or diffused ? Diffused  
Not diffused completely it has low distribution because it’s diffused when the patient has massive heart failure

2-Cardiomegaly “ do you see the borders?” the right is clear but the left not clear because there is fluid in the lung “blurred” or congested”

3- Costophrenic angle not clear “ because this image done while the patient is semi supine, However. The patient has pleural effusion”

### CHF: ( Just for your knowledge )



- Butterfly area “as it coming out from the hilum “ the area of consolidation “bat”

Scenario:

2 doctors argue: **DR.1** said consolidation due to infection and **Dr.2** said patient has cardiac failure, In this case **History** helps. patient who has **infection** has **fever** & patient with **HF** has **dyspnea and orthopnea** so there are differences in clinical presentation.

- if patient has **HF** there will be dramatic change in **CXR** within 1 hour after taking medication. However, if it’s **infection** won’t be any improvement within 1hour even with massive antibiotics

- So this is defiantly indication for **Acute pulmonary edema.**


### Important

#### ACUTE PULMONARY EDEMA:




seen when the interlobular septa in the pulmonary interstitium become prominent. caused by pulmonary edema


- **KERELY'S B-LINES:( Just for your knowledge )**



# Cardiac CT




- Coronary is better to be seen in CT: limited time, IV contrast “ a lot of details you will be able to see” .
- Will take Sagittal, axial and coronal images.
- **Notice the clear details:** muscle of the heart, lumen, chambers and septum also the fat between cardiac muscle and pericardium “very thin line “, space between pericardium and heart supposed to be filled with fat if it’s filled with water we call it **pericardial effusion** “sometimes after trauma rather than fluid you find blood”.




- Whenever we **encounter a large heart figure** , we should always be aware of the possibility of **pericardial effusion** simulating a large heart.
- On chest x-ray it looks as if this patient has dilated heart while on CT it is clear, that it is the pericardial effusion is responsible for the enlarged heart figure.
- Why isn’t cardiac Tamponade ? Different history
- In chest x-ray we see cardiac enlargement

# CARDIAC CHAMBERS CT



Axial slices through the heart



4-chamber view. RA=right atrium, RV=right ventricle, LA=left atrium, LV=left ventricle


we are able to see chambers, different phase of contrast , left side and right side, cardiac mobility “ pulsating heart”.

## Maximum Intensity Projection Soft Plaque in Proximal LAD


CT Coronary angiography “a true revolution in cardiac imaging”

We use it to see:

- Coronary arteries narrowing and vessels in general
  - Valves
  - Heart muscle thickness
- ❖ The narrowing of the vessel is called area of stenosis or plaque
- ❖ If the intensity of narrowed vessel is similar to muscle it called **soft plaque**
- ❖ Treated by placement of stent through cardiac catheter angiography also called balloon angiography or Angioplasty




NARROWED LUMEN



Cardiac stent

# Plaque visualization




## PLAQUE = VASCULAR NARROWING

- We are able to determine where is the narrowing and insert cardiac cath to put stent in the narrowing area, we can see clearly stenotic area and which vessel has stenosis and the degree.
- This appearance we used to have it only by catheter angiography, and we do this procedure only when the patient needs dilatation of the narrowing or replacement of stent.
- But if you need shorter technique do this one "MDCT"\* you can see stenosis very clearly as angiography, we can determine the site, measure the length or see it in 3D image.
- After applying stent no need to see how is it by cardiac cath, just follow up with CT.
- How are we able to see the whole vessel although it has stenosis?  
As the vessel not blocked totally the contrast will pass.


## Soft Plaque Visualization:

- Another example of soft plaque "not calcified", in this case they can apply balloon dilatation and put stent.
- by **CTA**: muscle, lumen of the heart, vessels, cause of narrowing While you can't see all these in "catheter angiography"



## Calcific plaque "hard":

- very dense, you are able to see the site in different parts of coronary vessels.
- you can see details of plaque and measure them also by computer you can see vessels spread out





# Visualization of vessels with CT + contrast

## (understand the Pictures)

- ★ Just look how the vessels appear with contrast
- ★ you won't be asked about names of vessels

