

# GI Bleeding Approach

# Case 1

- A 65 years old male referred for evaluation of 4 months HX of weight loss, fatigue , and weakness. He also gave history of passing dark stool intermittently for the last 3 months. He is known DM on insulin , hyperlipidemia on statin and occasionally aspirin

- What other information you would like to ask?

## ESSENTIALS OF DIAGNOSIS

- Symptoms: Coffee ground vomiting, hematemesis, melena, hematochezia, anemic symptoms
  - Past medical history: Liver cirrhosis, use of non-steroidal anti-inflammatory drugs
  - Signs: Hypotension, tachycardia, pallor, altered mental status, melena or blood per rectum, decreased urine output
  - Bloods: Anemia, raised urea, high urea to creatinine ratio
- 

- What is the likely diagnosis?

# Causes of UGIB

**Table 1**  
**Frequency of common causes of upper gastrointestinal bleeding**

<b>Diagnosis</b>	<b>Frequency (Percentage)</b>
Peptic ulcer disease, including duodenal and gastric ulcer	28–59
Variceal bleeding	4–14
Mucosal erosive disease, including esophagitis, gastritis, and duodenitis	1–31
Mallory-Weiss tear	4–8
Malignancy	2–4
Arteriovenous malformation	3
Gastric antral vascular ectasia	~ 1
Dieulafoy lesion	~ 1

Gibson et al. Gastrointest Endosc Clin N Am 2011;21:583-96.

- What will be the next step?



## Case 2

- A 42 years old male complaining of chronic recurrent epigastric pain which worsen recently especially when he is fasting
- For the last 2 days he started to have frequent vomiting associated with blood
- He is not known to have any chronic medical problems and not on any medications

- What is the best next step in the approach of such patient?

- Detailed HX
- Full Physical examination

- How would you assess the bleeding severity?

# Risk Stratification

Glasgow- Blatchford Score (GBS)

Rockall Score

Modified-GBS

AIMS65

**Table 1** | Glasgow–Blatchford score assessment criteria

	Risk factors at presentation	Threshold	Score
Urea	Blood urea nitrogen (mmol/l)	6.5–7.9	2
		8.0–9.9	3
		10.0–24.9	4
		≥25.0	6
CBC	Hemoglobin for men (g/l)	120–130	1
		100–119	3
		<100	6
	Hemoglobin for women (g/l)	100–120	1
		<100	6
Physical	Systolic blood pressure (mmHg)	100–109	1
		90–99	2
		<90	3
	Heart rate (bpm)	>100	1
History	Melena	Present	1
	Syncope	Present	2
	Hepatic disease	Present	2
	Cardiac failure	Present	2

Total score (0–23). Patients with scores >0 are considered to be at high risk. Permission obtained from Elsevier Ltd © Blatchford, O. *et al. Lancet* 356, 1318–1321 (2000).


## B Rockall Score

		Variable	Points
Complete Rockall Score	Clinical Rockall Score	Age	History 0 1 2
		<60 yr	
		60–79 yr	
		≥80 yr	2
		Shock	Physical 2
		Heart rate >100 beats/min	
		Systolic blood pressure <100 mm Hg	2
		Coexisting illness	History 2 3
		Ischemic heart disease, congestive heart failure, other major illness	
		Renal failure, hepatic failure, metastatic cancer	
Endoscopic diagnosis			
No lesion observed, Mallory–Weiss tear	0		
Peptic ulcer, erosive disease, esophagitis	1		
Cancer of upper GI tract	2		
Endoscopic stigmata of recent hemorrhage			
Clean base ulcer, flat pigmented spot	0		
Blood in upper GI tract, active bleeding, visible vessel, clot	2		

Hearnshaw et al. Aliment Pharmacol Ther 2010;32:215-24.


- What is the diagnosis and the associated risk factors?

# Spurting Blood



Gralnek et al. N Engl J Med 2008;359:928-37.

# Non-bleeding Visible Vessel




Gralnek et al. N Engl J Med 2008;359:928-37.

# Flat, Pigmented Spot



Gralnek et al. N Engl J Med 2008;359:928-37.

# Clean Base



Gralnek et al. N Engl J Med 2008;359:928-37.

Age >65  
Previous peptic ulcer  
Previous ulcer-related upper GI complication  
High-dose NSAIDs  
Multiple NSAID use  
Selection of NSAID (e.g., COX-1 vs. COX-2 inhibition)  
NSAID-related dyspepsia  
Aspirin (including cardioprotective dosages)  
Concomitant use of  
    NSAID plus low-dose aspirin  
    Oral bisphosphonates (e.g., alendronate)  
    Corticosteroids  
    Anticoagulant or coagulopathy  
    Antiplatelet drugs (e.g., clopidogrel)  
    Selective serotonin reuptake inhibitor  
Chronic debilitating disorders (e.g., cardiovascular disease, rheumatoid arthritis)  
*Helicobacter pylori* infection  
Cigarette smoking  
Alcohol consumption

<sup>a</sup>Combinations of risk factors are additive.

Data from references 1, 12–15, 20, and 29.

# H pylori

- Patients with bleeding peptic ulcers should be tested for H. pylori
  - Receive eradication therapy if present
  - Confirmation of eradication
- Negative H. pylori diagnostic tests obtained in the acute setting should be repeated


# Case 3

- A 52 years old lady presented to ER with one day history of vomiting of fresh blood. She also notices passing black tarry stool. She is feeling dizzy and unwell
- Past HX of jaundice no other medical problems and not on any medications
- Clinically jaundiced and pale
- Vital signs BP 100/70 pulse 110/min
- Abdomen examination showed liver span of 7 cm and spleen felt 3 fingers below costal margin with few spider nevi seen over chest

- what is the likely diagnosis of this case and list 4 common aetiology ?

## **Causes of liver cirrhosis:**

- 1) Viral Hepatitis B, C.
- 2) Alcoholic liver disease.
- 3) Non-alcoholic fatty liver disease (NAFLD).
- 4) Autoimmune hepatitis.
- 5) Primary biliary cirrhosis.
- 6) Secondary biliary cirrhosis (associated with chronic extrahepatic bile duct obstruction).
- 7) Primary sclerosing cholangitis.
- 8) Hemochromatosis
- 9) Wilson disease.
- 10) Alpha-1 antitrypsin deficiency.
- 11) Granulomatous disease (eg, sarcoidosis).
- 12) Type IV glycogen storage disease.
- 13) Drug-induced liver disease (eg, methotrexate, alpha methyl dopa, amiodarone).
- 14) Venous outflow obstruction (eg, Budd-Chiari syndrome, veno-occlusive disease).
- 15) Cardiac cirrhosis: chronic right-sided heart failure, tricuspid regurgitation.



### Symptoms of liver cirrhosis

- General malaise, fatigue
- Anorexia / weight loss
- Feeling of enlarged abdomen
- Swollen abdomen / legs
- Nose bleed / bleeding from lower limbs
- Jaundice / itch
- Hand tremors

### Physical findings

- Skin pigmentation
- Xanthoma
- Spider angioma
- Palmar erythema
- Finger clubbing (hepatopulmonary syndrome)
- Caput medusae
- Gynecomastia
- Fever
- Hepatoceleoma
- Hepatic halitosis (dimethyls-ulphide, ketons in the expired breath)
- Jaundice
- Ascites, lower thigh edema
- Hepatic encephalopathy
- Bleeding plaque / purpura

- What is the priority in the management of this patient?

# IV Fluid Resuscitation

- What is the target Hb and INR prior to the endoscopy for this cases?

# 3- Blood Transfusions

The role of transfusion in clinically stable patients with mild GI bleeding remains controversial, with uncertainty at which hemoglobin level transfusion should be initiated

Literature suggesting poor outcomes in patients managed with a liberal transfusion


Marik PE, Corwin HL. Crit Care Med 2008; 36: 2667 – 2674

Restellini S, Kherad O, Jairath V et al. Aliment Pharmacol Ther 2013; 37: 316 – 322

# 3- Blood Transfusions (cont'd)


The restrictive RBC transfusion had significantly improved survival and reduced rebleeding

# 3- Blood Transfusions (Cont'd)



# 3- Blood Transfusions (Cont'd)

**A Survival, According to Transfusion Strategy**



**No. at Risk**

Restrictive strategy	444	429	412	404	401	399	397	395	394	392
Liberal strategy	445	428	407	397	393	386	383	378	375	372

**B Death by 6 Weeks, According to Subgroup**

Subgroup	Restrictive Strategy no. of patients/total no. (%)	Liberal Strategy no. of patients/total no. (%)	Hazard Ratio (95% CI)	P Value
Overall	23/444 (5)	41/445 (9)	0.55 (0.33–0.92)	0.02
Patients with cirrhosis	15/139 (11)	25/138 (18)	0.57 (0.30–1.08)	0.08
Child–Pugh class A or B	5/113 (4)	13/109 (12)	0.30 (0.11–0.85)	0.02
Child–Pugh class C	10/26 (38)	12/29 (41)	1.04 (0.45–2.37)	0.91
Bleeding from varices	10/93 (11)	17/97 (18)	0.58 (0.27–1.27)	0.18
Bleeding from peptic ulcer	7/228 (3)	11/209 (5)	0.70 (0.26–1.25)	0.26

0.1      1.0      10.0

Restrictive Strategy Better      Liberal Strategy Better

# Patients receiving anticoagulants

Correction of coagulopathy is recommended

Endoscopy should not be delayed for a high INR  
unless the INR is supratherapeutic

# Timing and need for early endoscopy

- Definition of early endoscopy
  - Ranges from 6 to 24 hours AFTER INITIAL PRESENTATION
- May need to be delayed or deferred:
  - Active acute coronary syndromes
  - Suspected perforation

# Case 4

- A 47 years old male known to have alcoholic liver disease presented with hematemesis of large amount and dizziness after resuscitation an upper GI endoscopy done which showed multiple large oesophageal varix which was banded , however 12 hrs post endoscopy he continued to have melena with drop of Hb and hypotension

- What is the next step in the patient management?

# Gastroenterology



# Interventional Rad.



YOU ARE NOT ALONE


Intensive Care




Surgery




# Summary for steps of GI bleeding approach




## Algorithm for management of acute GI bleeding





# When to go to surgery?



# Conclusions

- \* Resuscitation should be initiated prior to any diagnostic procedure
- \* Gastrointestinal endoscopy allows visualization of the stigmata, accurate assessment of the level of risk and treatment of the underlying lesion
- \* Intravenous PPI therapy after endoscopy is crucial to decrease the risk of cardiovascular complications and to prevent recurrence of bleeding
- \* *Helicobacter pylori* testing should be performed in the acute setting