
Outbreak Investigation

Dr Armen Torchyan

Department of Family and Community Medicine



Objectives

- Understand what constitutes to an outbreak
- Distinguish between endemic and epidemic
- Learn the importance of investigating an outbreak
- List the steps of an outbreak investigation
- Describe types of studies used to investigate an outbreak
- Read an epidemic curve and use it in estimating the incubation period
- Calculate the attack rate from outbreak investigation data

Key terms

Sporadic: a disease that occurs infrequently and irregularly.

Endemic: the habitual presence of a disease within a given geographic area.

Hyperendemic: persistent, high levels of disease occurrence.

Epidemic: the occurrence of more cases of disease than expected in a given area or among a specific group of people over a particular period of time. Usually, the cases are presumed to have a common cause or to be related to one another in some way


Outbreak: epidemic limited to localized increase in the incidence of disease

Cluster: aggregation of cases in a given area over a particular period without regard to whether the number of cases is more than expected

Pandemic: An epidemic that has spread over several countries or continents, usually affecting a large number of people.

Pneumonia and Influenza Mortality for 122 U.S. Cities

Week ending 4/17/2010



2009-2010 Influenza Season – Week 15, ending April 17, 2010

Outbreak settings

- Most outbreaks are caused by exposure to contaminated food, water, or direct contact with others
 - Food borne outbreaks: most common
 - Waterborne outbreaks: infection occurs by either ingesting contaminated water or swimming in contaminated water
 - Community/institution acquired outbreaks: most widely varied
 - Others
- Illness can be minor “self-limited” diseases, or major deadly diseases
- Successful investigation of an outbreak can lead to positive advances in future protection of public health

Food-borne Outbreaks

- The most common outbreak is in a food borne setting.
- A food-borne outbreak is generally considered to exist if there are more than two reports of a similar illness from the same food source, frequently in a restaurant or at a community dinner.
- A food-borne outbreak may have a widely varied number of cases, and has no seasonal distribution.
- While the food-borne outbreak is the most common, in only about 50% of the outbreaks is the food culprit identified, so a large number of these outbreaks go unsolved.

Water-borne Outbreaks

- In disease that occurs from a water-borne outbreak, infection occurs by either ingesting water contaminated by pathogens or by swimming in water contaminated by pathogens.
- Most often, these outbreaks are common source types.
- The numbers of cases in these outbreaks can be variable and often unknown.
- The most common agents responsible for water-borne outbreaks are norovirus, Shigella, Giardia, Cryptosporidiosis, and E. coli.

Community/Institution Acquired

- The most widely varied of the outbreak settings
- Include most all types of infectious diseases, such as respiratory diseases and gastrointestinal diseases.
- Transmitted most often by person-to-person transmission in schools, hospitals, daycare, nursing homes, prisons, and high density living areas such as military barracks, hotels, and even airplanes.
- Some common agents that cause the diseases acquired in a community setting include norovirus, varicella, influenza, rhinovirus, parasites, and adenovirus.

Exercise

1. 22 cases of legionellosis occurred within 3 weeks among residents of a particular neighborhood (usually 0 or 1 per year)

Epidemic disease

2. Average annual incidence was 364 cases of pulmonary tuberculosis per 100,000 population in one area, compared with national average of 134 cases per 100,000 population

3. Over 20 million people worldwide died from influenza in 1918-1919

Hyperendemic

4. Single case of histoplasmosis was diagnosed in a community

Pandemic

5. About 60 cases of gonorrhea are usually reported in this region per week, slightly less than the national average


Sporadic

Endemic

Detecting outbreaks

<https://healthmap.org/en/>

- Regular, timely analysis of surveillance data
- Reports of cases of notifiable diseases
- Alert clinicians call the health department.
- Patients or community members can report to the health department




Epidemic Patterns

- **Common-source**
 - Point - everyone becomes ill within one incubation period
 - Continuous - exposed over a period of days, weeks, or longer
 - Intermittent - intermittent nature of the exposure
- **Propagated**
- Mixed
- Other


Common-source

- All cases of the infectious disease are acquired from the same source in a limited period of time and in a limited geographical location.
- It is also characterized by very minimal (or zero) transmission from person to person.
- Generally, a common source outbreak has a smaller number of cases than a propagated outbreak and is often caused by contaminated food or water.
- A typical example of a common source epidemic is a food borne illness caused by exposure to one specific food or restaurant.
- Common source epidemics are usually characterized by a dramatic single “peak” of cases.
- Many common source outbreaks go unreported since they are generally small in numbers and often don't come to the attention of public health authorities.


Common-source Point



Common-source Continuous




Common-source Intermittent



Propagated

- Characterized by an outbreak that continues over an extended period of time.
- This outbreak has individuals exposed to the original source, but then will also have secondary infections in individuals exposed to those initially ill people via person-to-person spread.
- The propagated epidemic usually lasts for a longer period of time and has various numbers of “peaks” of cases over time.
- The initial source often resolves, but the outbreak continues by affected persons infecting other persons.
- Propagated outbreaks often result in larger numbers of cases than common source outbreaks.
- Most outbreaks of respiratory diseases, such as influenza, are propagated outbreaks, as well as some food or water borne outbreaks such as those occurring from norovirus infections.

Propagated



Exercise

1. 21 cases of shigellosis among children and workers at a day care center over a period of 6 weeks, no external source identified (incubation period for shigellosis is usually 1–3 days)

Propagated

2. 36 cases of giardiasis over 6 weeks traced to occasional use of a supplementary reservoir (incubation period for giardiasis 3–25 days or more, usually 7–10 days)

Intermittent or continuous common source


3. 43 cases of norovirus infection over 2 days traced to the ice machine on a cruise ship (incubation period for norovirus is usually 24–48 hours)

Point source

Making Epidemic Curves

1. Plot the number of reported cases on the y-axis.
2. Plot the date/time of symptom onset on the x-axis.
3. Choose an appropriate time interval for the x-axis, or try several to see which best represents the data.
4. Include pre-outbreak time on the x-axis to show the “baseline” disease level and to show visually when the outbreak began.
5. Label the x and y-axes clearly.
6. Give the epi curve a descriptive, self-explanatory title.
7. Include more detailed information, such as cases by geographic location or by symptom, if helpful.
8. To be technically correct, make the bars touch each other (unless there are periods of time with no cases, in which case there will be space between the bars).

Figure 5. Epidemic curve from outbreak of norovirus gastroenteritis in U.S. Army trainees in 1998



Steps in an Outbreak Investigation

Each day, health departments learn about cases or outbreaks that require investigation.

- Although the U.S. CDC receives over 1000 reported outbreaks per year, this is likely only the tip of the iceberg
- Many times outbreaks are not seen in their entirety and never reported
- 75 million food-borne illnesses in the U.S. each year are estimated to be part of an outbreak that is not reported

Epidemiologists have an “outbreak cookbook”

- The 10 steps of an outbreak investigation assist epidemiologists in investigating an outbreak
- Allows investigators to have the best success in determining the cause of the outbreak and preventing future cases of the same disease

Steps in an Outbreak Investigation

Steps organized into categories that:

- Identify that a problem exists
- Measure the problem
- Find the responsible agent
- Prevent it from occurring further

Each step is dependent on the successful completion and information obtained in the previous step(s)

Steps in an Outbreak Investigation

1. Confirm the Diagnosis
 2. Confirm the Existence of an Outbreak/Epidemic
 3. Define a Case and Count Cases
 4. Orient Data in Terms of Person, Place and Time
 5. Determine Who Is at Risk
 6. Develop a Hypothesis and Test It
 7. Determine Control Measures
 8. Plan a More Systematic Study
 9. Execute Disease Control and Prevention Measures
 10. Prepare a Written Report
- Identify that a problem exists
- Measure the problem
- Find the responsible agent
- Prevent it from occurring further

Step 1: Confirm the Diagnosis

Symptoms need to be reviewed.

Disease must be properly diagnosed:

- “Re-look” at records and visit existing cases
- Ensure suspected illness is properly diagnosed
- Confirm laboratory results/rule out errors
- Visit/assess patients (cases)
- Summarize clinical and laboratory findings

Step 2: Confirm the Existence of an Outbreak/Epidemic

Very important to establish that the disease being seen in the community is in fact an outbreak


- Investigations can be costly and time consuming
- Normal rate of illness in the population must be known
- Time is critical
 - What is expected number of disease?
 - Use Health Department Surveillance data
 - Use hospital discharge records
 - Use vital records
 - Use registries
 - May need to use neighboring rates
 - Last resort: conduct a survey

Step 2: Confirm the Existence of an Outbreak/Epidemic

Establishing existence of a disease uses concept of *epidemic threshold*

- Normal rates of disease over time used to determine range of normal high and low limits
- Normal high limit used to determine if there is an excess
- Any amount of disease over threshold is considered to be excess of normal

Pneumonia and Influenza Mortality for 122 U.S. Cities Week ending 4/17/2010



2009-2010 Influenza Season – Week 15, ending April 17, 2010

Step 3: Define a Case and Count Cases

- Case classification (definition) should include:
 - Clinical symptoms (\pm 1 symptom can change accuracy)
 - Laboratory verification
 - Restrictions of person, place, and time
 - Do not include the exposure or risk factor you are interested in evaluating
 - Must classify if cases are “confirmed, suspected, or probable”
- Count cases:
 - Important to find and count all cases that exist
 - Cast a wide net
 - Health clinics, hospitals
 - Advertise
 - Ask other cases

Meningococcal Disease — Case Definition

Clinical case definition

An illness with sudden onset of fever ($>38.5^{\circ}\text{C}$ rectal or $>38.0^{\circ}\text{C}$ axillary) and one or more of the following: neck stiffness, altered consciousness, other meningeal sign or petechial or puerperal rash.

Laboratory criteria for diagnosis

Positive cerebrospinal fluid (CSF) antigen detection or positive culture.

Case classification

Suspected: A case that meets the clinical case definition.

Probable: A suspected case as defined above and turbid CSF (with or without positive Gram stain) or ongoing epidemic and epidemiological link to a confirmed case.

Confirmed: A suspected or probable case with laboratory confirmation.

Source: Pan American Health Organization. Case Definitions Meningococcal Disease. Epidemiological Bulletin 2002; 22(4):14–5.

Line listing

Case #	Report Date	Onset	Physician Diagnosis	Signs/Symptoms						Labs		Demographics	
				N	V	A	F	D	J	HAIgM	Other	Sex	Age
1	10/12/02	10/5/02	Hepatitis A	1	1	1	1	1	1	1	Low SGOT	M	37
2	10/12/02	10/4/02	Hepatitis A	1	0	1	1	1	1	1	Low Alt	M	62
3	10/13/02	10/4/02	Hepatitis A	1	0	1	1	1	1	1	Low SGOT	M	38
4	10/13/02	10/9/02	NA	0	0	1	0	?	0	NA	NA	F	44
5	10/15/02		Hepatitis A	1	1	1	1	1	0	1	Hbs/Ag-	M	17
6	10/16/02	10/6/02	Hepatitis A	0	0	1	1	1	1	1	SGOT=24	F	43

N=nausea V=vomiting A=elevated aminotransferase F=fever D=discreet onset J=jaundice HAIgM=hepatitis AlgM antibody test SGOT=serum glutamic oxaloacetic transaminase ALT=alanine aminotransferase Hbs=hepatitis B surface antigen Ag-=antigen negative

1="yes", 0="no"

* This table illustrates a line listing that might be used during an outbreak of hepatitis A. It was adapted from the CDC's "Excellence in Curriculum Integration through Teaching Epidemiology" program. Additional variables that might be helpful to include are drug use, occupation, meal at restaurant X, neighborhood of residence and sexual orientation.

Step 4: Orient Data in Terms of Person, Place, and Time

- Get to know your data
- Descriptive Epidemiology
 - Person: age, race, gender, medical status, exposures
 - Place: map cases (GIS)
 - Map attack rates, not numerators
 - Time: Epidemic curve
- Orienting groups of cases by time provides more information about the outbreak and possible cause
 - Graphing number of cases over time provides an **epidemic curve**
 - Earliest set of cases to appear on graph can identify date of first exposure
 - Can help identify type of outbreak and secondary attack rate


Figure 1.1 Spot map of deaths from cholera in Golden Square area, London, 1854 (redrawn from original)



Source: Snow J. *Snow on cholera*. London: Humphrey Milford: Oxford University Press; 1936.

Haiti Cholera Outbreak - Map of Cumulative Attack Rate With Data on Cases, Hospitalizations, and Deaths

Cumulative Attack Rate per Department through 21-Feb-2011



NOTES:

1. Port-au-Prince includes the following communes: Carrefour, Cité Soleil, Delmas, Kenscoff, Pétion Ville, Port-au-Prince, and Tabarre
2. Numbers in parentheses indicate the daily total

Data Sources: MSPP, Institut Haitien de Statistique et d'Informatique, Centre National de l'Information Géo Spatiale
Creation Date: 01-Mar-2011

Nord-Ouest

Population: 662,777
Cases: 15,591 (20)
Attack Rate: 2.4%
Hospitalizations: 9,178 (15)
Hospitalization Rate: 1.4%
Deaths: 238 (0)
Death Rate Per 100,000 pop: 36
First Confirmed Case: Oct 26, 2010

Nord

Population: 970,495
Cases: 26,311 (56)
Attack Rate: 2.7%
Hospitalizations: 26,311 (42)
Hospitalization Rate: 2.7%
Deaths: 623 (0)
Death Rate Per 100,000 pop: 64
First Confirmed Case: Oct 25, 2010

Nord-Est

Population: 358,277
Cases: 9,149 (40)
Attack Rate: 2.6%
Hospitalizations: 7,603 (29)
Hospitalization Rate: 2.1%
Deaths: 257 (0)
Death Rate Per 100,000 pop: 72
First Confirmed Case: Nov 13, 2010

HAITI

Population: 9,923,243
Cases: 245,183 (615)
Attack Rate: 2.5%
Hospitalizations: 132,293 (318)
Hospitalization Rate: 1.3%
Deaths: 4,625 (1)
Death Rate Per 100,000 pop: 47

Artibonite

Population: 1,571,020
Cases: 62,641 (130)
Attack Rate: 4.0%
Hospitalizations: 23,991 (48)
Hospitalization Rate: 1.5%
Deaths: 868 (0)
Death Rate Per 100,000 pop: 55
First Confirmed Case: Oct 21, 2010

Centre

Population: 678,626
Cases: 20,608 (98)
Attack Rate: 3.0%
Hospitalizations: 8,644 (40)
Hospitalization Rate: 1.3%
Deaths: 357 (0)
Death Rate Per 100,000 pop: 53
First Confirmed Case: Oct 24, 2010

Grande Anse

Population: 425,878
Cases: 14,002 (31)
Attack Rate: 3.3%
Hospitalizations: 10,213 (21)
Hospitalization Rate: 2.4%
Deaths: 824 (0)
Death Rate Per 100,000 pop: 193
First Confirmed Case: Nov 19, 2010

Nippes

Population: 311,497
Cases: 3,013 (17)
Attack Rate: 0.97%
Hospitalizations: 1,744 (11)
Hospitalization Rate: 0.56%
Deaths: 140 (1)
Death Rate Per 100,000 pop: 45
First Confirmed Case: Nov 18, 2010

Port-au-Prince

Population: 2,476,787
Cases: 58,429 (80)
Attack Rate: 2.4%
Hospitalizations: 23,969 (23)
Hospitalization Rate: 0.97%
Deaths: 553 (0)
Death Rate Per 100,000 pop: 22
First Confirmed Case: Unknown

Ouest


Population: 1,187,833
Cases: 20,280 (25)
Attack Rate: 1.7%
Hospitalizations: 11,839 (20)
Hospitalization Rate: 1%
Deaths: 309 (0)
Death Rate Per 100,000 pop: 26
First Confirmed Case: Oct 23, 2010

Sud


Population: 704,760
Cases: 10,135 (112)
Attack Rate: 1.4%
Hospitalizations: 6,204 (60)
Hospitalization Rate: 0.88%
Deaths: 228 (0)
Death Rate Per 100,000 pop: 32
First Confirmed Case: Nov 10, 2010

Sud-Est


Population: 575,293
Cases: 2,423 (6)
Attack Rate: 0.42%
Hospitalizations: 2,221 (6)
Hospitalization Rate: 0.39%
Deaths: 228 (0)
Death Rate Per 100,000 pop: 40
First Confirmed Case: Nov 14, 2010



Epidemic curve



Epidemic curve



Step 5: Determine Who is at Risk



Necessary to gather information from subjects who are not cases

Population at risk is important because:

- It will be the population used as controls
 - Approach subjects for a study to test the hypothesis to find the cause of the outbreak
- It defines the population for whom prevention and control measures will be targeted

Step 5: Determine Who is at Risk

- Where did the cases come from
- Using definition of cases, identify population with the same criteria
 - Geographic location
 - Time period
 - Population characteristic
- Look for any remaining cases in population at risk
- This population will also be the target of prevention and control measures



Step 6: Develop a Hypothesis and Test It

- Develop a hypothesis to confirm the cause of disease
 - Include suspected etiologic agent
 - Include mode of transmission
 - Identifies expected exposures to transmit the disease
 - Specifies population
- Test the hypothesis using a study design
 - Case-control study if the population is not well defined and speed of investigation is important
 - Cohort study if the population is small and well defined

Example of a Cohort Study

A community in Massachusetts experienced an outbreak of Salmonellosis. Health officials noted that an unusually large number of cases had been reported during a span of several days. Descriptive epidemiology was conducted, and hypothesis-generating interviews indicated that all of the disease people had attended a parent-teacher luncheon at a local school. The descriptive epidemiology convincingly indicated that the outbreak originated at the luncheon, but which specific dish was responsible? The investigators needed to establish which dish was responsible in order to clearly establish the source and to ensure that appropriate control measures were undertaken.

		Salmonellosis			Incidence ↓ 16/23 = 0.70
		Yes	No	Total	
Ate Cheese Appetizer (Exposed)	Yes	16	7	23	
	No	9	13	22	9/22 = 0.41

Risk Ratio = (Incidence in the exposed group) / (Incidence in the unexposed group)

Menu Item	Tot. Exp.	Attack Rate		RR	95% CI	p
		Exp.	Unexp.			
Cheese	23	70%	41%	1.70	0.96 – 3.01	0.053
Mushrooms	23	61%	55%	1.12	0.67 – 1.85	0.89
Pasta	30	53%	67%	0.80	0.49 – 1.31	0.50
Potato Salad	6	33%	62%	0.54	0.17 – 1.73	0.39
Veg.Lasagna	17	47%	64%	0.73	0.41 – 1.30	0.41
Chickn& Rice	14	43%	65%	0.66	0.34 – 1.28	0.30
Manicotti	27	93%	6%	16.67	2.47 – 112.30	<0.001
Veggies	19	63%	54%	1.17	0.72 – 1.92	0.10
Wings	13	46%	63%	0.74	0.39 – 1.41	0.50
Caesar Salad	6	17%	64%	0.26	0.04 – 1.58	0.04
Kielbasa	8	63%	57%	1.10	0.60 – 2.02	0.54
Chick.&Brocc	23	74%	41%	1.81	1.03 – 3.16	0.05
Chicken Parm	14	64%	55%	1.17	0.71 – 1.94	0.55
Calzone	16	50%	62%	0.81	0.46 – 1.42	0.43
Eggplnt Parm	7	57%	58%	0.99	0.49 – 1.98	0.64
Meatballs	13	46%	63%	0.74	0.39 – 1.41	0.31

Example of a Case-Control Study

Within a short period of time 20 cases of hepatitis A were identified in the Marshfield area. The epidemic curve suggested a point source epidemic, and the spot map showed the cases to be spread across the entire South Shore of Massachusetts. Hypothesis-generating interviews resulted in five food establishments that were candidate sources.

The investigators identified as many cases as possible, and they selected a sample of non-diseased people as a comparison group (the controls). The "controls" were matched to the cases with respect to age, gender, and neighborhood of residence. Investigators then ascertained the prior exposures of subjects in each group, focusing on food establishments and other possibly relevant exposures they had had during the past two months.

	Cases	Controls
Ate at Papa Gino's	10	19
Did not eat at Papa Gino's	9	19
	19	38

$$\text{Odds Ratio} = (10/19) / (9/19) = 1.1$$

	Cases	Controls
Ate at Ron's Grill	18	7
Did not eat at Ron's	1	29
	19	38

$$\text{Odds Ratio} = (18/7) / (1/29) = 75$$

Exercise

You are called to help investigate a cluster of 17 persons who developed brain cancer in an area over the past couple of years. Most, perhaps all, used cell phones. Which study design would you choose to investigate a possible association between cell phone use and brain cancer?

A case-control study is the design of choice

Step 7: Determine Control Measures


Important to control current outbreak and prevent future outbreaks

Control measures completely dependent on the identified source of the outbreak

- May include treating carriers of infection and vaccinating population at risk
- May include comprehensive training programs

Step 7: Determine Control Measures

- Destroy implicated food
- Close water source or beaches
- Treat carriers
- Vaccinate susceptible population
- Training
- Etc.



Step 8: Plan a More Systematic Study

It may be necessary to continue to study the outbreak with a more comprehensive design

- Initial study may be inconclusive
- Reconsider hypothesis
- Revisit patients
- Expand exposures
- Utilize additional lab specimens
- Additional or more refined control group
- Utilize a more comprehensive design
- Perform research to expand knowledge

Steps 9: Execute Disease Control and Prevention Measures

As investigation wraps up, it is important to use all information available to prevent the spread or resurgence of outbreak

Activities include:

- Implement the necessary control and prevention measures
 - Treat carriers
 - Vaccinate susceptible population
 - Etc.
- Surveillance for future disease occurrence
- Regular communication with affected population and health care facilities

Step 10: Prepare a Written Report

A written report should be prepared in a usual scientific format and should include information about:

- The setting and the methods used
- Results of any data collection and analysis
- The identified causative agent and source
- Recommendations for control and prevention

Report should be written for members of affected community

Summary

Disease outbreaks can be large or small

In order to determine source of outbreak, there are standard steps that should be taken to:

- Identify there is a problem
- Measure the outbreak
- Find the responsible agent
- Controlling the outbreak/prevention of further cases

References

Centers for Disease Control and Prevention. Principles of Epidemiology in Public Health Practice, 3rd Edition. Atlanta, GA: CDC; 2006 (updated in 2012)

Macera CA, Shaffer R, Shaffer PM. Introduction to Epidemiology: Distribution and Determinants of Disease. Clifton Park, NY: Delmar; 2012.

LaMorte WW. Outbreak Investigations. Boston University School of Public Health; 2016. Available online: <http://sphweb.bumc.bu.edu/otlt/MPH-Modules/PH/Outbreak/index.html>