



Radiology
Team 438

Nuclear medicine in Oncology


Lecture 26

Objectives

At the end of the lecture you will be able to answer the following questions:

- ❖ What are the tumor imaging and therapeutic radiopharmaceuticals?
- ❖ What are the nuclear medicine tumor imaging methods?
- ❖ What are the objectives of tumor imaging?
- ❖ What are the potential values of nuclear medicine tumor imaging methods?
- ❖ What is the role of nuclear medicine in the treatment of tumors?

Reviewed By



Noura Alturki
Jehad Alorainy

Color Index:

♦ Important

♦ Doctor's Notes

♦ Extra

♦ Female slides

♦ male slides

Team Leaders



Omar Aldosari



Leena Alnassar



Shahd Alsalamh

Done by:



&



Tumor Imaging

Isotopes show function not structure

» Tumor metabolic properties (changes in function)

To access how active is the tumor

Increased **vascularization**



Increased capillary **permeability**

Newly **proliferated** capillaries

Increased **blood flow**

Metabolically **active** cells; Consumes more protein and glucose for its growth which is the cause of weight loss so if we labeled glucose with a nuclear material their will be more concentration inside the tumor cells

Increased energy demand



» Tumor Specific useful properties

- ✓ High density of some **common receptors**
- ✓ Expression of several **specific receptors**
- ✓ Expression of **some specific tumor antigens**
Access antigen×antibody reaction

All these properties could be used for **imaging** and **therapy**

» Tumor Non-specific Diagnostic radiopharmaceuticals ★

PET or PET-CT:

- **F-18 FDG** (Fluorodeoxyglucose) - anaerobic metabolism

When there is a tumor with increased glucose consumption, there will be increased F-18 FDG concentration

Planar, SPECT or SPECT-CT:

- **Diphosphonates** - bone scan
- **Ga-67 citrate** - similar to FDG - localising agent
- **Tc99m Nanocolloid** - bone marrow scan
- **Tc99m MIBI / Thallium 201** - several tumors

Demonstrate tumor sites but are not specific for malignancy; Tells us there is a tumor but not what is it exactly

» Tumor Specific Diagnostic radiopharmaceuticals

PET or PET/CT:

- ★ **Gallium -68 - Octreotide analogues (Ga-68 DOTA) = somatostatin agent:**
 - For **neuroendocrine tumors**
- **Fluorine -18-fluorodeoxythymidine (F-18-FLT):** For tumor proliferation
- **Fluorine -18-fluoromisonidazole (F-18-FMISO):** For tumor hypoxia

Planar, SPECT or SPECT-CT: ★

- **I-123/131 MIBG** for neuroendocrine tumours
- **I-131** for differentiated thyroid carcinomas
- **In-111 or Tc99m octreotide** for tumours expressing somatostatin receptors.
- **Monoclonal antibodies** labelled with In-111, I-123/131 or Tc-99m

Binds directly to special tumor **antigens or receptors** or are accumulated by **special metabolic pathway**

» Therapeutic radiopharmaceutical

Non-specific

- **Sr-89, Sm-153, Re-189**
- **Bone pain palliation** (any metastatic pain is relieved)

Specific

- **I-131 : Thyroid cancer**, as specific diagnostic if tumor significantly accumulates
- **Y-90: Zevalin** monoclonal antibody for **B-cell lymphomas**


» NM Imaging modalities

■ **SPECT:** Single Photon Emission Computed Tomography and SPECT CT


(there is Abnormal uptake as seen in the arrow, but exact location unknown we use SPECT-CT)



Planar imaging
(Two dimensions)



Whole body Gallium scan: planar image of a patient with lymphoma




SPECT/CT Shows precise location which is, in this case, deep in the mediastinum (arrow)



← SPECT

■ **PET:** Positron Emission Tomography and PET CT

PET




planar; it's a gif :)


PET/CT



CT



Plane PET



Fused CT + PET

Role for Nuclear Medicine In Oncology



Diagnosis

- Specific or non-specific



Follow-up

- Early detection of recurrence



Staging

- Important for proper therapy



Treatment

- Specific or non-specific

» Non specific tumor imaging agents:

- **Tc-99m MDP bone scan:** Detection and follow up of bone metastasis
- **Gallium 67 (Replaced by F18-FDG) :** Staging ,Restaging & therapy assessment of HD , NHL , Lung cancer
- **Thallium 201 :** Tumor viability & tumor seeking. {Tc-99 m Agents (MIBI ,TETRO.)
- **F18 – FDG (Gold standard for tumour imaging) :** Staging ,Restaging & therapy assessment of HD , NHL , Lung cancer

» Specific tumor imaging agents:

- **In-111 (TC99m) Octreotide :** Neuroendocrine tumors
- **I -123 MIBG :** Neuroendocrine tumor
- **I -131 :** Lung mets. thyroid carcinoma

» Bone Scan In Oncology

■ Procedure

Radiopharmaceuticals:

- Technitium 99m Methylene DiPhosPhonate (Tc-99m MDP) has High sensitivity but none specific

Tissue accumulation depends on:

- Blood flow
- Capillary permeability
- Metabolic activity of osteoblasts and osteoclasts
- Mineral turnover

Dose: 500 to 800 MBq (Megabequerel) / 15- 20 mCi (millicurie)

Imaging time: 2 to 3 hours postinjection - WB + SPECT

Potentials of bone scan: Positivity many months before an abnormality can be detected on X ray

■ Indications

1 Metastatic disease: (Lung cancer, prostate, breast, thyroid, and renal tumours)

- Diagnosis.
- Initial staging.
- Restaging.
- Asses response to therapy.

2 Primary Bone Tumor: Main diagnosis is done by radiology

- Malignant or Benign
- Therapy planning for patients with primary bone malignancy (e.g. Osteogenic & Ewings sarcoma)

3 Soft tissue tumors :

- Primary
- Metastases

Imaging features

- A. Hot lesions** : Majority of bone tumors. Doesn't show the metastasis itself but shows the osteopathic reaction around the tumor
- B. Cold lesions** : Purely osteolytic tumors (renal cell carcinoma, thyroid cancer, anaplastic tumors), radiation therapy
- C. Superscan** : Diffuse increased skeletal uptake with no soft tissue or kidney activity (e.g. CA prostate, breast, ...etc).
- D. Normal distribution** : Marrow tumors (e.g. lymphomas, leukemia, multiple myeloma).
- E. Soft tissue uptake** : Soft tissue tumors may concentrate the tracer
- F. Flare phenomenon**: increased number of lesions in the case of effective therapy Healing process, indicate good response to treatment

» Bone Scan : In Metastatic Disease

Access of Non Osseous Tumors To Bone:

- Direct Extension.
- Retrograde venous flow.
- Arterial Circulation (after venous or lymphatic access).

Epithelial Tumors:

- Reach red marrow of axial skeleton via venous and arterial flow.
- ★ **Distribution of red marrow in adult** : calvarium, spine, pelvis, and proximal femurs and humerus
- 90% of metastasis from epithelial tumors are found in **red marrow**. In adults the red marrow is confined in the axial skeleton, therefore there is no red marrow below the knee and elbow. The metastasis is only in the axial skeleton. With some exception in lung CA where there is metastasis to the big toe and nobody can explain this.

Metastatic Foci:


- Grows in red marrow space.
- The surrounding bone remodels through osteoclastic (resorption) and osteoblastic (deposition) activity.
- The relative balance between resorption and deposition determine whether the lesion is hot (sclerotic), cold (lytic) or mixed pattern.
- The tracer does not concentrate in the metastatic foci (cancerous tissue) but in the surrounding reactive bone.

Sensitivity:

- In early stage superior to x-ray. Functional changes are faster than structural changes
- In advanced stage both have high sensitivity.
- The accuracy of bone scan not known because of the lack of reference standard.
- The sensitivity is agreed to be 90% or more.


Scan Patterns:

1. Solitary lesions.
2. Multiple focal lesions.
3. Diffuse involvement (Superscan).
4. Photon deficient lesions (cold lesions).
5. Flare phenomenon.
6. Normal (false negative).
7. Soft tissue lesions (tracer uptake in tumor).



Tumor Imaging

Tumor staging



CA breast
hot spots represents
bone metastasis

CA lung

CA stomach

» Diffuse involvement (Superscan) ★

Definition : Bone scan with diffuse symmetrical increased uptake and almost absence of soft tissue activity, lack of kidney activity and bone uptake seen in blood pool images.

Causes :

- a. **Bone metastases :** Prostate, breast, lung, bladder and lymphoma.
- b. **Non tumor causes:** HPT, osteomalacia, Paget's disease and fibrous dysplasia


Important clues : In metabolic bone disease the calvarium and long bones are involved unlike in bone metastases.

Superscan



CA prostate.

Diffuse increased uptake in the **axial skeleton**, nothing below knee or elbow



HPT

Diffuse increased uptake in the mandible and below knee and elbow

Tumor Imaging


Pure Lytic Lesions Caused by multiple myeloma, radiation, or by past surgery like this patient [tumor was removed(arrow)]



Bone Scan : Radiation Effect



Patient with bone metastasis







Follow up bone scan, shows cold areas, good response to radiation

- **A: Hours** following radiation : **Increased** uptake due to increased blood flow and vascular permeability.
- **B: 3-6 months** post radiation : **Decreased** uptake due to microvascular injury. Dose related (>2000 rads).
- **Following Radiotherapy** : Spontaneous ribs fracture may occur.

Tumor Imaging

Ewing's Sarcoma






- A= Increased vascularity of tumor
- B= Metabolic delayed phase, increased uptake
- C= No distant metastasis

» Bone Scan In Bone Tumors




Osteoid Osteoma

(Benign bone tumor. Usually affects **children**, present as **pain** in the site of the tumor relieved by aspirin)







- A=Focal area of increased blood flow
- B=Delayed scan shows nodes of intense uptake

Giant Cell Tumor



Soft Tissue Sarcoma

(Goal is to see is there the local extent of the tumor and is there any distant metastasis)



» Properties: (Introduced in seventies of 20th century for lymphomas)

1) Mechanisms of accumulation:

- Tumour viability
- Blood flow
- Capillary permeability
- Lymphatic drainage
- Binds to Transferrin receptors on the tumour cells

2) Non specific for infection-inflammation and tumors

3) Excretion: Kidneys and large bowel

» Imaging Protocol:

- **Patient preparation:** Laxatives for bowel preparation post injection, nothing else
- **Several weeks post tumor therapy (FN)** Radiation therapy and chemotherapy can alter the normal pattern of gallium distribution
- **180 MBq (4-5 mCi) is usually administered** Imaging follows after 48 - 72 hours WB + SPECT/SPECT CT, medium-energy collimator


» Normal Ga-67 scan

Normal scan

Accumulates in bone marrow and liver. Splenic uptake is variable. The kidneys are usually visualized and also lacrimal, salivary, nasopharyngeal and genital activity is often present. Female breasts can be visualized, but accumulation is physiologically symmetrical. Radioactivity is commonly seen in the colon.

Clinical indications


- Lymphoma
- Melanoma
- Lung cancer
- Hepatoma



» Gallium Scan in Lymphoma

- ✓ Staging
- ✓ Follow up and monitoring of therapy
- ✓ Detection of tumor recurrence
- ✓ Differentiate post therapy changes : tissue necrosis and fibrosis from local recurrence.

Affected lymph nodes above and below the diaphragm




» Gallium Scan in Lymphoma (Continue..)

Prediction of response to therapy

07.1




Before chemotherapy



After chemotherapy

Normalization of a positive pre-therapy scan : A negative scan after one cycle

Prediction of outcome




Partial response


Residual gallium uptake after treatment is a poor prognostic sign, indicates viable tumor and treatment should be modified.

» Gallium Scan NHL

Planar Vs SPECT CT



Lymph nodes in the mediastinum



In-111 octreoscan.



I123 MIBG Scan.

» Somatostatin Receptor Imaging (Indium-111 Octreoscan)

Insulinoma

Clinical History The patient is a 66-ys male with insulinoma, now being evaluated for evidence of recurrent and/or metastatic disease.

Findings : Multiple lung, mediastinum , liver and abdominal metastases.



Labeling: Liver, Spleen, Kidneys, Urinary bladder, Gallbladders

NORMAL STUDY

» I123 MIBG Scan (MIBG : Meta Iodo Benzyl Guanidine)



- Is a nor adrenaline analog
- Localizes in adrenergic tissues: catecholamines producing tumors and their metastases.
- Patient preparation: stop drugs interfering with MIBG uptake. Lugols solution to protect thyroid gland

Indications: Neuroendocrine tumors

(Pheochromocytoma, Paraganglioma, Insulinoma, Neuroblastoma ,Medullary thyroid carcinoma Carcinoid tumors)

■ MIBG In Pheochromocytoma


Bilateral Disease




A: large left adrenal tumor
B: smaller on the right

MIBG In Pheochromocytoma (Continue..)

Clinical History A 41 years old female patient with 2ndary hypertension. Right adrenal mass.? **Pheochromocytoma.**



Tumor is paraspinal, supranational.




Planar Vs SPECT CT


MIBG In Neuroblastoma (Wilms' tumour)

(effects children under 5 years, can be removed)

Planar Vs SPECT CT



Abnormal uptake in the right upper abdomen



Intense MIBG uptake, in the right paraspinal and subhepatic (arrow)


1131 MIBG Total body scan

1ry neuroblastoma /bone mets

1131 MIBG:when we label MIBG with 1131 we can use it to treat,so MIBG is used in both **treatment** and **diagnosis**

A: Primary

B: Metastasis treated with 1131 MIBG



Thyroid Metastases Study

(I-123 or I-131 as Sodium Iodide)

(I-123 is the gold standard for **imaging** while I-131 is used for **therapy**)

Indications

Detection and localization of persistent or recurrent local or distant functioning thyroid cancer

Patient Preparation

1) Stimulation of potentially functioning thyroid tissue:

A. Inject **recombinant human thyrotropin** on 2 consecutive days and administer the iodine on the third day .

B. Withdraw thyroid replacement hormones :

1. Thyroxine (T-4) for at least 4 weeks.

2. Triiodothyronine (T-3) for at least 10 days.

2) The patient must not have had i.v iodinated contrast material (IVP, CT with contrast, myelogram, angiogram) for at least 3 weeks .

3) The patient should be NPO for at least 4 hours prior to radiopharmaceutical administration and for at least 1 hour afterwards .

Tracer , Dose, & Technique of Administration

Radiopharmaceutical: **Oral** administration


a. I-123 as sodium iodide : 2 mCi

b. I-131 as sodium iodide : 2-10 mCi


Imaging using Gamma camera : Whole body scan

» Thyroid Cancer

I-123 WB Scan



Negative I-123 WB




I-123 WB Scan : Postoperative Thyroid remnants


» Thyroid Cancer (Continue...)

I-123 WB Scan: Postoperative Thyroid remnants


Planar Vs SPECT CT




Localized retrosternally




» Thyroid metastases study: (I-123 or I-131 as Sodium Iodide)



Local recurrence




Bone metastases




Lung metastases

I-131 Pre & Post therapy



Dec 04




(Resolution) March 06

What is PET - CT?

PET: Positron Emission Tomography.

CT: Computerized Tomography.

PET-CT is the fusion of functional and anatomic information acquired almost simultaneously from which we are able to visualize form and function.




PET CT= PET + CT = Function + Form

PET : How it is performed...?

Positron emitters (e.g. F18) labelled with biologically active natural compounds such as oxygen, carbon or glucose given intravenously and reacting in the body identically to their non-radioactive counterparts.

Positrons are emitted from F18 and react with tissue electrons....Anihilation occurs...

Two photons 511 kev each in opposite direction are emitted and detected by PET SCANNER giving an image of the normal and abnormal distribution of tracer in the body.



» Positron Emitting Isotopes:

1.Cyclotron produced isotopes:

Isotope	T/2
Oxygen-15	2 min
Nitrogen-13	10 min
Carbon-11	20 min
Fluorine-18	110 min

2.Generator produced isotopes:

Parent	T/2	daughter	T/2
Strontium -82	25 days	Rubidium-82	75 seconds
Zin-62	9.3 hrs	Copper-62	10 min
Germanium m-68	288 days	Gallium-68	68 min

*The ones in the red are the most used agents. Important to know them


» FDG : Fluoro-2-deoxy-D-Glucose

Uptake mechanism:

FDG is a glucose analogue used to assess glucose metabolism. FDG gets transported from intravascular space to the cells by the same mechanism as the glucose. In the cell, hexoKinase acts on both **FDG and glucose to form:**

- FDG-6-phosphatase (FDG-6-PO₄-)
- Glucose-6-phosphatase.

FDG-6-PO₄- cannot progress further into glucose metabolism and remains trapped intracellularly in proportion to glycolytic rate of the cell.



In tumors, there is high rate of glycolysis compared to normal cells as well as higher level of hexoKinase. FDG is labeled with F18

» Normal distribution

In a non fasting state

Brain & heart	High uptake
Liver	Less uptake
Kidneys	Unlike glucose, FDG is excreted with urine
GI	Mild clearance, faintly seen
Muscles	Low, increase with exercise



» FDG in oncology

Tumors do not have a blood tumor barrier, FDG transport into tumors occurs at a higher rate than in the surrounding normal tissues. FDG is de-phosphorylated and can then leave the cell. The de-phosphorylation occurs at a slower rate in tumors.

» Applications of FDG:

- Locating unknown primaries
- Differentiation of tumor from normal tissue
- Pre-operative staging of disease
 - (lung, breast, colorectal, melanoma, H&N, pancreas)
- Recurrence vs necrosis
- Recurrence vs post-operative changes (limitations with FDG)
- Monitoring response to therapy

➤ PET CT F18 FDG imaging protocol


Fasting : 4 - 6 hours

Dose : Inject 10 mCi F18 FDG

Wait (uptake phase): 45 -60 min then scan

Scanning time : 30 min to complete PET CT study


SUV : Standard uptake value (N:0.5-2.5 and Tumors > 3.0)



➤ FDG PET




FDG PET
Normal




FDG PET
Staging of NHL
(stage 4 lymphoma)

Bilateral
mediastinum and
hilarious lymph
nodes para aortic
lymph nodes in the
abdomen

➤ FDG PET CT




FDG PET-CT
Normal



FDG PET-CT
Staging Of Lymphoma

⇒ FDG PRT CT NHL: stage IV



- Extensive lymph nodes involvement
- Spleen involvement
- Para aortic involvement



wow, such empty

Hodgkin's lymphoma


FDG PETCT : Hodgkin's lymphoma

Post 6 cycles of chemotherapy for assessment.



Baseline

(24.10.2018) = Lymph nodes with spleen involvement



Interim

(1.1.2019) = Few remnants



Final

(19.5.2019) = Complete Resolution

Partial metabolic response (Deauville Score 5).



Baseline

(31.3.2019)



Interim

(9.6.2019) = Partial response




Final

(4.8.2019) = Not completely resolved


Progressive disease (Deauville Score 5)

Baseline
(19.2.2019)




Interim
(17.7.2019)
= Got worse

FDG PETCT Rectal cancer with liver metastases




FDG PET CT : Nasopharyngeal cancer with LN metastasis





FDG PET CT : RCC with bone and lung metastases

309756 FDG-avid left renal mass coupled with FDG-avid locoregional nodal disease and multiple FDG avid pulmonary metastasis. Solitary sclerotic area of hypermetabolism at the right iliac bone, worrisome for bone metastasis.





» FDG PET CT : Metastatic Lung Cancer

Multiple distant metastasis in the lung, liver, cervical and bone



» FDG PET – brain tumor

post therapy Two foci on CT, only one viable tumor



Indications of PET CT

Breast Cancer*	Staging*, restaging*, and monitoring response to therapy*
Colorectal Cancer	Diagnosis*, staging* and restaging*
Esophageal Cancer	Diagnosis*, staging* and restaging*
Head & Neck Cancers (excluding CNS and thyroid)	Diagnosis*, staging* and restaging*
Lung Cancer (Non-Small Cell)	Diagnosis*, staging* and restaging*
Lymphoma	Diagnosis*, staging* and restaging*
Melanoma (Excludes evaluation of regional nodes)	Diagnosis*, staging* and restaging*
Solitary Pulmonary Nodule	Characterization of indeterminate single pulmonary nodule
Thyroid Cancer*	Restaging
Cervical Cancer*	Staging as an adjunct to conventional imaging

» Somatostatin receptor PET tracers Ga-68 DOTANOC

(Gold standard for somatostatin receptor)

Radiopharmaceutical: DOTANOC, DOTATOC or DOTATATE is labeled with Ga-68

Dose: 3-5 mCi given intravenously.

PET Imaging time: 45-60 min postinjection

Clinical value: higher lesion detection rate than is achieved with (18)F-fluorodihydroxyphenyl-L-alanine PET, somatostatin receptor SPECT, CT, or MR imaging.

Sensitivity: 70-100% (depends on density of somatostatin receptors in the tumor)

Indications:

Tumours with high expression of receptors of somatostatin


- ❖ Gastroenteropancreatic tumours
 - e.g. carcinoids, gastrinoma, insulinoma, glucagonoma, VIPoma, etc.
- ❖ Sympathoadrenal system tumours
 - pheochromocytoma, paraganglioma, neuroblastoma, ganglioneuroma
- ❖ Medullary thyroid carcinoma
- ❖ Pituitary adenoma
- ❖ Medulloblastoma
- ❖ Merkel cell carcinoma
- ❖ Small-cell lung cancer
 - mainly primary tumours
- ❖ Meningioma

» Normal distribution 68Ga-DOTA peptide PET/CT




- Normal tracer uptake is seen in the pituitary, salivary glands, thyroid, liver, spleen, adrenals, pancreas, kidneys, ureters, and bladder

» 68Ga-DOTA PET



- NET with multiple metastatic disease confined to the liver and abdominal cavity



- NET with extensive metastatic lesions throughout the body

» Ga-68 DOTANOC PET superior to In-111 Octreoscan


» Carcinoid tumor :

Positive ^{68}Ga -DOTA-NOC and
Negative ^{111}In -Octreoscan.

» ^{68}Ga DOTA-NOC Findings:


Multiple metastatic lesions in the liver.
(The pituitary also expresses somatostatin receptors and is visualized in the ^{68}Ga PET image, along with normal uptake in the spleen, kidneys, and bladder.)

» Indication of ^{68}Ga DOTA-NOC: The ^{68}Ga PET scan was performed because the patient's symptoms were inconsistent with the ^{111}In -Octreoscan findings.




» Pheochromocytoma

Ga68 DOTANOC superior to MIBG



SPECT CT - I123 MIBG



PET CT Ga68 DOTANOC
(Higher sensitivity)

» Radionuclide Therapy

Properties of the Ideal Therapeutic Radiopharmaceutical


1. **Pure beta minus emitter.** =(All I want you to know from this section)
2. Medium/high energy (>1 meV).
3. Effective half-life = moderately long, e.g., days.
4. High target:non-target ratio
5. Minimal radiation dose to patient and Nuclear Medicine personnel
6. Patient Safety
7. Inexpensive, readily available radiopharmaceutical.
8. Simple preparation and quality control if manufactured in house.

» Radionuclide Therapy (cont)

Dose not important :)

Agent	Indication	Dose
★ I131 :	Thyroid cancer	100-200 mCi
¹³¹ MIBG :	Neuroblastoma	100-300 mCi
Strontium-89	Bone metastasis	40-60uCi/kg
Sm-153-EDTMP:	Bone metastasis	1.0 mCi per kg
Phosphorus- 32 :	Polycythaemia	2.3 mCi/m ²
Y-90-Ibritumomab Tiuxetan [Zevalin®]:	B-Cell NHL	0.3 mCi/kg
> Platelet count > 150,000 cells/mL: 0.4 mCi/kg > Platelet count 100,000-150,000 cells/mL: The dose should never exceed 32 mCi (1,184 MBq)		

» Strontium-89 Therapy for Palliation of Bony Metastases



History :

A 65 Ys ,M, with CA prostate and widespread bone metastases and severe bony pain.

Admitted for palliative Strontium-89 therapy.

Procedure :

- Bone metastases was confirmed by bone scan. The patient was given 40 mCi of Strontium-89 I.V. according to body weight of the patient. The patient experienced one day of exacerbated pain which was controlled by opiates but the following day showed gradual pain relieve.


» NM tumor imaging

- Functional
- Sensitive
- Whole body evaluation
- Specific : Some tumors
- Targeted therapy

» objectives of NM tumor imaging


- Diagnosis
- Staging
- Guiding biopsy
- Follow up and therapy monitoring
- Detection of recurrence.

tumor non- specific diagnostic radiopharmaceuticals	tumor specific diagnostic radiopharmaceuticals	therapeutic radiopharmaceuticals
<ul style="list-style-type: none"> • F-18 FDG • Diphosphonates • ga-67 citrate • tc99m nanocolloid • Tc99m NIBI/ thallium 201 	<ul style="list-style-type: none"> • (Ga-68 DOTA) <ul style="list-style-type: none"> • (F-18-FLT) • (F-18-FMISO) 	<ul style="list-style-type: none"> • non- specific: <ul style="list-style-type: none"> ○ Sr-89, Sm-153, Re-189 • specific <ul style="list-style-type: none"> ○ I-131 ○ Y-90
Demonstrate tumor sites but are not specific for malignancy	Binds directly to special tumor antigens or receptors or are accumulated by special metabolic pathway	



1-this is the image is a result of a 5 month treatment done to a patient with hodgkin's lymphoma. it shows:

- a. complete resolution
- b. partial resolution
- c. it got worse
- d. i don't know



2-in the normal distribution of FDG which of the following represents the uptake from highest to lowest?

- a. kidneys> brain> heart> liver
- b. brain> heart> liver> GI
- c. liver> kidneys> brain> heart
- d. brain>liver> GI> heart

3-neuroblastoma metastasis is treated with


- a. Sr-89
- b. Y-90
- c. I-131
- d. Re-189

4-which one is the Gold standard for tumour imaging

- a. Gallium 67
- b. Thallium 201
- c. F18 - FDG
- d. Tc-99m MDP

5-A 55 year old male came with insulinoma and an Indium-111 Octreoscan was done and showed metastasis to the

- a. Liver
- b. Kidney
- c. Gallbladder
- d. Urinary bladder



6-Fluorine-18 has a T/2 of?

- a. 20m
- b. 100m
- c. 110m
- d. 120m

Answers
1)b
2)b
3)c
4)c
5)a
6)c